

10/826,439

EAST Search History

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	0	("2004006104").PN.	US-PGPUB	OR	OFF	2006/05/11 07:14
L2	1	("20040006104").PN.	US-PGPUB	OR	OFF	2006/05/11 07:19
L3	1	("6482949").PN.	US-PGPUB; USPAT	OR	OFF	2006/05/11 08:30
L4	5	"2005007099"	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	OFF	2006/05/11 08:30
S1	985	544/344 OR 544/347 OR 544/353	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	OFF	2006/05/11 07:13
S2	72	S1 AND (ANTIVIRAL OR HCV OR HEPATITIS)	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	ON	2006/05/10 11:58
S3	1	("6989451").PN.	USPAT	OR	OFF	2006/05/10 12:31
S4	1	("3510487").PN.	USPAT	OR	OFF	2006/05/10 12:42
S5	1	("3656953").PN.	USPAT	OR	OFF	2006/05/10 13:29
S6	1	("6518423").PN.	USPAT	OR	OFF	2006/05/10 13:39
S7	1	("6103720").PN.	USPAT	OR	OFF	2006/05/10 13:43
S8	1	("5874587").PN.	USPAT	OR	OFF	2006/05/10 13:45
S9	1	("5969150").PN.	USPAT	OR	OFF	2006/05/10 13:45

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPAT1623ZCT

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR 7):2

***** Welcome to STN International *****

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 DEC 23 New IPC8 SEARCH, DISPLAY, and SELECT fields in USPATFULL/
USPAT2
NEWS 4 JAN 13 IPC 8 searching in IPAT, IPIUDS, and IPIUDB
NEWS 5 JAN 13 New IPC 8 SEARCH, DISPLAY, and SELECT enhancements added to
INFADOC
NEWS 6 JAN 17 Pre-1988 INPI data added to MARRPAT
NEWS 7 JAN 17 IPC 8 in the MPI family of databases including WPIFV
NEWS 8 JAN 30 Saved answer limit increased
NEWS 9 FEB 21 STN AnaVist, Version 1.1, lets you share your STN AnaVist
Visualization results
NEWS 10 FEB 22 The IPC thesaurus added to additional patent databases on STN
NEWS 11 FEB 22 Updates in EPFULL; IPC 8 enhancements added
NEWS 12 FEB 27 New STN AnaVist pricing effective March 1, 2006
NEWS 13 FEB 28 MEDLINE/LMEDLINE reload improves functionality
NEWS 14 FEB 28 TOXCENTER reloaded with enhancements
NEWS 15 FEB 28 REGISTRY/REGISTRY enhanced with more experimental spectral
property data
NEWS 16 MAR 01 INSPEC reloaded and enhanced
NEWS 17 MAR 03 Updates in PATDPA; addition of IPC 8 data without attributes
NEWS 18 MAR 08 X.25 communication option no longer available after June 2006
NEWS 19 MAR 22 EMBASE is now updated on a daily basis
NEWS 20 APR 03 New IPC 8 fields and IPC thesaurus added to PATDPAFULL
NEWS 21 APR 03 Bibliographic data updates resume; new IPC 8 fields and IPC
thesaurus added in PCTFULL
NEWS 22 APR 04 STN AnaVist \$500 visualization usage credit offered
NEWS 23 APR 12 LINSPEC, learning database for INSPEC, reloaded and enhanced
NEWS 24 APR 12 Improved structure highlighting in PQHIT and QHIT display
in MARRPAT
NEWS 25 APR 12 Derwent World Patents Index to be reloaded and enhanced during
second quarter; strategies may be affected
NEWS 26 MAY 10 CA/Cplus enhanced with 1900-1906 U.S. patent records
NEWS EXPRESS FEBRUARY IS CURRENT VERSION FOR WINDOWS IS V8.01a,
CURRENT MACINTOSH VERSION IS V6.0c (ENG) AND V6.0c (JP),
AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005.
V8.0 AND V6.01 USERS CAN OBTAIN THE UPGRADE TO V8.01a AT
<http://download.cas.org/express/v8.0-Discover/>
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific


on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

>>Testing the current file.... screen

ENTER SCREEN EXPRESSION OR (END):end

>> Uploading C:\Program Files\Stnexp\Queries\ANRONGUICHOW ANTIVIRALS.str



chain nodes :
11 13 15 16 18 19
ring nodes :
1 2 3 4 5 6 7 8 9 10
ring/chain nodes :
20
chain bonds :
5-11 11-13 11-15 15-16 16-18 16-19 18-20
ring bonds :
1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10
exact/norm bonds :
11-13 11-15 15-16
exact bonds :
5-11 16-18 16-19 18-20
normalized bonds :
1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10

G1:C,S

G2:O,N

G3:C,H,S,P

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:CLASS 13:CLASS 15:CLASS 16:CLASS 18:CLASS 19:CLASS 20:CLASS

L1 STRUCTURE UPLOADED

>> que L1

L2 QUE L1

>> D L1

L1 HAS NO ANSWERS

L1 STR

research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

COMPLETE THE STN SURVEY - APRIL 27 THROUGH MAY 31

Dear valued STN customer,

In an effort to enhance your experience with STN, we would
like to better understand what you find useful. Please take
approximately 5 minutes to complete a web survey.

If you provide us with your name, login ID, and e-mail address, you
will be entered in a drawing to win a free iPod(R). Your responses
will be kept confidential and will help us make future improvements
to STN.

Take survey: <http://www.zoomerang.com/survey.xgi?p=WEB2259HONKUNTU>

Thank you in advance for your participation.

***** STN Columbus *****

FILE 'HOME' ENTERED AT 12:13:09 ON 10 MAY 2006

FILE REG	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 12:13:15 ON 10 MAY 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 9 MAY 2006 HIGHEST RN 883631-57-0
DICTIONARY FILE UPDATES: 9 MAY 2006 HIGHEST RN 883631-57-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.


TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *

Structure search iteration limits have been increased. See HELP SLIMITS
for details.

REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document. For information



THIS IS
FOR CL-1, 2, 3, 4, 6

G1 C,S

G2 O,N

G3 [G1],[G2]

Structure attributes must be viewed using STN Express query preparation.

>> S L1

SAMPLE SEARCH INITIATED 12:13:34 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 530 TO ITERATE

100.0% PROCESSED 530 ITERATIONS 12 ANSWERS
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 9219 TO 11981
PROJECTED ANSWERS: 33 TO 447

L3 12 SEA SSS SAM L1

>> S L1 SSS FULL

FULL SEARCH INITIATED 12:13:40 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 10767 TO ITERATE

100.0% PROCESSED 10767 ITERATIONS 329 ANSWERS
SEARCH TIME: 00.00.01

L4 329 SEA SSS FUL L1


>>Testing the current file.... screen

ENTER SCREEN EXPRESSION OR (END):end

>> Uploading C:\Program Files\Stnexp\Queries\ANRONGUICHOW ANTIVIRALS.str

STN SEARCH TRANSCRIPT

10/826,439



```

chain nodes :
11 13 15 16 18 19
ring nodes :
1 2 3 4 5 6 7 8 9 10
ring/chain nodes :
20
chain bonds :
5-11 11-13 11-15 15-16 16-18 16-19 18-20
ring bonds :
1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10
exact/norm bonds :
11-13 11-15 15-16
exact bonds :
5-11 16-18 16-19 18-20
normalized bonds :
1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10

```

G1:C,S

G2:O,N

G3:C,H,S,P

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:CLASS 13:CLASS 15:CLASS 16:CLASS 18:CLASS 19:CLASS 20:CLASS

L5 STRUCTURE UPLOADED


=> que L5

L6 QUE L5

=> D L5

L5 HAS NO ANSWERS

L5 STR



G1:C,S

G2:O,N

G3:C,H,S,P

Structure attributes must be viewed using STN Express query preparation.

=> S L5
SAMPLE SEARCH INITIATED 12:16:14 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 530 TO ITERATE

100.0% PROCESSED 530 ITERATIONS 50 ANSWERS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 9219 TO 11981
PROJECTED ANSWERS: 833 TO 1807

L7 50 SEA SSS SAM L5

=> S L5 SSS FULL
FULL SEARCH INITIATED 12:16:19 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 10767 TO ITERATE

100.0% PROCESSED 10767 ITERATIONS 1540 ANSWERS
SEARCH TIME: 00.00.01

L8 1540 SEA SSS FUL L5


=> S L8 NOT L4

L9 1213 L8 NOT L4

=>Testing the current file.... screen

ENTER SCREEN EXPRESSION OR (END):end

=> Uploading C:\Program Files\Stnexp\Queries\ANRONGWUCHOW ANTIVIRALS.etr



```

chain nodes :
11 13 15 16 18 19
ring nodes :
1 2 3 4 5 6 7 8 9 10
ring/chain nodes :
20
chain bonds :
5-11 11-13 11-15 15-16 16-18 16-19 18-20
ring bonds :
1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10
exact/norm bonds :
11-13 11-15 15-16
exact bonds :
5-11 16-18 16-19 18-20
normalized bonds :
1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10

```

G1:C,S

G2:O,N

G3:C,H,S,P

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:CLASS 13:CLASS 15:CLASS 16:CLASS 18:CLASS 19:CLASS 20:CLASS

L10 STRUCTURE UPLOADED


=> que L10

L11 QUE L10

=> D L10

L10 HAS NO ANSWERS

L10 STR



G1:C,S

G2:O,N

G3:C,H,S,P

Structure attributes must be viewed using STN Express query preparation.

=> S L10 SUB=L9 FULL
FULL SUBSET SEARCH INITIATED 12:21:05 FILE 'REGISTRY'
FULL SUBSET SCREEN SEARCH COMPLETED - 987 TO ITERATE

100.0% PROCESSED 987 ITERATIONS 790 ANSWERS
SEARCH TIME: 00.00.01

L12 790 SEA SUB=L9 SSS FUL L10

FILE CAPLUS

COST IN U.S. DOLLARS

FULL ESTIMATED COST

SINCE FILE ENTRY TOTAL
379.00 379.21

FILE 'CAPLUS' ENTERED AT 12:22:02 ON 10 MAY 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 10 May 2006 VOL 144 ISS 20
FILE LAST UPDATED: 9 May 2006 (20060509/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> S L4 OR L12
40 L4
151 L12
L13 181 L4 OR L12

← ALL SEARCHES TOGETHER
DISPLAYED 1-181

L13 ANSWER 1 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2006:361356 CAPLUS
TITLE: Preparation of disulfide dyes for dyeing human keratin fibers
INVENTOR(S): Daubresse, Nicolas; Genain, Gilles
PATENT ASSIGNEE(S): Fr.
SOURCE: U.S. Pat. Appl. Publ., 39 pp.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2006080791	A1	20060420	US 2005-249357	20051014
FR 2876574	A1	20060421	FR 2004-10864	20041014
EP 1647580	A1	20060419	EP 2005-292159	20051013
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IR, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, BA, HR, IS, YU				
JP 2006111624	A2	20060427	JP 2005-299150	20051013
PRIORITY APPL. INFO.: FR 2004-10864 A 20041014 US 2004-629308P P 20041119				


AB Disclosed herein is a dyeing composition comprising a particular disulfide dye and a method of dyeing human keratin fibers, such as hair, using this composition. This composition makes it possible to obtain particularly fast chromatic colorations. E.g., I was prepared from cystamine-2HCl and Reactive Blue 44. I and other prepared dyes were tested on gray hair.

IT INDEXING IN PROGRESS

IT 883566-67-4P
RL: BSU (Biological study, unclassified); COS (Cosmetic use); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)
(Preparation of disulfide dyes for dyeing human keratin fibers)


RN 883566-67-4 CAPLUS

CN INDEX NAME NOT YET ASSIGNED



PAGE 1-A

4 Na




AB This invention relates to novel compds. useful in the treatment of diseases associated with TRPV4 channel receptor. E.g., I was prepared from 2-O-DAB (BOC)-OH DCHA and ClCO2Et giving an intermediate which was treated with phthalimide and Ph3P and DEAD giving phenylmethyl (2R)-4-[[1,1-dimethylethoxy]carbonyl]amino]-2-[[1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)methyl]butanoate which was treated with HCl and N-(1-benzothien-2-ylcarbonyl)-L-leucine. The resulting intermediate was brominated and treated with 2-chloro-4-fluorobenzenesulfonyl chloride and the resulting intermediate hydrazinolized to give I. Tablets were prepared containing I.

IT 878797-73-0P
RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(Preparation of acyclic 1,3-diamines for use in treatment of diseases associated with TRPV4 channel receptor)

RN 878797-73-0 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[(1S)-1-[[[3-[[[2-chloro-4-fluorophenyl]sulfonyl]amino]propyl]amino]carbonyl]-3-methylbutyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L13 ANSWER 3 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2006:162602 CAPLUS
TITLE: Nucleic acid intercalators and avidin probes derived from luminescent cyclometalated iridium(III)-dipyridoquinoline and -dipyridophenazine complexes
AUTHOR(S): Lo, Kenneth Kam-Wing; Chung, Chi-Kuang; Zhu, Nanyong
CORPORATE SOURCE: Department of Biology and Chemistry, City University of Hong Kong, Kowloon, Hong Kong, Peop. Rep. China
SOURCE: Chemistry--A European Journal (2006), 12(5), 1500-1512
CODEN: CHEUJD; ISSN: 0947-6539
PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA
DOCUMENT TYPE: Journal
LANGUAGE: English
OI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-B



L13 ANSWER 2 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2006:240647 CAPLUS
DOCUMENT NUMBER: 144:311900
TITLE: Preparation of acyclic 1,3-diamines for use in treatment of diseases associated with TRPV4 channel receptor.
INVENTOR(S): Casillas, Linda N.; Jeong, Jae Uk; Marquis, Robert W.
PATENT ASSIGNEE(S): SmithKline Beecham Corporation, USA
SOURCE: PCT Int. Appl., 152 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006029210	A2	20060316	WO 2005-US31873	20050907
M: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, OM, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
PRIORITY APPL. INFO.: US 2004-607678P P 20040907				

OI

AB Six luminescent cyclometalated cationic redox-active luminescent 2-phenylpyridine iridium(III)-complexes with substituted pyrazino-annulated phenanthroline bidentate ligands were prepared; the DNA and avidin intercalation were assayed by emission titration. Reaction of [Ir2(ppp)4Cl2] (ppp = 2-phenylpyridine) with ligands L2 gave complexes 1-PP6 (1-3; L2 = dpq, R = H; L2 = dpq, R = CONH2; L = dpq, R = CONH(CH2)2NHQ, where Q = CO(CH2)4CSH7N2OS, biotinyl) and II [4-6; L2 = dpq, R1 = R2 = H; L2 = dpq, R1-R2 = benzo; L2 = dpq, R2 = H, R1 = CONH(CH2)2NHQ] were designed as luminescent intercalators for DNA and probes for avidin. The crystal structure of complex 4 is reported. The photophysics and electrochem. properties of the complexes 1-4 were also investigated. The binding of these complexes to double-stranded calf thymus DNA and synthetic double-stranded oligonucleotides poly(dA)-poly(dT) and poly(dG)-poly(dC) was investigated by spectroscopic titrations. The interactions between the two biotin-containing complexes 3 and 6 and avidin were studied by 4'-hydroxybenzoic acid (HABA) assays and emission titrations.


IT 882571-97-3P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(Preparation of cyclometalated luminescent iridium pyrazino-annulated phenanthroline 2-phenylpyridine complexes as DNA intercalators and avidin complexants)

RN 882571-97-3 CAPLUS


CN Quinoxalino[2,3-f][1,10]phenanthroline-11-carboxamide, N-[2-[[[5-[[[3aS,4S,6aR]-hexahydro-2-oxo-1H-thieno[3,4-d]imidazol-4-yl]-1-oxopentyl]amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



IT 882571-95-1P
RL: BSU (Biological study, unclassified); CPS (Chemical process); PREP (Physical, engineering or chemical process); PYP (Physical process); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); PROC (Process)
(redox potential, luminescence spectra; preparation of cyclometalated luminescent iridium pyrazino-annulated phenanthroline 2-phenylpyridine)

complexes as DNA intercalators and avidin complexants)
RN 882571-95-1 CAPLUS
CN INDEX NAME NOT YET ASSIGNED

PAGE 2-A



CM 1

CRN 882571-94-0

CMF C53 H46 Ir N10 O3 S

CCI CCS

PAGE 1-A




CM 2

CRN 16919-18-9

CMF F6 P

CCI CCS



REFERENCE COUNT: 115 THERE ARE 115 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 4 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STM
ACCESSION NUMBER: 2006:152549 CAPLUS
DOCUMENT NUMBER: 144:232928
TITLE: Preparation of heterocyclic compounds as novel antimalaria agents
INVENTOR(S): Nakamoto, Kazutaka; Matsukura, Masayuki; Tanaka, Keigo; Inoue, Satoshi; Tawakada, Itaru; Haneda, Toru; Ueda, Morihiro; Abe, Shinya; Sagane, Koji
PATENT ASSIGNER(S): Eisai Co., Ltd., Japan
SOURCE: PCT Int. Appl., 326 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:


PATENT NO. KIND DATE APPLICATION NO. DATE
WO 2006016548 A1 20060216 WO 2005-JP14505 20050808
W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HD, IL, IN, IS, JP, KE, KG, KM, KN, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MG, MH, MI, MN, MO, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CP, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TO, BW, GH, GM, KS, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
WO 2005033079 A1 20050414 WO 2004-JP14063 20040927
W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HD, IL, IN, IS, JP, KE, KG, KM, KN, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MG, MH, MI, MN, MO, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CP, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TO

PRIORITY APPL. INFO.: JP 2004-232617 A 20040809

WO 2004-JP14063 A 20040927
JP 2005-82760 A 20050322
JP 2003-342273 A 20030930
JP 2004-68186 A 20040310

OTHER SOURCE(S): MARPAT 144:232928

GI



AB Antimalaria agents containing comds. represented by the formula (I) (wherein A1 = each optionally substituted 3-pyridyl or 6-quinolyl; X1 = -C(Y1)-NH-; Y1 = O; R = each optionally substituted furyl, thienyl, or phenyl; provided that A1 may have one to three substituents and R has one or two substituents), salts of the comds., or hydrates of either are disclosed. Thus, a solution of 2-aminonicotinic acid and [5-(3-chlorobenzyl)furan-2-ylmethyl]amine in DMF was treated with benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate and Et3N and stirred at 80° for 40 min to give 3-amino-N-[5-(3-chlorobenzyl)furan-2-ylmethyl]nicotinamide (II). II showed min. inhibitory concentration of 6.25 µg/mL against yeast expressing plasmidom GWT1 gene (optGWT1).

IT 849810-07-3P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of heterocyclic comds. such as nicotinamide

quinolinecarboxamide derivs. as antimalaria agents)


RN 849810-87-3 CAPLUS

6-Quinoxalinecarboxamide, N-[(3-phenoxyphenyl)methyl]-, mono(trifluoroacetate) (9CI) (CA INDEX NAME)

CM 1

CRN 849810-86-2

CMF C22 H17 N3 O2



CM 2

CRN 76-05-1

CMF C2 H F3 O2



REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 5 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STM
ACCESSION NUMBER: 2006:54368 CAPLUS
DOCUMENT NUMBER: 144:150635
TITLE: Preparation of amino acid amide derivatives as inhibitors of histone deacetylase
INVENTOR(S): Chakravarty, Prasun K.; Colletti, Steven L.; Ingenito, Raffaele; Jones, Philip; Meinke, Peter T.; Muraglia, Ester; Petrocchi, Alessia; Rowley, Michael; Scarpelli, Rita; Steinkuhler, Christian
PATENT ASSIGNER(S): Istituto di Ricerche di Biologia Molecolare p Angeletti S.p.A., Italy; Merck & Co. Inc.
SOURCE: PCT Int. Appl., 161 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE
WO 2006005941 A1 20060119 WO 2005-GB2729 20050711
W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HD, IL, IN, IS, JP, KE, KG, KM, KN, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MG, MH, MI, MN, MO, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CP, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TO, BW, GH, GM, KS, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
US 2004-587177P P 20040712
US 2004-610707P P 20040917

OTHER SOURCE(S): MARPAT 144:150635

AB The invention relates to compds. R1(CH2)0-3NR5COCH[NR4-X-(CH2)0-3R3](CH2)1-6COR2 [X is CH2, CO, SO2, CONH, CO2, C(S)NH or CONHSO2; R1 is (un)substituted carbalkoxy, amino groups, aryl, aryloxy, cycloalkyl, aryl or heterocyclyl; R2 is H, (un)substituted alkyl, carbonyl, CF3, cycloalkyl, aryl or heterocyclyl; R3 is H, CF3, oxo, OH, CN, halo, amino groups, (un)substituted carboxylic ester, acyl, sulfonyl groups, etc.; R4 is H or alkyl; R5 is H or together with R1(CH2)0-3N forms (un)substituted piperazinyl that are inhibitors of histone deacetylase (HDAC) and are useful for treating cancer, neurodegenerative diseases, schizophrenia, stroke and other diseases. Thus, (2S)-2-[[[(5-methoxy-2-methyl-1H-indol-3-yl)acetyl]amino]-8-oxo-N-[2-(2-phenyl-1H-indol-3-yl)ethyl]nonanamide was prepared by a multistep sequence involving reactions of Me 8-oxononanoate, ethylenediol, (S)-(-)-4-benzyl-2-oxazolidinone, 2-(2-phenyl-1H-indol-3-yl)ethanaminium chloride, and 5-methoxy-2-methyl-3-indolylacetic acid. Compds. of the invention were found to have HDAC inhibitory activity (IC50 < 30 nM).

IT 874154-44-6P 874159-11-2P 874159-15-6P
874159-36-1P 874159-71-4P 874159-77-0P
874160-17-5P 874160-23-3P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses).

(preparation of amino acid amide derivs. as inhibitors of histone deacetylase)

RN 874154-44-6 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[[[(1S)-7-oxo-1-[[[2-(2-phenyl-1H-indol-3-yl)ethyl]amino]carbonyl]octyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 874159-11-2 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[[[(1S)-1-[[[3-(3-acetylphenyl)amino]carbonyl]-7-oxononyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 874159-15-6 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[[[(1S)-1-[[[cyclopentylamino]carbonyl]-7-oxononyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 874159-36-1 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[[[(1S)-7-oxo-1-[[[3-(pyridinylamino)carbonyl]nonyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 874159-71-4 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[[[(1S)-7-oxo-1-[[[2-(2-phenyl-4-thiazolyl)methyl]amino]carbonyl]nonyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 874159-77-0 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[[[(1S)-1-[[[2-(naphthalenylamino)carbonyl]-7-oxononyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 874160-17-5 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[[[(1S)-7-oxo-1-[[[2-(phenylethyl)amino]carbonyl]nonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 874160-23-3 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[[[(1S)-7-oxo-1-[[[2-(3-phenyl-1-pyrrolidinyl)ethyl]amino]carbonyl]nonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 6 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2006:16547 CAPLUS

DOCUMENT NUMBER: 144:274236

TITLE: Synthesis and antiprotezoal activity of some new synthetic substituted quinoxalines

AUTHOR(S): Hui, Xu; Desrivot, Julie; Bories, Christian; Loiseau, Philippe M.; Franck, Xavier; Hocquemiller, Reynald; Figgere, Bruno

CORPORATE SOURCE: Address Laboratoire de Pharmacognosie et Groupe Chimiotherapie Antiparasitaire (associe au CNRS-BioCIS) Faculte de Pharmacie, Universite de Paris-Sud, Chateaufort-Malebry, 92296, Fr.

SOURCE: Bioorg. Med. Chem. Lett. (2006), 16(4), 815-820

CODEN: BMCLEB; ISSN: 0960-894X

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 144:274236

AB A set of 29 6-aminoquinoxalines and 6-quinoxalinecarboxamides are prepared and evaluated in vitro against several protozoal parasites (Leishmania donovani, Trypanosoma brucei brucei, and Trichomonas vaginalis); four compds. are active as antileishmanial agents with IC50 values of < 20 μM. While none of the brominated quinoxalines or 2,3-diphenylquinoxalines prepared are active as antiprotezoal agents, no other clear structure-activity relationship among the quinoxalines prepared is found.


IT 878290-07-4P 878290-12-1P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)


(preparation of aminoquinoxalines and quinoxalinecarboxamides, their antiprotezoal structure-activity relationships, and their activities against Leishmania donovani, Trypanosoma brucei brucei, and Trichomonas vaginalis)

RN 878290-07-4 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[[[2,3-diphenyl-6-quinoxaliny]carbonyl]-N-(3-methoxy-2-dibenzofurenyl)-2,3-diphenyl- (9CI) (CA INDEX NAME)



RN 87290-12-1 CAPLUS
CN 6-Quinoxalinecarboxamide, N-((2,3-diphenyl-6-quinoxalinyloxy)carbonyl)-N-(9-ethyl-9H-carbazol-3-yl)-2,3-diphenyl- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


1.13 ANSWER 7 OF 181 CAPLUS COPYRIGHT 2004 ACS ON STM
ACCESSION NUMBER: 2005:135084 CAPLUS
DOCUMENT NUMBER: 144:88701
TITLE: Charge-transport materials, methods of fabrication thereof, and methods of use thereof
INVENTOR(S): Marler, Seth; Kaafarani, Bilal; Barlow, Steve; Kippelen, Bernhard; Domercq, Benoit; Zhang, Qing; Kondo, Takeshi
PATENT ASSIGNEE(S): Georgia Tech Research Corporation, USA
SOURCE: PCT Int. Appl., 219 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005123737	A2	20051229	WO 2005-US20998	20050614
WO 2005123737	C2	20060406		

M: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GR, GM, HR, HU, ID, IL, IN, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MH, MI, MN, MO, MP, MQ, MR, MU, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PK, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
RN: BM, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GO, GW, ML, MR, NE, SN, TD, TG


PRIORITY APPLN. INFO.: US 2004-579308P P 20040614
AB Briefly described, embodiments of this disclosure include charge-transport materials (e.g., 2,3,8,9,14,15-hexakis(dodecylsulfonyl)-5,6,11,12,17,18-hexaazatrinaphthylene), methods of forming charge-transport materials, and methods of using the charge-transport materials. The charge-transport materials can be used in organic electronic devices such as organic light-emitting diodes, lasers, photovoltaic cells, photodetectors, active and passive electronic devices, and memories.
IT 872140-83-5P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,15-tricarboxylic acid tri(pentafluorophenyl)methyl ester 872140-84-6P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,14-tricarboxylic acid tri(pentafluorophenyl)methyl ester
RL: DEV (Device component use); IMF (Industrial manufacture); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses) (production of charge-transport materials containing hexaazatrinaphthylene for organic electronic devices)

RN 872140-83-5 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris((pentafluorophenyl)methyl) ester (9CI) (CA INDEX NAME)




RN 872140-84-6 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,14-tricarboxylic acid, tris((pentafluorophenyl)methyl) ester (9CI) (CA INDEX NAME)


(production of charge-transport materials containing hexaazatrinaphthylene for organic electronic devices)
RN 444579-17-3 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, triethyl ester (9CI) (CA INDEX NAME)



RN 872140-78-8 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,14-tricarboxylic acid, triethyl ester (9CI) (CA INDEX NAME)




RN 872140-79-9 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tridodecyl ester (9CI) (CA INDEX NAME)




PAGE 2-A

IT 444579-17-3P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,15-tricarboxylic acid triethyl ester 872140-78-8P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,14-tricarboxylic acid triethyl ester 872140-79-9P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,15-tricarboxylic acid tridodecyl ester 872140-80-2P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,14-tricarboxylic acid tridodecyl ester 872140-81-3P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,15-tricarboxylic acid tris-(2,3,3,3,4,4,4-heptafluorobutyl) ester 872140-82-6P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,14-tricarboxylic acid tris-(2,2,3,3,4,4,4-heptafluorobutyl) ester 872140-83-7P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,15-tricarboxylic acid tris-(2-methyl-butyl) ester 872140-86-8P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,14-tricarboxylic acid tris-(2-methyl-butyl) ester 872140-87-9P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,15-tricarboxylic acid tris-(2-naphthalen-1-yl-ethyl) ester 872140-88-0P, 872140-89-1P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,15-tricarboxylic acid tribenzylester 872140-90-4P, 5,6,11,12,17,18-Hexaaza-trinaphthylene-2,8,14-tricarboxylic acid tribenzylester
RL: IMF (Industrial manufacture); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)




RN 872140-80-2 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,14-tricarboxylic acid, tridodecyl ester (9CI) (CA INDEX NAME)



RN 872140-81-3 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris(2,2,3,3,4,4,4-heptafluorobutyl) ester (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B

—CF₂—CF₃

—CF₂—CF₃

RN 872140-82-4 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,14-tricarboxylic acid, tris(2,2,3,3,4,4,4-heptafluorobutyl) ester (9CI) (CA INDEX NAME)




PAGE 1-A


PAGE 1-B

—CF₂—CF₃

RN 872140-85-7 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris(2-methylbutyl) ester (9CI) (CA INDEX NAME)





RN 872140-86-8 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,14-tricarboxylic acid, tris(2-methylbutyl) ester (9CI) (CA INDEX NAME)



RN 872140-87-9 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris[2-(1-naphthalenyl)ethyl] ester (9CI) (CA INDEX NAME)

PAGE 1-A






PAGE 2-A




PAGE 3-A

RN 872140-88-0 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,14-tricarboxylic acid, tri[2-(1-naphthalenyl)ethyl] ester (9CI) (CA INDEX NAME)




PAGE 1-A


PAGE 2-A



RN 872140-89-1 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tri(phenylmethyl) ester (9CI) (CA INDEX NAME)




RN 872140-90-4 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,14-tricarboxylic acid, tri(phenylmethyl) ester (9CI) (CA INDEX NAME)



L13 ANSWER 8 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2005:1247836 CAPLUS
DOCUMENT NUMBER: 144:128936
TITLE: Design, Synthesis, and Binding Affinities of Potential Positron Emission Tomography (PET) Ligands for Visualization of Brain Dopamine D3 Receptors
AUTHOR(S): Leopoldo, Marcello; Lacivita, Enza; De Giorgio, Paola; Colabufio, Nicola A.; Niso, Mauro; Berardi, Francesco; Perrone, Roberto
CORPORATE SOURCE: Dipartimento Farmaco-Chimico, Universita degli Studi di Bari, Bari, 70125, Italy
SOURCE: Journal of Medicinal Chemistry (2006), 49(1), 358-365
CODEN: JMCMAH; ISSN: 0022-2623
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English


OTHER SOURCE(S): CASREACT 144:128936
OI




AB The synthesis of compds. I [R1 = 7-methoxybenzofuran-2-yl, quinoxalin-6-yl, 3-(2-pyrimidinyl)phenyl, 5-(2-furyl)-3-pyrazolyl, etc.; R2 = 2-MeOC6H4, 2-benzimidazolyl, 5-methoxy-2-benzisoxazolyl, etc.], structurally related to the high-affinity dopamine D3 receptor ligand N-[4-(4-(2,3-dichlorophenyl)piperazin-1-yl)butyl]-7-methoxy-2-benzofurancarboxamide (II), is reported. All compds. were specifically designed as potential PET radioligands for brain D3 receptors visualization, having lipophilicity within a range for high brain uptake and weak nonspecific binding (2 < ClogP < 3.5) and bearing a methoxy substituent for easy access to labeling with the positron emitter isotope ¹¹C. 1 (R1 = 4-(4-morpholinyl)phenyl, 4-(1-imidazolyl)phenyl, 5-(2-furyl)-3-pyrazolyl; R2 = 5-methoxy-2-benzisoxazolyl) displayed good D3 receptor affinities (Ki values 38.0, 22.6, and 21.3 nM, resp.) and were selective over D2 receptor. Moreover, these compds. were able to permeate the Caco-2 cell monolayer, differently from compound II. Although the goal to identify potential PET radioligands with subnanomolar affinities for D3 receptor was not achieved, the proposed strategy could be a starting point for future developments.

IT 873662-58-9P 873662-69-2P
RL: PAC (Pharmacological activity); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
[preparation, lipophilicity and brain dopamine D3 receptor binding affinities of N-(aryl)piperazinylbutyl heteroarylcarboxamides as potential positron emission tomog. ligands]

RN 873662-58-9 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[4-[4-(5-methoxy-1,2-benzisoxazol-3-yl)-1-piperazinyl]butyl]- (9CI) (CA INDEX NAME)



RN 873662-69-2 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[4-[4-(2-methoxyphenyl)-1-piperazinyl]butyl]- (9CI) (CA INDEX NAME)




IT 873662-82-9P 873662-89-6P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation, lipophilicity and brain dopamine D3 receptor binding affinities of N-(arylpiperaziny)butyl heteroarylcarboxamides as potential positron emission tomog. ligands)

RM 873662-82-9 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[4-[4-(5-methoxy-1,2-benzisoxazol-3-yl)-1-piperaziny]butyl]-, ethanedioate (1:2) (9CI) (CA INDEX NAME)

CM 1

CRN 873662-58-9
CMF C25 H28 N6 O3




CM 2

CRN 144-62-7
CMF C2 H2 O4



RM 873662-89-6 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[4-[4-(2-methoxyphenyl)-1-piperaziny]butyl]-, dihydrochloride (9CI) (CA INDEX NAME)



●2 HCl

REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L13 ANSWER 9 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2005:1224419 CAPLUS
DOCUMENT NUMBER: 143:454394
TITLE: Preparation of quinoxalin-2-one derivatives as herbicide safeners
INVENTOR(S): Schaper, Wolfgang; Willms, Lothar; Rosinger, Christopher; Hacker, Erwin; Rose, Eckhard; Schmutzler, Dirk
PATENT ASSIGNEE(S): Bayer Cropscience GmbH, Germany
SOURCE: U.S. Pat. Appl. Publ., 97 pp.
CODEN: USXKCO
DOCUMENT TYPE: Patent

LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2005256000	A1	20051117	US 2005-127016	20050511
DE 102004023332	A1	20060119	DE 2004-102004023332	20040512
WO 2005112630	A1	20051201	WO 2005-EP4445	20050426

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, BZ, CF, CG, CI, CM, GN, GQ, GW, ML, MR, NE, NG, TD, TG


PRIORITY APPLN. INFO.: DE 2004-102004023332A 20040512
OTHER SOURCE(S): MARPAT 143:454394
OI



AB The quinoxalin-2-one derivs. I [X = O or S; Y = halo, cyano, nitro, alkyl, alkenyl, alkynyl, etc.; n = 0, 1, 2, 3 or 4; R1 = H, OH, NH2, alkylamino, dialkylamino, (un)substituted alkyl, alkenyl, alkynyl or alkoxy, cycloalkyl, cycloalkenyl, aryl or heterocyclyl; R2 = H or (un)substituted alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl or heterocyclyl] or I salts are prepared as herbicide safeners.

IT 869312-47-0P
RL: AGR (Agricultural use); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation as herbicide safener)

RM 869312-47-0 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 1,2-dihydro-2-oxo-3-(2-thienyl)-, ethyl ester (9CI) (CA INDEX NAME)




L13 ANSWER 10 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2005:1184873 CAPLUS
DOCUMENT NUMBER: 144:98799
TITLE: High Charge-Carrier Mobility in an Amorphous


AUTHOR(S): Hexaazatrinaphthylene Derivative
Kaaferani, Bilal R.; Kondo, Takeshi; Yu, Junsheng; Zhang, Qing; Dattilo, Davide; Risko, Chad; Jones, Simon C.; Barlow, Stephen; Domercq, Benoit; Amy, Fabrice; Kahn, Antoine; Bredas, Jean-Luc; Kippelen, Bernard; Marder, Seth R.
CORPORATE SOURCE: Center for Organic Photonics and Electronics (COPE), School of Chemistry and Biochemistry and School of Electrical and Computer Engineering, Georgia Institute of Technology, Atlanta, GA, 30332, USA
SOURCE: Journal of the American Chemical Society (2005), 127(47), 16358-16359
CODEN: JACSAT; ISSN: 0002-7863
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English

AB An isomeric mixture of a tris(pentafluorobenzyl ester) derivative of hexaazatrinaphthylene forms stable amorphous films with an effective charge-carrier mobility of 0.02 cm²/Vs, while the pure 2,8,15-isomer exhibits widely differing morphologies and carrier mobilities (0.001-0.07 cm²/Vs), depending critically on the processing conditions.

IT 872140-83-5P 872140-94-6P
RL: DEV (Device component use); PRP (Properties); SPN (Synthetic preparation); PREP (Preparation); USES (Uses)
(carrier mobility in amorphous hexaazatrinaphthylene derivative)
RM 872140-83-5 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris[(pentafluorophenyl)methyl] ester (9CI) (CA INDEX NAME)




RM 872140-84-6 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,14-tricarboxylic acid, tris[(pentafluorophenyl)methyl] ester (9CI) (CA INDEX NAME)




PAGE 1-A

PAGE 2-A

IT 444579-17-3 872140-78-8
RL: PRP (Properties)
(carrier mobility in amorphous hexaazatrinaphthylene derivative)
RM 444579-17-3 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, triethyl ester (9CI) (CA INDEX NAME)



RN 872140-78-8 CAPLUS
CN DiQuinoxaline[2,3-a:2',3'-c]phenazine-2,8,14-tricarboxylic acid, triethyl ester (9CI) (CA INDEX NAME)





REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L13 ANSWER 11 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2005:1050874 CAPLUS
DOCUMENT NUMBER: 143:326207
TITLE: Preparation and pharmaceutical compositions of pyrrolidine derivatives as inhibitors of dipeptidyl peptidase-iv (DPP-iv)
INVENTOR(S): Akritopoulou-Zanze, Irini; Darczak, Daria; Dinges, Jurgen; Djuric, Stevan W.; Hoff, Ethan D.; Kopecka, Hana A.; Patel, Jyoti R.; Pei, Zhonghua; Shuai, Qi; Sarria, Kathy; Sham, Hing L.; Wiedeman, Paul E.
PATENT ASSIGNER(S): USA
SOURCE: U.S. Pat. Appl. Publ., 51 pp.
CODEN: USXXCO
DOCUMENT TYPE: Patent

LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2005215603	A1	20050929	US 2004-795622	20040308
PRIORITY APPL. INFO.:			US 2004-795622	20040308
OTHER SOURCE(S):		MARPAT 143:326207		




AB Title compds. I [R1 = aryl, alkyl, cycloalkyl, etc.; D = CO, O, SO2, CONH, etc.; L = bond, -CH2-, aryl, etc.; A = CO, NHSO2, NHCO, etc.; X = CHF, CH2, O, S, etc.], and their pharmaceutically acceptable salts, are prepared and disclosed as inhibitors of dipeptidyl peptidase IV (DPP-IV). Thus, e.g., II was prepared by amidation of (2S,4R)-4-aminomethyl-2-(thiazolidine-3-carbonyl)pyrrolidine-1-carboxylic acid tert-Bu ester (preparation given) with benzoyl chloride. I were found to inhibit DPP-IV induced fluorescence with inhibitory consts. in a range of about 0.0005 μ M to about 7 μ M. I should prove useful for the prevention or treatment of diabetes, especially type II diabetes, as well as hyperglycemia, syndrome X, hyperinsulinemia, obesity, atherosclerosis, and various immunomodulatory diseases.
IT 865294-88-8P 865296-56-6P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(drug candidate; preparation of pyrrolidine derivs. as inhibitors of dipeptidyl peptidase-iv (DPP-iv))
RN 865294-88-8 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[[(3S,5S)-5-(3-thiazolidinylcarbonyl)-3-pyrrolidinyl]methyl]- (9CI) (CA INDEX NAME)
Absolute stereochemistry.





RN 865296-56-6 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[[(3S,5S)-5-(1-pyrrolidinylcarbonyl)-3-pyrrolidinyl]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L13 ANSWER 12 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2005:1028081 CAPLUS
DOCUMENT NUMBER: 143:326207
TITLE: Preparation and pharmaceutical compositions of pyrrolidine derivatives as inhibitors of dipeptidyl peptidase-iv (DPP-iv)
INVENTOR(S): Akritopoulou-Zanze, Irini; Darczak, Daria; Dinges, Jurgen; Djuric, Stevan W.; Hoff, Ethan D.; Kopecka, Hana A.; Patel, Jyoti R.; Pei, Zhonghua; Shuai, Qi; Sarria, Kathy; Sham, Hing L.; Wiedeman, Paul E.
PATENT ASSIGNER(S): USA
SOURCE: U.S. Pat. Appl. Publ., 50 pp.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2005209249	A1	20050922	US 2005-75319	20050308
PRIORITY APPL. INFO.:			US 2004-551079P	P 20040308
OTHER SOURCE(S):		MARPAT 143:326202		




AB Title compds. I [R1 = aryl, alkyl, cycloalkyl, etc.; D = CO, O, SO2, CONH, etc.; L = bond, -CH2-, aryl, etc.; A = CO, NHSO2, NHCO, etc.; X = CHF, CH2, O, S, etc.], and their pharmaceutically acceptable salts, are prepared and disclosed as inhibitors of dipeptidyl peptidase IV (DPP-IV). Thus, e.g., II was prepared by amidation of (2S,4R)-4-aminomethyl-2-(thiazolidine-3-carbonyl)pyrrolidine-1-carboxylic acid tert-Bu ester (preparation given) with benzoyl chloride. I were found to inhibit DPP-IV induced fluorescence with inhibitory consts. in a range of about 0.0005 μ M to about 7 μ M. I should prove useful for the prevention or treatment of diabetes, especially type II diabetes, as well as hyperglycemia, syndrome X, hyperinsulinemia, obesity, atherosclerosis, and various immunomodulatory diseases.
IT 865294-88-8P 865296-56-6P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(drug candidate; preparation of pyrrolidine derivs. as inhibitors of dipeptidyl peptidase-iv (DPP-iv))
RN 865294-88-8 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[[(3S,5S)-5-(3-thiazolidinylcarbonyl)-3-pyrrolidinyl]methyl]- (9CI) (CA INDEX NAME)
Absolute stereochemistry.




RN 865296-56-6 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[[(3S,5S)-5-(1-pyrrolidinylcarbonyl)-3-pyrrolidinyl]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L13 ANSWER 13 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 2005:989948 CAPLUS
 DOCUMENT NUMBER: 143:433305
 TITLE: Asymmetric cooperativity in tandem hybridization of enantiomeric metal complex-tethered short fluorescent DNA probes
 AUTHOR(S): Kitamura, Yusuke; Ihara, Toshihiro; Okada, Kenji; Tsujimura, Yusuke; Shirasaka, Yoshinori; Tazaki, Masato; Jyo, Akinori
 CORPORATE SOURCE: Department of Applied Chemistry and Biochemistry, Kumamoto University, Kumamoto, 860-8555, Japan
 SOURCE: Chemical Communications (Cambridge, United Kingdom) (2005), (36), 4523-4525
 CODEN: CHCOFS; ISSN: 1359-7345
 PUBLISHER: Royal Society of Chemistry
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The complex [Ru(phen)2(dppz)]2+ (phen = 1,10-phenanthroline, dppz = dipyrro[3,2-a:2',3'-c]phenazine) was attached to the 5' end of a short oligonucleotide to form conjugates, the Δ-isomer of which showed a high cooperativity during the recognition of the repetitive sequence, while the Λ-isomer did not.
 IT 663942-78-7
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USSS (Uses) (asym. cooperativity during tandem hybridization of enantiomeric metal complex-tethered short fluorescent DNA probes to human telomere repetitive DNA)
 RN 663942-78-7 CAPLUS
 CN Adenosine, 2'-deoxy-5'-O-[[[6-[[[dipyrro[3,2-a:2',3'-c]phenazin-11-ylcarbonyl]amino]hexyl]oxy]hydroxyphosphinyl]cytidyl- (3' → 5')-2'-deoxycytidyl- (3' → 5')-2'-deoxycytidyl- (3' → 5')-thymidyl- (3' → 5')-2'-deoxyadenyl- (3' → 5')-2'-deoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



PAGE 1-A





PAGE 1-B



PAGE 2-A

IT 868744-57-4 868744-70-1 868745-17-9
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USSS (Uses) (asym. cooperativity in tandem hybridization of enantiomeric metal complex-tethered short fluorescent DNA probes)
 RN 868744-57-4 CAPLUS
 CN Ruthenate(4-), [5'-O-[[[6-[[[dipyrro[3,2-a:2',3'-c]phenazin-11-yl-κN4,κN5]carbonyl]amino]hexyl]oxy]hydroxyphosphinyl]-2'-deoxycytidyl- (3' → 5')-2'-deoxycytidyl- (3' → 5')-2'-deoxycytidyl- (3' → 5')-thymidyl- (3' → 5')-2'-deoxyadenyl- (3' → 5')-2'-deoxyadenosinate(6-)]bis(1,10-phenanthroline-κN1,κN10)-, pentahydrogen, (OC-6-33)- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



PAGE 2-A




● 5 H⁺





RN 868744-70-1 CAPLUS
 CN Ruthenate (4-), [5'-O-[[[6-[[[dipyrido[3,2-a:2',3'-c]phenazin-11-yl-
 <N4,<N5>carbonylamino]hexyl]oxy]hydroxyphosphinyl]-2'-
 deoxycytidyl- (3' ->5') -2'-deoxycytidyl- (3' ->5') -2'-
 deoxycytidyl- (3' ->5') -thymidyl- (3' ->5') -2'-deoxyadenyl-
 (3' ->5') -2'-deoxyadenosin-6-yl]]bis(1,10-phenanthroline-
 <N1,<N10>-), pentahydrogen, (OC-6-33-Δ)- (9CI) (CA INDEX
 NAME)




● 5 H⁺



RN 868745-17-9 CAPLUS
 CN Ruthenate (4-), [5'-O-[[[6-[[[dipyrido[3,2-a:2',3'-c]phenazin-11-yl-
 <N4,<N5>carbonylamino]hexyl]oxy]hydroxyphosphinyl]-2'-
 deoxycytidyl- (3' ->5') -2'-deoxycytidyl- (3' ->5') -2'-
 deoxycytidyl- (3' ->5') -thymidyl- (3' ->5') -2'-deoxyadenyl-
 (3' ->5') -2'-deoxyadenosin-6-yl]]bis(1,10-phenanthroline-
 <N1,<N10>-), pentahydrogen, (OC-6-33-Δ)- (9CI) (CA
 INDEX NAME)



● 5 H⁺



REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 14 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2005:800492

DOCUMENT NUMBER: 143:386895

TITLE: A phase-switch purification approach for the expedient removal of tagged reagents and scavengers following their application in organic synthesis

AUTHOR(S): Liu, Jason; Baxendale, Ian R.; Leithwaite, Russell A.; Ley, Steven V.

CORPORATE SOURCE: Department of Chemistry, University of Cambridge, Cambridge, CB2 1EW, UK

SOURCE: Organic & Biomolecular Chemistry (2005), 3(17), 3140-3160

CODEN: OBCRAK; ISSN: 1477-0520

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

AB In this paper a variety of expedient chemical transformations and purifications achieved via a generic catch and release methodology, based on a synthetically inert bipyridyl chelating tag that can be selectively captured with a resin-bound copper(II) species, were reported. Utilizing this approach it was possible to derive many of the same benefits associated with both solid phase synthesis and supported reagent methods.


IT 866789-75-5P

RL: PUR (Purification or recovery); SPN (Synthetic preparation); PREP (Preparation)


(Preparation of amides using amines and carboxylic acid as reactants and N-(cyclohexylcarboxymethyl)bipyridine amine as coupling agent and study of phase-switch purification approach for expedient removal of tagged reagents and scavengers)

RN 866789-75-5 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[(1,3,5-trimethyl-1H-pyrazol-4-yl)methyl]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 61 THERE ARE 61 CITED REFERENCES AVAILABLE FOR THIS



RN 853914-90-6 CAPLUS


CN Copper(I), (ethyl dipyrro[3,2-a:2',3'-c]phenazine-11-carboxylate-kN4,kN5)bis(triphenylphosphine)-, (T-4)-, tetrafluoroborate(1-) (9CI) (CA INDEX NAME)

CM 1

CRN 853914-89-3

CMF C57 H44 Cu N4 O2 P2

CCI CCS



CM 2

CRN 14874-70-5

CMF B F4

CCI CCS



IT 862288-29-7

RL: PRP (Properties) (Raman spectroscopy and DFT calcs. in study of ground- and excited states of Cu(I) and Re(I) complexes with dipyrrophenazine ligands)

RN 862288-29-7 CAPLUS

CN Rhenate(1-), tricarboxylchloro(ethyl dipyrro[3,2-a:2',3'-c]phenazine-11-carboxylate-kN4,kN5)-, (OC-6-44)- (9CI) (CA INDEX NAME)

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 15 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2005:504199

DOCUMENT NUMBER: 143:319127

TITLE: Photoexcitation in Cu(I) and Re(I) Complexes Containing Substituted Dipyrro[3,2-a:2',3'-c]phenazine: A Spectroscopic and Density Functional Theoretical Study

AUTHOR(S): Walsh, Penny J.; Gordon, Keith C.; Lundin, Natasha J.; Blackman, Allan G.

CORPORATE SOURCE: Department of Chemistry, University of Otago, Dunedin, N. Z.

SOURCE: Journal of Physical Chemistry A (2005), 109(26), 5933-5942

CODEN: JPCAPH; ISSN: 1089-5639

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English


AB Copper(I) and rhenium(I) complexes [Cu(PPh3)2(dppz-11-COOEt)]BF4, [Cu(PPh3)2(dppz-11-Br)]BF4, [Re(CO)3Cl(dppz-11-COOEt)] and [Re(CO)3Cl(dppz-11-Br)] (dppz-11-COOEt = dipyrro[3,2-a:2',3'-c]phenazine-11-carboxylic Et ester, dppz-11-Br = 11-bromo-dipyrro[3,2-a:2',3'-c]phenazine) have been studied using Raman, resonance Raman, and transient resonance Raman (TR2) spectroscopy, in conjunction with computational chemical DFT (B3LYP) frequency calcs. with a 6-31G(d) basis set for the ligands and copper(I) centers and an effective core potential (LANL2DZ) for rhenium in the rhenium(I) complexes show close agreement with the exptl. nonresonance Raman spectra. Modes that are phenazine-based, phenanthroline-based, and delocalized across the entire ligand structure were identified. The nature of the absorbing chromophores at 356 nm for ligands and complexes was established using resonance Raman spectroscopy in concert with vibrational assignments from calcs. This anal. reveals that the dominant chromophore for the complexes measured at 356 nm is ligand-centered (LC), except for [Re(CO)3Cl(dppz-11-Br)], which appears to have addnl. chromophores at this wavelength. Calcs. on the reduced complexes, undertaken to model the metal-to-ligand charge transfer (MLCT) excited state, show that the reducing electron occupies a ligand MO that is delocalized across the ligand structure. Resonance Raman spectra (λexc = 514.5 nm) of the reduced rhenium complexes show a similar spectral pattern to that observed in [Re(CO)3Cl(dppz)]⁺; the measured bands are therefore attributed to ligand radical anion modes. These bands lie at 1583-1593 cm⁻¹ for [Re(CO)3Cl(dppz-11-COOEt)] and 1611 cm⁻¹ for [Re(CO)3Cl(dppz-11-Br)]. The thermally equilibrated excited states are examined using nanosecond-TR2 spectroscopy (λexc = 354.7 nm). The TR2 spectra of the ligands provide spectral signatures for the 3LC state. A band at 1382 cm⁻¹ is identified as a marker for the 3LC states of both ligands. TR2 spectra of the copper and rhenium complexes of dppz-11-Br show this 3LC band, but it is not prominent in the spectra of [Cu(PPh3)2(dppz-11-COOEt)]⁺ and [Re(CO)3Cl(dppz-11-COOEt)]. Calcs. suggest that the lowest triplet states of both of the rhenium(I) complexes and [Cu(PPh3)2(dppz-11-Br)] are metal-to-ligand charge transfer in nature, but the lowest triplet state of [Cu(PPh3)2(dppz-11-COOEt)]⁺ appears to be LC in character.

IT 767350-96-9 853914-90-6

RL: PRP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); PROC (Process) (Raman spectroscopy and DFT calcs. in study of ground- and excited states of Cu(I) and Re(I) complexes with dipyrrophenazine ligands)

RN 767350-96-9 CAPLUS

CN Rhodium, tricarboxylchloro(ethyl dipyrro[3,2-a:2',3'-c]phenazine-11-carboxylate-kN4,kN5)-, (OC-6-44)- (9CI) (CA INDEX NAME)




IT 853914-87-1

RL: PRP (Properties) (ligand; Raman spectroscopy and DFT calcs. in study of ground- and excited states of Cu(I) and Re(I) complexes with dipyrrophenazine ligands)

RN 853914-87-1 CAPLUS

CN Dipyrro[3,2-a:2',3'-c]phenazine-11-carboxylic acid, ethyl ester (9CI) (CA INDEX NAME)



REFERENCE COUNT: 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 16 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2005:339438

DOCUMENT NUMBER: 143:70403

TITLE: Complexes of Functionalized Dipyrro[3,2-a:2',3'-c]phenazine: A Synthetic, Spectroscopic, Structural, and Density Functional Theory Study

AUTHOR(S): Lundin, Natasha J.; Walsh, Penny J.; Howell, Sarah L.; McGarvey, John J.; Blackman, Allan G.; Gordon, Keith C.

CORPORATE SOURCE: Department of Chemistry, MacDiarmid Institute for Advanced Materials and Nanotechnology, University of Otago, Dunedin, N. Z.

SOURCE: Inorganic Chemistry (2005), 44(10), 3551-3560

CODEN: INOCAT; ISSN: 0020-1669

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal


LANGUAGE: English

OTHER SOURCE(S): CASREACT 143:70403


AB The ligands 11-bromodipyrro[3,2-a:2',3'-c]phenazine and Et dipyrro[3,2-a:2',3'-c]phenazine-11-carboxylate were prepared and coordinated to Ru(II), Re(I), and Cu(I) metal centers. The electronic effects of substitution of dipyrro[2,3-a:3',2'-c]phenazine (dppz) were studied by spectroscopy and electrochem., and some photophys. properties were studied. The crystal structures of [Re(I)(CO)3Cl] (L = Et dipyrro[3,2-a:2',3'-c]phenazine-11-carboxylate or 11-bromodipyrro[3,2-a:2',3'-c]phenazine) are presented. D. functional theory calcs. on the complexes show only small deviations in bond lengths and angles (most bonds within 0.02 Å, most angles within 2°) from the

crystallog. data. Also, the vibrational spectra of the strongest Raman and IR bands are predicted to within an average 6 cm⁻¹ for [Re(L)(CO)3Cl] and [Cu(L)(triphenylphosphine)2]BF4 (in the 1000-1700 cm⁻¹ region). Spectroscopic and electrochem. evidence suggest that reduction of the complex causes structural changes across the entire dppz ligand. This is unusual as dppz-based ligands typically have electrochem. properties that suggest charge localization with reduction on the phenazine portion of the ligand. The excited-state lifetimes of the complexes were measured, and they range from ca. 200 ns for the [Ru(L)(2,2'-bipyridine)2](PF6)2 complexes to over 2 μs for [Cu(11-bromodipyrido[3,2-a:2',3'-c]phenazine)(PPh3)2](BF4) at room temperature. The emission spectra suggest that the unusually long-lived excited states of the Cu complexes result from metal-to-ligand charge transfer (MLCT) transitions as they are completely quenched in MeOH. Electroluminescent films may be fabricated from these compds.; they show MLCT state emission even at low doping levels (<0.1% by weight in poly(vinylcarbazole) polymer matrix).

IT 767350-96-99
 RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); TEM (Technical or engineered material use); PRSP (Preparation); PROC (Process); USES (Uses) (preparation, crystal structure, exptl. and calculated mol. structure, fluorescence, reduction potentials, vibrational spectra and application in electroluminescent film)
 RN 767350-96-9 CAPLUS
 CN Rhenium, tricarbonylchloro(ethyl dipyrro[3,2-a:2',3'-c]phenazine-11-carboxylate-κN4,κN5)-, (OC-6-44)- (9CI) (CA INDEX NAME)




IT 853914-87-1P
 RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PRSP (Preparation); PROC (Process); RACT (Reactant or reagent) (preparation, mol. structure from DFT calcs., fluorescence, reduction potentials and complexation with copper, rhenium and ruthenium)
 RN 853914-87-1 CAPLUS
 CN Dipyrro[3,2-a:2',3'-c]phenazine-11-carboxylic acid, ethyl ester (9CI) (CA INDEX NAME)



IT 853914-91-7P 853914-95-1P
 RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); PRSP (Preparation); PROC (Process); RACT (Reactant or reagent) (preparation, mol. structure from DFT calcs., fluorescence, reduction potentials and complexation with copper, rhenium and ruthenium)

process); PRP (Properties); SPN (Synthetic preparation); PRSP (Preparation); PROC (Process) (preparation, mol. structure from DFT calcs., fluorescence, reduction potentials of)
 RN 853914-91-7 CAPLUS
 CN Copper(1+), (ethyl dipyrro[3,2-a:2',3'-c]phenazine-11-carboxylate-κN4,κN5)bis(triphenylphosphine)-, (T-4)-, tetrafluoroborate(1-), monohydrate (9CI) (CA INDEX NAME)


CM 1
 CRN 853914-90-6
 CMF C57 H44 Cu N4 O2 P2 . B F4
 CM 2
 CRN 853914-89-3
 CMF C57 H44 Cu N4 O2 P2
 CCI CCS




CM 3
 CRN 14674-70-5
 CMF B F4
 CCI CCS



RN 853914-95-1 CAPLUS
 CN Ruthenium(2+), bis(2,2'-bipyridine-κN1,κN1')(ethyl dipyrro[3,2-a:2',3'-c]phenazine-11-carboxylate-κN4,κN5)-, (OC-6-31)-, bis(hexafluorophosphate(1-)) (9CI) (CA INDEX NAME)
 CM 1
 CRN 853914-94-0
 CMF C41 H30 N8 O2 Ru
 CCI CCS



CM 2
 CRN 16919-18-9
 CMF F6 P
 CCI CCS




REFERENCE COUNT: 60 THERE ARE 60 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 17 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2005:324138 CAPLUS
 DOCUMENT NUMBER: 142:392428
 TITLE: Preparation of heterocyclic compounds as antifungal agents
 INVENTOR(S): Nakamoto, Kazutaka; Tsukada, Itaru; Tanaka, Keigo; Matsukura, Masayuki; Haneda, Toru; Inoue, Satoshi; Ueda, Norihiro; Abe, Shinya; Hata, Katsura; Watanabe, Naoki
 PATENT ASSIGNOR(S): Eisai Co., Ltd., Japan
 SOURCE: PCT Int. Appl., 418 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE

WO 2005033079	A1	20050414	WO 2004-JP14063	20040927
W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
WO 2006016548	A1	20060216	WO 2005-JP14505	20050808
W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MX, MY, NA, NO, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, ML, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, GN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
PRIORITY APPL. INFO.:			JP 2003-342273	A 20030930
			JP 2004-68186	A 20040310
			JP 2004-232617	A 20040809
			WO 2004-JP14063	A 20040927
			JP 2005-82760	A 20050322


OTHER SOURCE(S): MARPAT 142:392428
 GI



AB The title compds., e.g. I [ring A1 is optionally substituted 3-pyridyl, optionally substituted quinolyl, etc.; X1 is NHCO, etc.; and ring E is furyl, thienyl, pyrrolyl, Ph, pyridyl, tetrazolyl, thiazolyl, or pyrazolyl; provided that A1 may have one to three substituents and E has one or two substituents], are prepared 2,6-Diamino-N-(5-(4-fluorophenoxy)furan-2-ylmethyl)nicotinamide was prepared in a multistep process. Compds. of this invention in vitro showed MIC values of 0.1 μg/mL to 6.25 μg/mL against Candida.

IT RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation of heterocyclic compds. as antifungal agents)
 RN 849810-87-3 CAPLUS
 CN 6-Quinoxalinecarboxamide, N-[1-(3-pyridinylmethyl)-, mono(trifluoroacetate) (9CI) (CA INDEX NAME)

CM 1
 CRN 849810-86-3
 CMF C22 H17 N3 O2



CH 2

CRN 76-05-1
CNF C2 H F3 O2




REFERENCE COUNT: 56 THERE ARE 56 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 18 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2005:324132 CAPLUS
DOCUMENT NUMBER: 142:392427
TITLE: Preparation of N-heterocyclyl amides and sulfonamides as p38 kinase inhibitors
INVENTOR(S): Dugar, Sundee; McEnroe, Glen
PATENT ASSIGNEE(S): Scienc Inc., USA
SOURCE: PCT Int. Appl., 195 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005033072	A2	20050414	WO 2004-US32403	20040930
WO 2005033072	A3	20060112		

W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GR, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MH, MI, MK, MN, NA, NL, NO, NZ, OM, OS, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SV, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CO, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG


PRIORITY APPLN. INFO.: US 2003-507633P P 20030930
OTHER SOURCE(S): MARPAT 142:392427
GI



AB The title compds. I [R1 = alkyl, cycloalkyl, heterocycloalkyl, aryl; L = CO, SO2; X = O, CO, (un)substituted CH2, NH; n = 0-3; R2 = H, alkyl, aryl, etc.; Y = (un)substituted NH2, OH; one of Z1 and Z2 = CH, and the other is either CH or N], useful for inhibiting p38 kinase, were prepared. E.g., a multi-step synthesis of (18)-II, starting from 4-amino-2-chloropyridine and 2-naphthyl chloride, was given. The compds. I were tested against p38 kinase in the diluted whole blood assay (biol. data were given for representative compds. I). The pharmaceutical composition comprising the compound I is disclosed.


IT 849746-04-9P 849746-62-5P 849746-71-6P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of N-heterocyclyl amides and sulfonamides as p38 kinase inhibitors)
RN 849746-04-9 CAPLUS
CN 6-Quinoxalinecarboxamide, N-ethyl-N-[2-[[[(1S)-1-phenylethyl]amino]-4-pyrimidinyl]-9CI] (CA INDEX NAME)

Absolute stereochemistry.




RN 849746-62-5 CAPLUS
CN 1-Piperidinecarboxylic acid, 4-[[[2-[[[(1S)-1-phenylethyl]amino]-4-pyrimidinyl]-6-quinoxalinyloxy]amino]methyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 849746-71-6 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-[[[(1S)-1-phenylethyl]amino]-4-pyrimidinyl]-N-(4-piperidinylmethyl)-9CI] (CA INDEX NAME)

Absolute stereochemistry.



L13 ANSWER 19 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2005:122800 CAPLUS
DOCUMENT NUMBER: 142:191201
TITLE: Antimicrobial biaryl compounds
INVENTOR(S): Jefferson, Elizabeth Anne; Swayze, Eric S.; Seth, Punit P.; Robinson, Dale E.
PATENT ASSIGNEE(S): USA
SOURCE: U.S. Pat. Appl. Publ., 25 pp., Cont.-in-part of U.S. Ser. No. 630,122.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2005032805	A1	20050210	US 2004-914356	20040809
US 6849660	B1	20050201	US 2000-630122	20000801

PRIORITY APPLN. INFO.: US 2000-630122 A2 20000801
OTHER SOURCE(S): MARPAT 142:191201

AB Provided are antibacterial biaryl compds. having micromolar MIC activity against Gram-neg. and Gram-pos. pathogens, including a methicillin-resistant *S. aureus* strain. Other embodiments of invention are methods of treating bacterial infection in a mammal by administering to the mammal an effective amount of a compound described herein. The inhibitory effect of some of the compds. on bacterial translation was determined.

IT 797770-86-6
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(antibacterial biaryl compds. in relation to inhibiting bacterial translation and overcoming methicillin resistance)

RN 797770-86-6 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[1-[4-(3-(dimethylamino)propoxy)phenyl]-2-oxo-2-(1-piperazinyl)ethyl]-1,2,3,4-tetrahydro-2,3-dioxo-9CI] (CA INDEX NAME)




L13 ANSWER 20 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2005:99161 CAPLUS
DOCUMENT NUMBER: 142:198098
TITLE: Preparation of quinoxalinecarboxamides as antivirals
INVENTOR(S): An, Haoyun; Rong, Frank; Wu, Jim; Harris, Clayton; Chow, Suetying
PATENT ASSIGNEE(S): USA
SOURCE: U.S. Pat. Appl. Publ., 18 pp.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2005026923	A1	20050203	US 2004-826439	20040415
PRIORITY APPLN. INFO.:			US 2003-463357P	20030415
OTHER SOURCE(S):			MARPAT 142:198098	
GI				

APPLICANTS!

10/826,439



I




II

AB Title compds. [I; Z = NH, O; X = OH, NH2, OR, NHR, NR2, SR; R1, R2 = H, (substituted) alkyl, alkenyl, alkynyl, aryl, fused aryl, heterocyclyl, fused heterocyclyl; R1R2 = atoms to form a 5-6 membered ring; R3 = (substituted) alkyl, alkenyl, alkynyl, aryl, fused aryl, heterocyclyl, fused heterocyclyl; R, R4 = H, (substituted) alkyl, alkenyl, alkynyl, aryl, fused aryl, heterocyclyl, fused heterocyclyl], were prepared Thus, title compound (II), (preparation from L-5-hydroxytryptophan, 3,4-diaminobenzoic acid, and 4-fluorobenzyl given) showed inhibitory activity with IC50 <10 µM in an hepatitis C virus (HCV) NS5B replicon assay.

IT 835922-72-0P 835922-73-1P 835922-74-2P
835922-76-4P 835922-78-6P 835922-79-7P
835922-81-1P 835922-82-2P 835922-95-7P
835922-96-8P 835922-97-9P 835922-98-0P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


RN 835922-72-0 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[(1S)-2-amino-1-[(5-hydroxy-1H-indol-3-yl)methyl]-2-oxoethyl]-2,3-bis(4-fluorophenyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 835922-73-1 CAPLUS
CN L-Tryptophan, N-[(2,3-bis(4-fluorophenyl)-6-quinoxaliny]carbonyl]-5-hydroxy-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 835922-74-3 CAPLUS
CN L-Tryptophan, N-[(2,3-bis(4-fluorophenyl)-6-quinoxaliny]carbonyl]-5-hydroxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 835922-76-4 CAPLUS
CN L-Tryptophan, N-[(2,3-bis(4-fluorophenyl)-6-quinoxaliny]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 835922-78-6 CAPLUS
CN L-Tryptophan, N-[(2,3-bis(4-fluorophenyl)-6-quinoxaliny]carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 835922-79-7 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[(1S)-2-amino-1-[(1H-indol-3-yl)methyl]-2-oxoethyl]-2,3-bis(4-fluorophenyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 835922-81-1 CAPLUS
CN L-Tryptophan, N-(2-phenaziny]carbonyl)-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 835922-82-2 CAPLUS
CN L-Tryptophan, N-(2-phenaziny]carbonyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 835922-95-7 CAPLUS
CN L-Tryptophan, N-[(2-cyclohexyl-3-phenyl-6-quinoxaliny]carbonyl]-5-hydroxy-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 835922-96-8 CAPLUS
CN L-Tryptophan, N-[(2-cyclohexyl-3-phenyl-6-quinoxaliny]carbonyl]-5-hydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 835922-97-9 CAPLUS
CN L-Tryptophan, N-[(3-cyclohexyl-2-phenyl-6-quinoxaliny]carbonyl]-5-hydroxy-, methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.

APPLICANTS



RN 835922-98-0 CAPLUS
CN L-Tryptophan, N-[(3-cyclohexyl-2-phenyl-6-quinoxalinyloxy)carbonyl]-5-hydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



IT 835922-83-3 835922-84-4 835922-85-5
835922-86-6 835922-87-7 835922-89-9
835922-90-2 835922-92-4
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(preparation of quinoxalinecarboxamides as antivirals)


RN 835922-83-3 CAPLUS
CN L-Tryptophan, N-[(2,3-diphenyl-6-quinoxalinyloxy)carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 835922-84-4 CAPLUS
CN L-Tryptophan, N-[(2,3-diphenyl-6-quinoxalinyloxy)carbonyl]-5-hydroxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 835922-85-5 CAPLUS
CN L-Tryptophan, N-[(2,3-diphenyl-6-quinoxalinyloxy)carbonyl]-5-hydroxy-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 835922-86-6 CAPLUS
CN L-Tryptophan, N-[(2,3-di-2-pyridinyl-6-quinoxalinyloxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 835922-87-7 CAPLUS
CN L-Tryptophan, N-[(2,3-di-2-pyridinyl-6-quinoxalinyloxy)carbonyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

↑
↓
APPLICANTS




RN 835922-89-9 CAPLUS
CN L-Tryptophan, N-[(2-(4-chlorophenyl)-3-(4-methylphenyl)-6-quinoxalinyloxy)carbonyl]-5-hydroxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 835922-90-2 CAPLUS
CN L-Tryptophan, N-[(2-(4-chlorophenyl)-3-(4-methylphenyl)-6-quinoxalinyloxy)carbonyl]-5-hydroxy-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 835922-92-4 CAPLUS
CN L-Tryptophan, N-[(2,3-di-2-furanyl-6-quinoxalinyloxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



L13 ANSWER 21 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2005:71069 CAPLUS
DOCUMENT NUMBER: 142:176856
TITLE: Preparation of quinoxaline and pyrido[2,3-b]pyrazine derivatives as PKB inhibitors for treatment of cancers
INVENTOR(S): Kawakami, Joel; Duncton, Matthew; Sherman, Dan; He, Hai-Ying; Kiselyov, Alexander; Pytowski, Broniek
PATENT ASSIGNEE(S): Inclosure Systems Incorporated, USA
SOURCE: PCT Int. Appl., 126 pp.
CODEN: PIXK02
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005007099	A2	20050127	WO 2004-US21834	20040709
WO 2005007099	A3	20050414		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, US, UZ, VC, VN, YU, ZA, ZM, ZW
R: BM, BM, GM, KE, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
PRIORITY APPL. INFO.: US 2003-486339 P 20030710
OTHER SOURCE(S): MARPAT 142:176856
OI



AB Title compds. represented by the formula I [wherein X = N or C; R1, R2 = independently H, (cyclo)alkyl, alkoxy, heterocyclyl(alkyl), (hetero)aryl, (hetero)alkyl, (un)substituted amino; R3-R5 = independently H, cyano, (hetero)aryl, (cyclo)alkyl, etc.; with a proviso] were prepared as PKB inhibitors. For example, reaction of 4,5-diaminopyrimidine with


2,2'-thienyl gave II in 19% yield. I were tested for inhibition of PKB in PKBa, PKBb and PKBy in vitro kinase assay. Thus, I and their pharmaceutical compns. are useful as PKB inhibitors for the treatment of cancers, or the inhibition of tumor growth.

IT 443111-01-19, 2,3-Bis(thiophen-2-yl)quinoxaline-6-carboxylic acid [3-(morpholin-4-yl)propyl]amide 443111-36-39, 2,3-Bis(thiophen-2-yl)quinoxaline-6-carboxylic acid [2-(2-chlorophenyl)ethyl]amide 443111-45-39, 2,3-Bis(thiophen-2-yl)quinoxaline-6-carboxylic acid [2-(morpholin-4-yl)ethyl]amide 832080-84-99, 2-Chloro-3-(thiophen-2-yl)-6-quinoxalinecarboxylic acid [2-(2-chlorophenyl)ethyl]amide 832080-85-99, 2-[[2-(Pyridin-4-yl)ethyl]amino]-3-(thiophen-2-yl)quinoxaline-6-carboxylic acid [2-(2-chlorophenyl)ethyl]amide 832080-86-19, 2-[(Benzyl)(methyl)amino]-3-(thiophen-2-yl)quinoxaline-6-carboxylic acid [2-(2-chlorophenyl)ethyl]amide 832082-00-59, 2,3-Bis(thiophen-2-yl)quinoxaline-6-carboxylic acid [2-(2-methoxyphenyl)ethyl]amide 832082-01-69, 2,3-Bis(thiophen-2-yl)quinoxaline-6-carboxylic acid [2-(2-tolyl)ethyl]amide 832082-02-79, 2,3-Bis(thiophen-2-yl)quinoxaline-6-carboxylic acid [2-(2-pyridyl)ethyl]amide 832082-03-89, 2,3-Bis(thiophen-2-yl)quinoxaline-6-carboxylic acid [2-(3-pyridyl)ethyl]amide 832082-04-99, 2,3-Bis(thiophen-2-yl)quinoxaline-6-carboxylic acid [2-(4-pyridyl)ethyl]amide 832082-05-09, 2,3-Bis(thiophen-2-yl)quinoxaline-6-carboxylic acid N-[3-(phenylpropyl)amide]


RI: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USBS (Uses)

(preparation of quinoxaline and pyrido[2,3-b]pyrazine derive. as PKB inhibitors for treatment of cancers)


RN 443111-01-1 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[3-(4-morpholinyl)propyl]-2,3-di-2-thienyl- (9CI) (CA INDEX NAME)




RN 443111-36-2 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(2-chlorophenyl)ethyl]-2,3-di-2-thienyl- (9CI) (CA INDEX NAME)




RN 443111-45-3 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(4-morpholinyl)ethyl]-2,3-di-2-thienyl- (9CI) (CA INDEX NAME)




RN 832082-01-6 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(2-methylphenyl)ethyl]-2,3-di-2-thienyl- (9CI) (CA INDEX NAME)




RN 832082-02-7 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(2-pyridinyl)ethyl]-2,3-di-2-thienyl- (9CI) (CA INDEX NAME)




RN 832082-03-8 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(3-pyridinyl)ethyl]-2,3-di-2-thienyl- (9CI) (CA INDEX NAME)




RN 832082-04-9 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(4-pyridinyl)ethyl]-2,3-di-2-thienyl- (9CI) (CA INDEX NAME)




RN 832082-05-0 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[3-(phenylpropyl)-2,3-di-2-thienyl- (9CI) (CA INDEX NAME)




RN 832080-84-9 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-chloro-N-[2-(2-chlorophenyl)ethyl]-3-(2-thienyl)- (9CI) (CA INDEX NAME)





RN 832080-85-0 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(2-chlorophenyl)ethyl]-2-[[2-(4-pyridinyl)ethyl]amino]-3-(2-thienyl)- (9CI) (CA INDEX NAME)




RN 832080-86-1 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(2-chlorophenyl)ethyl]-2-(methyl(phenylmethyl)amino)-3-(2-thienyl)- (9CI) (CA INDEX NAME)




RN 832082-00-5 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(2-methoxyphenyl)ethyl]-2,3-di-2-thienyl- (9CI) (CA INDEX NAME)




L13 ANSWER 23 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2005:11522 CAPLUS
DOCUMENT NUMBER: 142:26107
TITLE: Preparation and redox properties of N,N,N-1,3,5-trialkylated flavin derivatives and their activity as redox catalysts
AUTHOR(S): Linden, Auri A.; Hermanns, Nina; Ott, Sascha; Krueger, Lars; Baackvall, Jan-E.
CORPORATE SOURCE: Department of Organic Chemistry, Stockholm University, Stockholm, 106 91, Swed.
SOURCE: Chemistry--A European Journal (2005), Volume Date 2004, 11(1), 112-119
CODEN: CEJUED; ISSN: 0947-6539
PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA
DOCUMENT TYPE: Journal
LANGUAGES: English
OTHER SOURCE(S): CASREACT 142:261107
AB Eight different flavin derive. have been synthesized and the electronic effects of substituents in various positions on the flavin redox chemical were investigated. The redox potentials of the flavins, determined by cyclic voltammetry, correlated with their efficiency as catalysts in the H2O2 oxidation of Me p-tolyl sulfide. Introduction of electron-withdrawing groups increased the stability of the reduced catalyst precursor.
IT 845753-36-0P 845753-39-1P 845753-44-0P 845753-46-0P
RL: PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and redox properties of N,N,N-1,3,5-trialkylated flavin derive. and activity as redox catalysts)
RN 845753-36-8 CAPLUS
CN Benzo[9]pteridine-7-carboxylic acid, 1,2,3,4-tetrahydro-1,3-dimethyl-2,4-dioxo-, butyl ester (9CI) (CA INDEX NAME)



RN 845753-39-1 CAPLUS
CN Benzo[5,6-b]pyridine-7-carboxylic acid, 1,2,3,4-tetrahydro-2,4-dioxo-, butyl ester (9CI) (CA INDEX NAME)



RN 845753-44-8 CAPLUS
CN Benzo[5,6-b]pyridine-8-carboxylic acid, 1,2,3,4-tetrahydro-2,4-dioxo-, butyl ester (9CI) (CA INDEX NAME)




RN 845753-46-0 CAPLUS
CN Benzo[5,6-b]pyridine-8-carboxylic acid, 1,2,3,4-tetrahydro-1,3-dimethyl-2,4-dioxo-, butyl ester (9CI) (CA INDEX NAME)

REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RS FORMAT

L13 ANSWER 23 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2004:1154708 CAPLUS
DOCUMENT NUMBER: 142:93843
TITLE: Preparation of pyrido[1,2-a]pyrimidin-4-ones as anticancer agents
INVENTOR(S): Wang, Weibo; Constantine, Ryan N.; Lagriton, Liana M.;

PATENT ASSIGNEE(S): Pecchi, Sabina; Burger, Matthew T.; Desai, Manoj C.
SOURCE: Chiron Corporation, USA
PCT Int. Appl., 78 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
MO 2004113335	A2	20041229	MO 2004-US19158	20040617
MO 2004113335	A3	20050324		
N: AR, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RN: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2004249730	A1	20041229	AU 2004-249730	20040617
CA 2528771	AA	20041229	CA 2004-2528771	20040617
US 2005085490	A1	20050421	US 2004-870707	20040617
EP 1636225	A2	20060322	EP 2004-776639	20040617
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IS, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK				
PRIORITY APPLN. INFO.: US 2003-480180P P 20030620 MO 2004-US19158 W 20040617				
OTHER SOURCE(S): MARPAT 142:93843 OI				



AB The title compds. I [R1 = H, alkyl, aryl, etc.; R2, R3 = H, alkyl, aryl, etc.; or R2 and R3 taken together with the carbon atom to which they are attached form a 3-7 membered carbocyclic or heterocyclic ring; R4 = H, alkyl, aryl, etc.; R5 = H, alkyl, aryl, etc.; R6-R9 = H, halo, NO2, etc.], useful, either alone or in combination with at least one addnl. therapeutic agent, in the prophylaxis or treatment of proliferative diseases, were prepared. E.g., a multi-step synthesis of II, starting from 2-aminopyridine and Et 4-chloroacetate, was given. Certain compds. I were shown to have a KSP inhibitory activity at an IC50 of less than about

25 µM. The compds. that include a pharmaceutically acceptable carrier and one or more of the pyrido[1,2-a]pyrimidinyl compds. I, either alone or in combination with at least one addnl. therapeutic agent, were disclosed.
IT 817205-84-8P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PRP (Preparation); USES (Uses)
(preparation of pyrido[1,2-a]pyrimidin-4-ones as anticancer agents)

RN 817205-84-8 CAPLUS
CN 6-Quinoxalinecarboxamide, N-(3-aminopropyl)-N-[1-[4-oxo-3-(phenylmethyl)-4H-pyrido[1,2-a]pyrimidin-2-yl]propyl]- (9CI) (CA INDEX NAME)




L13 ANSWER 24 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2004:976488 CAPLUS
DOCUMENT NUMBER: 142:110137
TITLE: Immunochemical detection of 3-deoxyglucosone in serum
AUTHOR(S): Uchida, Yoshiaki; Kurano, Yoshihiro; Endo, Tomohiro; Aoyama, Misao; Ito, Satoru
CORPORATE SOURCE: Research and Development Division, Fujirebio Inc., Machiej, Tokyo, 193-0031, Japan
SOURCE: Biochemical and Biophysical Research Communications (2004), 325(3), 1090-1098
CODEN: BBRCA9; ISSN: 0006-291X
PUBLISHER: Elsevier
DOCUMENT TYPE: Journal
LANGUAGE: English

AB 3-Deoxyglucosone (3-DG) is a metabolite of glucose that is thought to lead to the production of advanced glycation end products in diabetes. The previous assay for 3-DG in serum was based on a multi-step protocol, including derivatization, extraction, HPLC separation, and detection. In the current studies, we established a monoclonal antibody that recognizes the 3-DG-derivative, which is generated by the reaction of 3-DG and a 2,3-diamino-benzene derivative. Attachment of a biotin moiety to the 2,3-diamino-benzene ring via a linker allowed development of a highly sensitive chemiluminescent enzyme immunoassay for 3-DG equivalent. Unlike the previous assay, this method does not require extraction of 3-DG derivative from serum. Treatment of 3-DG in serum with the DAB-link-biotin produced a quinoxaline derivative, which was specifically recognized by the monoclonal antibody. Using this assay, we found that serum 3-DG was higher in streptozotocin-induced diabetic rats than in normal control rats (25±5.6 vs. 9.8±1.1 µg/L). This simple assay may allow the monitoring of conditions leading to the accumulation of advanced glycation end products and evaluation of the risk of complications in diabetic patients.


IT 824960-70-5
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(serum 3-deoxyglucosone is higher in streptozotocin-induced diabetic rats)

RN 824960-70-5 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[4-oxo-4-[(2-(2-pyridinylthio)ethyl)amino]butyl]-3-(2,3,4-trihydroxybutyl)- (9CI) (CA INDEX NAME)



IT 824960-65-8 824960-66-9 824960-67-0
824960-68-1 825637-48-7
RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
(serum 3-deoxyglucosone is higher in streptozotocin-induced diabetic rats)
RN 824960-65-8 CAPLUS
CN L-Lysine, N2-[6-[[5-[[3aS,4S,6aR]-hexahydro-2-oxo-1H-thieno[3,4-d]imidazol-4-yl]-1-oxopentyl]amino]-1-oxohexyl]-N6-[[2-(2,3,4-trihydroxybutyl)-6-quinoxaliny]carbonyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 824960-66-9 CAPLUS
CN L-Lysine, N2-[6-[[5-[[3aS,4S,6aR]-hexahydro-2-oxo-1H-thieno[3,4-d]imidazol-4-yl]-1-oxopentyl]amino]-1-oxohexyl]-N6-[[2-(2,3,4-trihydroxybutyl)-6-quinoxaliny]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




PAGE 1-B




RN 824960-67-0 CAPLUS
CN L-Lysine, N2-[6-[[[5-[(3aS,4S,6aR)-hexahydro-2-oxo-1H-thieno[3,4-d]imidazol-4-yl]-1-oxopentyl]amino]-1-oxohexyl]-N6-[(2-methyl-6-quinoxaliny]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




PAGE 1-B




RN 824960-68-1 CAPLUS
CN L-Lysine, N6-[[[2-(2,3-dihydroxypropyl)-6-quinoxaliny]carbonyl]-N2-[6-[[[5-[(3aS,4S,6aR)-hexahydro-2-oxo-1H-thieno[3,4-d]imidazol-4-yl]-1-oxopentyl]amino]-1-oxohexyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




PAGE 1-B




RN 825637-48-7 CAPLUS
CN L-Lysine, N2-[6-[[[5-[(3aS,4S,6aR)-hexahydro-2-oxo-1H-thieno[3,4-d]imidazol-4-yl]-1-oxopentyl]amino]-1-oxohexyl]-N6-[[[2-[(1R,2S,3R)-1,2,3,4-tetrahydroxybutyl]-6-quinoxaliny]carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 25 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2004:927005 CAPLUS
DOCUMENT NUMBER: 141:395806
TITLE: Preparation of quinoxaliny macrocyclic hepatitis C
serine protease inhibitors
INVENTOR(S): Nakajima, Suanne; Sun, Ying; Tang, Datong; Xu, Gouyou;
Porter, Brian; Or, Yat Sun; Wang, Zhe; Miao, Zhenwei
PATENT ASSIGNEE(S): Enanta Pharmaceuticals, Inc., USA
SOURCE: PCT Int. Appl., 131 pp.
CODEN: PIXK22
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004093798	A2	20041104	WO 2004-US11841	20040416
WO 2004093798	A3	20051208		
W: AB, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MG, MK, MN, MU, MZ, NA, NI, NO, NZ, OM, PA, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, OH, OM, KE, LS, MW, MZ, SD, SI, SE, TZ, UG, ZM, ZW, AM, AE, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2004231987	A1	20041104	AU 2004-231987	20040416
CA 2522561	AA	20041104	CA 2004-2522561	20040416
US 2004266668	A1	20041230	US 2004-826743	20040416
EP 1615613	A2	20060118	EP 2004-750236	20040416
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
PRIORITY APPL. INFO.: US 2003-418759 A 20030418 US 2003-509071P P 20030418 WO 2004-US11841 W 20040416				

OTHER SOURCE(S): MARPAT 141:395806
OI



AB The invention relates to macrocyclic compds. I [A is H, CO2R1, CO2R2,

CONHR2, CSNHR2 or SO2R2; G is OH, alkoxy, NHSO2R1, COR2, CO2R1 or CONHR2; L is S, SCH2, SO2, O, COCH2, CHMeCH2, etc.; m, n = 0-2; p = 0-4; R2 is a bond or H2; R1 is H, (un)substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl or heterocycloalkyl; R2 is any group given for R1 or mono- or dialkylamino or -arylamino; R3, R4 not defined; X and Y taken together with the carbon atoms to which they are attached form (un)substituted aryl or heteroaryl; W is absent, O, S, NH, C(O)NR1 or NR1; Z is H, -CN, -SCN, -NCO, -NCS, NHNH2, N3, halo, cycloalkyl, aryl, etc.) or their pharmaceutically-acceptable salts, esters or prodrugs which inhibit serine protease activity, particularly the activity of hepatitis C virus (HCV) NS3-NS4a protease. The compds. of the invention interfere with the life cycle of the hepatitis C virus and are also useful as antiviral agents. Thus, macrocycle II (Boc = tert-butoxycarbonyl) was prepared via peptide coupling and ring-closing metathesis reactions.


IT 787600-64-0P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USRS (Uses)



(preparation of quinoxaliny cyclic peptides as hepatitis C serine protease inhibitors)

RN 787600-64-0 CAPLUS
CN Cyclopropa[e]pyrrolo[1,2-a][1,4]diazacyclopentadecine-14a(5H)-carboxylic acid, 6-[[[(1,1-dimethylethoxy)carbonyl]amino]-1,2,3,6,7,8,9,10,11,13a,14,15,16,16a-tetradecahydro-5,16-dioxo-2-[[6-[[[phenylmethyl]amino]carbonyl]-3-(2-thienyl)-2-quinoxalinyloxy]-, (2R,6S,13aS,14aR,16aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry unknown.

PAGE 1-A





L13 ANSWER 26 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2004:85739 CAPLUS
 DOCUMENT NUMBER: 141:343478
 TITLE: Use of small molecule compounds for immunopotentialization
 INVENTOR(S): Valiente, Nicholas
 PATENT ASSIGNEE(S): Chiron Corporation, USA
 SOURCE: PCT Int. Appl., 146 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004087153	A2	20041014	WO 2004-US10331	20040329
WO 2004087153	A3	20050317		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GR, GU, HK, HN, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: BW, GH, GM, KE, LS, MW, MG, SD, SE, SL, SZ, TG, TZ, UG, ZM, ZW, AM, AZ, BY, BG, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, FR, GB, GR, GU, HK, HN, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

CA 2520124 AA 20041014 CA 2004-2520124 20040329
 US 2005136065 A1 20050623 US 2004-814480 20040329
 SP 1608369 A2 20051228 SP 2004-785593 20040329
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK
 PRIORITY APPLN. INFO.: US 2003-458888 P 20030328
 WO 2004-US10331 W 20040329


OTHER SOURCE(S): MARPAT 141:343478
 GI



AB The invention provides immunostimulatory compns. comprising a small mol. immunopotentiator (SMIP) compound and methods of administration thereof. Also provided are methods of administering a SMIP compound in an effective amount to enhance the immune response of a subject to an antigen. Further provided are compns. and methods of administering SMIP compds. alone or in combination with another agent for the treatment of cancer, infectious diseases and/or allergies/asthma. Preparation of selected compds., e.g. I, is included.

IT 854634-24-9
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (small mol. compds. for immunopotentialization)

RN 654634-24-9 CAPLUS
 CN Benzo[b]phenazine-2-carboxylic acid, 6,11-dihydro-6,11-dioxo-, ethyl ester (9CI) (CA INDEX NAME)




L13 ANSWER 27 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2004:791916 CAPLUS
 DOCUMENT NUMBER: 142:3310
 TITLE: Optimizing the antibacterial activity of a lead structure discovered by SAR by MS technology
 AUTHOR(S): Elisabeth A.; Geth, Punit P.; Robinson, Dale E.; Winter, Dana K.; Miyaji, Alysia; Riesen, Lisa

CORPORATE SOURCE: M.; Osgood, Stephen A.; Bertrand, Myra; Swayze, Eric E.
 Ibis Therapeutics, Isis Pharmaceuticals, Inc., Carlsbad, CA, 92008, USA
 SOURCE: Bioorganic & Medicinal Chemistry Letters (2004), 14(21), 5257-5261
 CODEN: BMCLSE; ISSN: 0960-894X
 PUBLISHER: Elsevier B.V.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 142:3310

AB We report on lead optimization of a compound that was originally discovered to bind bacterial 23S rRNA near the L11 binding site and inhibit translation in vitro, but lacked detectable antibacterial activity. In this study, we were able to generate compds. with antibacterial activity against Gram-neg. and Gram-pos. pathogens, including a methicillin-resistant *Staphylococcus aureus* strain.

IT 797770-86-6P
 RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PRSP (Preparation)
 (optimizing the antibacterial activity of a lead structure)
 RN 797770-86-6 CAPLUS
 CN 6-Quinoxalinecarboxamide, N-[1-[4-[3-(dimethylamino)propoxy]phenyl]-2-oxo-2-(1-piperazinyl)ethyl]-1,2,3,4-tetrahydro-2,3-dioxo- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L13 ANSWER 28 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2004:515470 CAPLUS
 DOCUMENT NUMBER: 141:71152
 TITLE: Preparation of biphenylaminobenzoates and related compounds as modulators of peroxisome proliferator activated receptor γ (PPAR γ) type receptors as drugs and cosmetics.
 INVENTOR(S): Clary, Laurence; Collette, Pascal; Rivier, Michel; Jonard, Andre
 PATENT ASSIGNEE(S): Galderma Research & Development, S.N.C., Fr.
 SOURCE: PCT Int. Appl., 90 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004052840	A1	20040624	WO 2003-EP15010	20031111

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GR, GU, HK, HN, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW


LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: BW, GH, GM, KE, LS, MW, MG, SD, SE, SL, SZ, TG, TZ, UG, ZM, ZW, AM, AZ, BY, BG, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, FR, GB, GR, GU, HK, HN, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 FR 2848553 A1 20040618 FR 2002-15751 20021212
 CA 2506732 AA 20040624 CA 2003-2506732 20031211
 AU 2003102909 A1 20040630 AU 2003-102909 20031211
 EP 1572629 A1 20050914 EP 2003-068292 20031211
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK
 BR 2003016193 A 20050927 BR 2003-16193 20031211
 JP 2006050979 T2 20060323 JP 2004-558089 20031211
 US 2006009484 A1 20060112 US 2005-149551 20050610
 PRIORITY APPLN. INFO.: FR 2002-15751 A 20021212
 WO 2003-434322 P 20031219
 WO 2003-EP15010 W 20031211

OTHER SOURCE(S): MARPAT 141:71352
 GI




AB Title compds. {I: R1 = Q1, Q2, COR5; R2 = H, halo, OH, NO2, alkyl, alkoxy, polyether, amino, aryl, aralkyl, heteroaryl, heterocyclyl; R3 = (CH2)t(NR15)u(C(O,N))vR16, alkyl, aryl, aralkyl, heteroaryl, heterocyclyl, 9-fluorenylmethyl; R4 = H, alkyl, aryl, aralkyl, heteroaryl, heterocyclyl; R5 = OH, O(CH2)nR6, amino, etc.; R6 = aryl, aralkyl, heteroaryl, heterocyclyl, NHCO2R7, NHCO2R7, NR7R8; A = (CH2)m(NR10)p(CO)qR9, (CH2)n(NR10)p(CS)qR9; D = O, S, CH2, NR11; R8, R10, R11 = H, alkyl; X = O, S, CH2, NR9; R9 = H, alkyl, aralkyl; R7, R12, R15 = H, alkyl, aryl, aralkyl, heteroaryl, heterocyclyl; R16 = R15, NHCO2R7, NHCO2R7, NR7R8; V = O, S, NO; Q = H, alkyl; W = N, CR12; m, p, q, r = 0, 1; n = 1-3; t, u, v = 0-4; with proviso(s), were prepared. Thus, N-(4'-bromobiphenyl-3-ylmethyl)-N-methyl-6-(2-methoxyethoxymethyl)-2-carboxamide (preparation given), Pd(OAc)2, Me anthranilate, and CaCl2 were successively introduced into a solution of BINAP in PhMe followed by heating at 100° for 8 h to give 83 Me 2-[3'-[[[6-(2-methoxyethoxymethyl)naphthalene-2-carbonyl]methyl]amino]methyl]biphenyl-4-ylaminobenzoate. This was stirred 8 h with NaOH in THF/MeOH/H2O to give 504 2-[3'-[[[6-(2-methoxyethoxymethyl)naphthalene-2-carbonyl]methyl]amino]methyl]biphenyl-4-ylaminobenzoic acid. The latter in a crossover-curve PPAR transactivation test showed PPAR γ activity with Kd apparent = 30 nM.
 IT 706779-77-3P 706779-85-3P 706779-92-2P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PRSP (Preparation); USES (Uses)
 (claimed compound; preparation of biphenylaminobenzoates and related compds. as modulators of peroxisome proliferator activated receptor γ)


RN 706779-77-3 CAPLUS
CN Benzoic acid, 2-[[3'-[[methyl(6-quinoxalinylylcarbonyl)amino]methyl]](1,1'-biphenyl)-4-yl]amino]- (9CI) (CA INDEX NAME)



RN 706779-85-3 CAPLUS
CN Benzoic acid, 3-[[3'-[[methyl(6-quinoxalinylylcarbonyl)amino]methyl]](1,1'-biphenyl)-4-yl]oxy]-, 2-(4-morpholinyl)ethyl ester (9CI) (CA INDEX NAME)



RN 706779-92-2 CAPLUS
CN Benzoic acid, 3-[[3'-[[methyl(6-quinoxalinylylcarbonyl)amino]methyl]](1,1'-biphenyl)-4-yl]oxy]-, 2-(4-morpholinyl)ethyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 29 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2004:492324 CAPLUS
DOCUMENT NUMBER: 141:59705
TITLE: New amino biphenyl compounds as modulators of peroxisome proliferator-activated γ -receptors (PPAR γ) for cosmetic or pharmaceutical compositions

INVENTOR(S): Clary, Laurence; Collette, Pascal; Rivier, Michel;
Jomard, Andre
PATENT ASSIGNEE(S): Galderma Research & Development, Fr.
SOURCE: Fr. Demande, 47 pp.


DOCUMENT TYPE: CODEN: FRXXBL
LANGUAGE: Patent
FAMILY ACC. NUM. COUNT: French
PATENT INFORMATION: 2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2848553	A1	20040618	FR 2002-15751	20021212
CA 2506732	AA	20040624	CA 2003-2506732	20031211
WO 2004052840	A1	20040624	WO 2003-EP15010	20031211
W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MX, MY, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MM, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 200302909	A1	20040630	AU 2003-302909	20031211
EP 1572629	A1	20050914	EP 2003-088292	20031211
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003016193	A	20050927	BR 2003-16193	20031211
JP 2006050979	T2	20060323	JP 2004-558089	20031211
US 2006009484	A1	20060112	US 2005-149551	20050610
PRIORITY APPLN. INFO.: FR 2002-15751 A 20021212 US 2002-43482P P 20021219 WO 2003-EP15010 W 20031211				


OTHER SOURCE(S): MARPAT 141:59705
AB New amino biphenyl compds. as PPAR γ receptor modulators are prepared and used for cosmetics or pharmaceuticals compns. for the treatment of cardiovascular diseases, immune diseases and/or diseases related to the metabolism of the lipids. Thus, tablets comprised a biphenylaminobenzoic acid 0.001, starch 0.114, dicalcium phosphate 0.020, silica 0.020, lactose 0.030, talc 0.010, and Mg stearate 0.005 g. The synthesis of amino biphenyl compds. is described.

IT 706779-77-3P 706779-85-3P 706779-92-2P
RL: COS (Cosmetic use); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(amino biphenyl compds. as modulators of PPAR γ receptors for cosmetic or pharmaceutical compns.)


RN 706779-77-3 CAPLUS
CN Benzoic acid, 2-[[3'-[[methyl(6-quinoxalinylylcarbonyl)amino]methyl]](1,1'-biphenyl)-4-yl]amino]- (9CI) (CA INDEX NAME)



RN 706779-85-3 CAPLUS
CN Benzoic acid, 3-[[3'-[[methyl(6-quinoxalinylylcarbonyl)amino]methyl]](1,1'-biphenyl)-4-yl]oxy]-, 2-methylpropyl ester (9CI) (CA INDEX NAME)



RN 706779-92-2 CAPLUS
CN Benzoic acid, 3-[[3'-[[methyl(6-quinoxalinylylcarbonyl)amino]methyl]](1,1'-biphenyl)-4-yl]oxy]-, 2-(4-morpholinyl)ethyl ester (9CI) (CA INDEX NAME)



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L13 ANSWER 30 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2004:467877 CAPLUS
DOCUMENT NUMBER: 141:38617
TITLE: Preparation of biphenyls which activate peroxisome proliferator activated receptor- γ (PPAR γ) receptors for use in drugs and cosmetics.

INVENTOR(S): Clary, Laurence; Bouix-Peter, Claire; Rivier, Michel;
Collette, Pascal; Jomard, Andre
PATENT ASSIGNEE(S): Galderma Research & Development, S.N.C., Fr.
SOURCE: PCT Int. Appl., 73 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004048351	A2	20040610	WO 2003-EP15002	20031121
WO 2004048351	A3	20040812		
W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MM, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				


ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
FR 2847580	A1	20040528	FR 2002-14793	20021125
FR 2847580	B1	20060303		
CA 2506523	AA	20040610	CA 2003-2506523	20031121
AU 2003294025	A1	20040618	AU 2003-294025	20031121
EP 1567509	A2	20050831	EP 2003-789439	20031121
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003015951	A	20050913	BR 2003-15951	20031121
US 2006004048	A1	20060105	US 2005-135499	20050524
PRIORITY APPLN. INFO.: FR 2002-14793 A 20021125 US 2002-430698P P 20021204 WO 2003-EP15002 W 20031121				

OTHER SOURCE(S): MARPAT 141:38617
GI



AB Title compds. [I; R1 = alkyl, (substituted) Ph, N-protected α -amino acid, etc.; R2 = Q1, Q2, COR3; R3 = H, halo, alkyl, OH, alkoxy, aralkoxy, aryloxy, polyether residue, NO2, amino; R4 = alkyl, aryl, aralkyl, heteroaryl, heterocyclyl, etc.; A = (CH2)_n(NR12)2(CO)2W, etc.; W, X, Y, Z = O, 1; D = O, E, CH2, NR13; V = O, N, S; W = N, CR13; Y = N, C; R7 = H, alkyl, aryl, aralkyl, heteroaryl, heterocyclyl; R8 = O(CH2)_nVR9, OH, alkoxy, aralkoxy, aryloxy, aryl, aralkyl, heteroaryl, heterocyclyl, etc.; R9 = aryl, aralkyl, heteroaryl, heterocyclyl, etc.; R10 = H, alkyl, aryl, aralkyl, heteroaryl, heterocyclyl; R12, R13 = H, alkyl; R14 = halo; v = 1-3; were prepared. Thus, HATU, PS-carbodiimide resin, 4'-[2-ethoxy-2-(5-propyl)-1,2,4-oxadiazol-2-yl]ethyl]biphenyl-3-ylmethyl]methanamine (preparation given), and 6-(2-methoxyethoxymethoxy)naphthalene-2-carboxylic acid were stirred together in DMF/CH2Cl2 for 18 h at room temperature to give a residue which was stirred 5 h with MP-carbonate resin in DMF/CH2Cl2 to give 100% N-[4'-[2-ethoxy-2-(5-propyl)-1,2,4-oxadiazol-2-yl]ethyl]biphenyl-3-ylmethyl]-N-methyl-6-(2-methoxyethoxymethoxy)naphthalene-2-carboxamide. In a crossed-curve PPAR γ activation test, the latter showed Kd app = 250 nM.

IT 692780-94-2P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(claimed compound: preparation of biphenyls which activate peroxisome proliferator activated receptor- γ receptors for use in drugs and cosmetics)
RN 692780-94-2 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[4'-[2-ethoxy-2-(4-morpholinyl)-3-oxopropyl]](1,1'-biphenyl)-3-yl]methyl]-N-methyl- (9CI) (CA INDEX NAME)



L13 ANSWER 31 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2004:433750 CAPLUS

DOCUMENT NUMBER:
141:7131

TITLE: Preparation of quinazolines and analogs as Akt inhibitors and indoles as protein kinase inhibitors for use in synergistic combination therapy for the treatment of cancer

INVENTOR(S): Barnett, Stanley F.; Defeo-Jones, Deborah D.; Hartman, George D.; Huber, Hans E.; Stirdivant, Steven M.; Heimbrook, David C.

PATENT ASSIGNER(S): USA
SOURCE: U.S. Pat. Appl. Publ., 121 pp., which

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004102360	A1	20040527	US 2003-678565	20031003
PRIORITY APPLN. INFO.:			US 2002-422312P	P 20021030
			US 2003-460911P	P 20030407

OTHER SOURCE(S):
GI


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The present invention relates to methods of treating cancer using a combination of at least two Akt inhibitors I [wherein Q = (un)substituted heterocyclyl, aryl; U, V, W, and X = independently CH, N; Y, Z = independently CH, N, provided that at least one of Y and Z = N; n = 0-3; p = 0-2; q = 0-4; R1, R2, R3, R4 = independently halo, CN, OH, CHO, NO2, or (un)substituted (cyclo)alkyl(oxy), alkenyl(oxy), alkynyl(oxy), heterocyclyl(oxy), acyl, carboxy, carbamoyl(oxy), ureido, sulfamoyl, etc.; R3, R4 = independently H, (perfluoro)alkyl; or CR3R4 = cycloalkyl, heterocyclyl; and pharmaceutically acceptable salts or stereoisomers thereof] or a combination of I and a protein kinase inhibitor II [wherein Q = H2, O; X = C, N, SO2-2, O; m = 0-2; p = 0-2; q = 0-4; R1 = independently H, halo, or (un)substituted (cyclo)alkyl, heterocyclyl, aryl, carbamoyl, amino, acyl, sulfamoyl, carboxy, etc.; R2 = H or (un)substituted (cyclo)alkyl(oxy), amino, aryloxy, heterocyclyloxy, alkenyloxy, alkynyloxy, etc.; R5 = independently H, halo, NO2, CN, or (un)substituted alkyl, alkenyl, alkynyl, carboxy, acyl, sulfamoyl, carbamoyl, ureido, amino, etc.; and pharmaceutically acceptable salts or stereoisomers thereof], optionally in combination with a third compound. Examples include syntheses for I and II and assays demonstrating Akt inhibitor activity, antitumor activity, and the synergistic effect of combinations of Akt inhibitors and/or protein kinase inhibitors on caspase 3 activity. For instance, III-HCl was prepared in an 8-step reaction sequence culminating with the cyclization of 4-(2-aminoprop-2-yl)benzil and o-phenylenediamine using glacial acetic acid in H2O, followed by work up

with chloroform and ethanolic HCl. III-HCl, a selective Akt1 and Akt2 inhibitor, demonstrated a 3.2-fold in caspase 3 activation over control compared to a 1.2-fold increase for a protein kinase inhibitor. Combination treatment produced a 9-fold increase in caspase 3 activation.
IT 612847-33-3P 612847-34-4P 612847-42-4P
612847-43-5P 612848-56-3P 612848-57-4P
612848-59-6P 612848-61-0P 695816-06-9P
RL: PAC (Pharmacological activity); SPH (Synthetic preparation); THU (Therapeutic use); BLOL (Biological study); PREP (Preparation); USES (Uses)

(antitumor agent; preparation of quinazolines and analogs as Akt inhibitors and indoles as protein kinase inhibitors for use in synergistic combination therapy for treatment of cancer)

RN 612847-33-3 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl- (9CI) (CA INDEX NAME)



PAGE 1-A


PAGE 1-B



RN 612847-34-4 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-, bis(trifluoroacetate) (9CI) (CA INDEX NAME)

CM 1

CRN 612847-33-3
CMF C40 H38 N8 O2



PAGE 1-A

PAGE 1-B




CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 612847-42-4 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl- (9CI) (CA INDEX NAME)



RN 612847-43-5 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl- (9CI) (CA INDEX NAME)




RN 612848-56-3 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 612847-43-5
CMF C40 H43 N7 O2



CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 612848-57-4 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 612847-42-4
CMF C40 H43 N7 O2



CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 612848-59-6 CAPLUS

CN D-arabino-Hexose, 2-deoxy-2-[[[2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl]-6-quinoxaliny]carbonyl]amino]-, (2S)-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1


CRN 612848-58-5
CMF C40 H40 N6 O7

Absolute stereochemistry.



PAGE 1-A

PAGE 1-B



CM 2

CRN 76-05-1
CMF C2 H F3 O2



RN 612848-61-0 CAPLUS


CN D-arabino-Hexose, 2-deoxy-2-[[[3-[4-[[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl-6-quinoxaliny]carbonyl]amino]-, (2S)-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1


CRN 612848-60-9
CMF C40 H40 N6 O7

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 695816-06-9 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-2-[4-[[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-methyl-3-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 695816-05-8
CMF C41 H45 N7 O2



CM 2

CRN 76-05-1
CMF C2 H F3 O2



L13 ANSWER 33 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2004:432769 CAPLUS

DOCUMENT NUMBER: 140:429035

TITLE: Preparation of biaromatic compounds as PPAR receptors and their use in cosmetic or pharmaceutical compositions

INVENTOR(S): Clary, Laurence; Bouix, Peter Claire; Rivier, Michel; Collette, Pascal; Jomard, Andre

PATENT ASSIGNER(S): Galderma Research & Development, Fr.

SOURCE: Fr. Demande, 45 pp.

CODEN: FRXXBL

DOCUMENT TYPE: Patent

LANGUAGE: French


FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2847580	A1	20040528	FR 2002-14793	20021125
FR 2847580	B1	20060303		
CA 2506523	AA	20040610	CA 2003-2506523	20031121
WO 2004048351	A2	20040610	WO 2003-EP15002	20031121
WO 2004048351	A3	20040812		
W: AB, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HK, HN, ID, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, OH, OM, KE, LS, MW, KE, SD, SE, SE, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003294025	A1	20040618	AU 2003-294025	20031121
EP 1567509	A2	20050831	EP 2003-789439	20031121
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, SI, ST, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003015951	A	20050913	BR 2003-15951	20031121
US 2006004048	A1	20060105	US 2005-135499	20050524
PRIORITY APPL. INFO.: FR 2002-14793 A 20021125				
US 2002-430698P P 20031204				
WO 2003-EP15002 N 20031121				

OTHER SOURCE(S): MARPAT 140:429035

Q1



I


AB The invention relates to new biarom. compds. I and their method of preparation, and their use in cosmetic or pharmaceutical compns. intended for use in human or veterinary medicine (such as cardiovascular diseases, immunity diseases and/or diseases related to the metabolism of the lipids). Thus, N-[(4'-[2-ethoxy-2-(5-propyl-1,3,4)oxadiazol-2-yl]-ethyl]-biphenyl-3-dimethyl-N-methyl-4-phenoxy-benzamide (II) was prepared by the reaction of 4-Phenoxy-benzoic acid with [4'-[2-ethoxy-2-(5-propyl-5 [1,3,4)oxadiazol-2-yl]-ethyl]-biphenyl-3-ylmethyl]-methyl-amine. The specific affinity of the composition for PPAR- γ is shown. A tablet contained II 0.001, starch 0.114, dicalcium phosphate 0.020, silica 0.020, lactose 0.030, talc 0.010, and magnesium stearate 0.005 g.

IT 692780-94-2

RL: COS (Cosmetic use); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) Preparation of biarom. compds. as activators of PPAR receptors and their uses in cosmetic or pharmaceutical compns.]

RN 692780-94-2 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[[4'-[2-ethoxy-3-(4-morpholinyl)-3-oxopropyl][1,1'-biphenyl]-3-yl]methyl]-N-methyl- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 33 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2004:430796 CAPLUS

DOCUMENT NUMBER: 141:7139

TITLE: Preparation of indolylquinoxalinones for treating hyperproliferative disorders and diseases associated with angiogenesis

INVENTOR(S): Ladouceur, Gaetan H.; Bear, Brian; Bi, Cheng; Brittelli, David R.; Burke, Michael J.; Chen, Gang; Cook, James; Dumas, Jacques; Sibley, Robert; Turner, Michael R.

PATENT ASSIGNER(S): Bayer Pharmaceuticals Corporation, USA

SOURCE: PCT Int. Appl., 217 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE

MO 2004043950 A1 20040527 MO 2003-US16003 20031110

W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GR, GU, GM, HR, HU, ID, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MG, MK, MN, MW, MX, MY, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SN, TD, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RM: BW, GH, GM, KE, LS, MM, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, NG, SN, TD, TG

CA 2505819 AA 20040527 CA 2003-2505819 20031110

AU 2003290744 A1 20040603 AU 2003-290744 20031110

EP 1565455 A1 20050824 EP 2003-783328 20031110

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, NL, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, NG, SN, TD, TG

BR 2003016169 A 20050927 BR 2003-16169 20031110

CN 1738814 A 20060222 CN 2003-80108639 20031110

JP 2006050840 T2 20060323 JP 2005-507146 20031110

US 2006040111 A 20060105 US 2005-534215 20050506

NO 2005002796 A 20050609 NO 2005-2796 20050609

PRIORITY APPL. INFO.: US 2002-425490P P 20021112


US 2003-460915P P 20030407

US 2003-484202P P 20030630

MO 2003-US16003 W 20031110

OTHER SOURCE(S): MARPAT 141:7139

GI



AB The invention relates to title compds. I [wherein Ar = 6-membered aromatic ring containing 0-2 N atoms; R1 and R2 = independently H, halo, CF3, acyl, piperidinyl, piperazinyl, morpholinyl, or (un)substituted alkyl, alkoxy, amino, pyrrolidinyl, Ph, etc.; R3 = H, alkyl, OH, NO2, NH2, alkylamino, sulfonylamino, or (un)substituted benzoylamino; R4 = H, OH, halo, CN, acyl, alkylamino, trialkylethoxy, tetrazolyl, thienyl, pyrrolyl, pyrimidinyl, oxazolyl, furanyl, or (un)substituted alkyl, alkenyl, alkynyl, alkoxy, amino, oxadiazolyl, Ph, pyridyl, (oxy), carbonyl; R11 and R12 = independently H, F, or Cl with the proviso that when one of R11 and R12 = F or Cl, the other must be H; and pharmaceutically acceptable salts and esters thereof]. The invention also relates to the use of I and their pharmaceutical compns. for treating hyperproliferative disorders and diseases associated with angiogenesis (no data). Examples include representative syntheses for compds. of the invention, pharmaceutical compns. comprising them, and tumor model assays (no specific data given). For instance, N-Boc-indole was coupled with di-Me oxalate using t-BuLi to give tert-Bu 2-[methoxy(oxo)acetyl]-1H-indole-1-carboxylate (72%). Cyclization of the dione with 1,2-phenylenediamine in AcOH afforded the quinoxalinone II (77%).


IT 694331-65-29

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(antiproliferative and angiogenesis inhibitor; preparation of indolylquinoxalinones for treating hyperproliferative disorders and diseases associated with angiogenesis)

RN 694531-65-2 CAPLUS

CN 6-Quinoxalinecarboxamide, 3-(3-amino-1H-indol-2-yl)-1,2-dihydro-N-(2-methoxyethyl)-N-methyl-2-oxo- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 34 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2004:379708 CAPLUS

DOCUMENT NUMBER: 141:322446

TITLE: Picosecond time-resolved infrared investigation into the nature of the lowest excited state of fac-[Re(III)(CO)3(CO2Et-dppp)] [CO2Et-dppp = dipyrro[3,2a:2',3']c]phenazine-11-carboxylic ethyl ester)

AUTHOR(S): Kuimova, Marina K.; Grilla, David C.; Matousek, Pavel; Parker, Anthony W.; Sun, Xue-Zhong; Towrie, Michael; George, Michael W.

CORPORATE SOURCE: School of Chemistry, University of Nottingham, Nottingham, NG7 2RD, UK

SOURCE: Vibrational Spectroscopy (2004), 35(1-2), 219-223

CODEN: VISPEK; ISSN: 0924-2031

PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The photophysics of the fac-[Re(III)(CO)3(CO2Et-dppp)] [CO2Et-dppp = dipyrro[3,2a:2',3']c]phenazine-11-carboxylic ethyl ester) was studied with picosecond time-resolved IR (ps-TRIR) spectroscopy in metal carbonyl (2100-1800 cm⁻¹) and organic ester (1800-1600 cm⁻¹) spectral regions. The ps-TRIR spectra in both regions give evidence for the formation of a metal-to-ligand charge transfer (MLCT) excited state. The magnitude of ν(C=O) shift in the metal carbonyl region of the excited state relative to those of the ground state allows the excited state to be described more precisely as a dπ(Re) → π*(phenazine) MLCT state.


IT 767350-96-9P

RL: PREP (Physical, engineering or chemical process); PRP (Properties); DYP (Physical process); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)

(photophysics of tricarbonylchloro(dipyrro[phenazinecarboxylic ethyl ester])rhenium complex studied with picosecond time-resolved IR spectroscopy)

RN 767350-96-9 CAPLUS

CN Rhenium, tricarbonylchloro(ethyl dipyrro[3,2a:2',3']c]phenazine-11-carboxylate-κN4,κN5)-, (OC-6-44)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 35 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2004:372867 CAPLUS

DOCUMENT NUMBER: 140:375191

TITLE: Preparation of heteroaryl-hexanoic acid amides which are CCRI antagonists useful as immunomodulatory agents

INVENTOR(S): Brown, Matthew P.; Gaveco, Anderson S.; Gladue, Ronald P.; Kath, John C.; Posa, Christopher S.

PATENT ASSIGNEE(S): Pfizer Inc, USA

SOURCE: U.S. Pat. Appl. Publ., 63 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE

US 2004087571 A1 20040506 US 2003-687015 20031016

MO 2004039375 A1 20040513 MO 2003-184614 20031020

W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GR, GU, GM, HR, HU, ID, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MG, MK, MN, MW, MX, MY, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RM: GH, GM, KE, LS, MM, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, NG, SN, TD, TG


AU 2003267800 A1 20040525 AU 2003-267800 20031020

PRIORITY APPL. INFO.: US 2002-425799 P 20021030

MO 2003-184614 W 20031020

OTHER SOURCE(S): MARPAT 140:375191

GI



AB The title compds. II: R1 = (un)substituted heteroaryl; R2 = (un)substituted phenyl; R3 = (un)substituted naphthyl; Y = (un)substituted alkyl or heteroaryl-(CH2)m-; m = 0-4; R3 = H, (un)substituted alkyl.

cycloalkyl-(CH2)n-, heterocycloalkyl-(CH2)n-, heteroaryl-(CH2)n-, aryl-(CH2)n-; n = 0-6; R3 and the carbon to which it is attached form (un)substituted and/or fused 5-7 membered carbocyclic ring; Y = heteroaryl, heterocycloalkyl, (un)substituted H2N-sulfonyl, C(X)NH2; X = O, S, (un)substituted NH; R4 = H, alkyl, OH, alkoxy, hydroxyalkyl, alkoxyCO, cycloalkyl-(CH2)p-, (un)substituted heterocycloalkyl-(CH2)p-, heteroaryl-(CH2)p-, phenyl-(CH2)p- or naphthyl-(CH2)p-; p = 0-4 which are CCRI antagonists useful as immunomodulatory agents, were prepared e.g. a multi-step synthesis of quinoxaline-2-carboxylic acid [1(S)-benzyl-4(R)-benzylcarbamoyl-7-fluoro-2(S)-hydroxy-7-methyloctylamide, was given. All of the compds. I that were tested showed IC50 of <25 μM in the chemotaxis assay.

IT 212789-54-3P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of heteroaryl-substituted hexanamides as CCRI antagonists useful as immunomodulatory agents)

RN 212789-54-3 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[(1S,2S,4R)-4-(aminocarbonyl)-7-fluoro-2-hydroxy-7-methyl-1-(phenylmethyl)octyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L13 ANSWER 36 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2003:991296 CAPLUS

DOCUMENT NUMBER: 140:41822

TITLE: Preparation of acylamino(formyl)propanoic acids as caspase-1 inhibitors

INVENTOR(S): Alledia, Martin; Fahr, Bruce; Oslob, Johan; Raimundo, Brian C.; Romanowski, Michael J.

PATENT ASSIGNEE(S): Sunesis Pharmaceuticals, Inc., USA

SOURCE: PCT Int. Appl., 88 pp.

CODEN: PIXX02

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE

MO 2003103599 A2 20031218 MO 2003-US18021 20030605


MO 2003103599 A3 20040708

W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GR, GU, GM, HR, HU, ID, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MG, MK, MN, MW, MX, MY, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW

RM: GH, GM, KE, LS, MM, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, NG, SN, TD, TG

AU 2003238948 A1 20031222 AU 2003-238948 20030605


RN 612847-43-5 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl- (9CI)
(CA INDEX NAME)



RN 612848-56-3 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 612847-43-5
CMF C40 H43 N7 O2



CM 2


CRN 76-05-1
CMF C2 H F3 O2



RN 612848-57-4 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 612847-42-4
CMF C40 H43 N7 O2




CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 612848-58-5 CAPLUS
CN D-Arabinohexose, 2-deoxy-2-[[[2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl-6-quinoxaliny]carbonyl]amino]-, (2S)-, (2R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



PAGE 1-A

PAGE 1-B




RN 612848-59-6 CAPLUS
CN D-Arabinohexose, 2-deoxy-2-[[[2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl-6-quinoxaliny]carbonyl]amino]-, (2S)-, (2R)-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1


CRN 612848-58-5
CMF C40 H40 N6 O7

Absolute stereochemistry.



PAGE 1-A

PAGE 1-B




CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 612848-60-9 CAPLUS
CN D-Arabinohexose, 2-deoxy-2-[[[2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl-6-quinoxaliny]carbonyl]amino]-, (2S)-, (2R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



PAGE 1-A




RN 612848-61-0 CAPLUS
CN D-Arabinohexose, 2-deoxy-2-[[[2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl-6-quinoxaliny]carbonyl]amino]-, (2S)-, (2R)-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1


CRN 612848-60-9
CMF C40 H40 N6 O7

Absolute stereochemistry.



PAGE 1-A

PAGE 1-B



CM 2


CRN 76-05-1
CMF C2 H F3 O2



RN 616868-43-0 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-

yl)-1-piperidinyl)methyl]phenyl]-N-(3-(1H-imidazol-1-yl)propyl)-3-phenyl-
(9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B




RN 616868-44-1 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl)methyl]phenyl]-N-(3-(1H-imidazol-1-yl)propyl)-3-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 616868-43-0
CMF C40 H38 N8 O2

PAGE 1-A



PAGE 1-B




CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 616868-45-2 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl)methyl]phenyl]-3-phenyl-, ethyl ester (9CI) (CA INDEX NAME)



RN 616868-46-3 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl)methyl]phenyl]-3-phenyl-, ethyl ester, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 616868-45-2
CMF C36 H33 N5 O3




CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 616868-47-4 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl)methyl]phenyl]-2-phenyl-, ethyl ester (9CI) (CA INDEX NAME)



RN 616868-48-5 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl)methyl]phenyl]-2-phenyl-, ethyl ester, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 616868-47-4
CMF C36 H33 N5 O3




CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 616868-49-6 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl)methyl]phenyl]-N-(2-hydroxyethyl)-3-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)



RN 616868-50-9 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl)methyl]phenyl]-N-(2-hydroxyethyl)-3-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1

CRN 616868-49-6
CMF C36 H34 N6 O3




CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 616868-51-0 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl)methyl]phenyl]-N-(2-hydroxyethyl)-2-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)



RN 616868-52-1 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl)methyl]phenyl]-N-(2-hydroxyethyl)-2-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1


CRN 616868-51-0
CMF C36 H34 N6 O3



CM 2
CRN 76-05-1
CMF C2 H F3 O2




RN 616868-53-2 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(2-hydroxy-2-phenylethyl)-3-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)



RN 616868-54-3 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(2-hydroxy-2-phenylethyl)-3-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1
CRN 616868-53-2
CMF C42 H38 N6 O3




CM 2
CRN 76-05-1
CMF C2 H F3 O2




RN 616868-55-4 CAPLUS

CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(2-hydroxy-2-phenylethyl)-2-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)



RN 616868-56-5 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(2-hydroxy-2-phenylethyl)-2-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)


CM 1
CRN 616868-55-4
CMF C42 H38 N6 O3



CM 2
CRN 76-05-1
CMF C2 H F3 O2




RN 616868-67-8 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(5-hydroxypentyl)-3-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)



RN 616868-68-9 CAPLUS

CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(5-hydroxypentyl)-3-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)


CM 1
CRN 616868-67-8
CMF C39 H40 N6 O3



CM 2
CRN 76-05-1
CMF C2 H F3 O2




RN 616868-69-0 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(5-hydroxypentyl)-2-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)



RN 616868-70-3 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(3-(dimethylamino)-2,2-dimethylpropyl)-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)


CM 1
CRN 616868-69-0
CMF C39 H40 N6 O3



CM 2
CRN 76-05-1
CMF C2 H F3 O2



RN 616868-71-4 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-[3-(dimethylamino)-2,2-dimethylpropyl]-3-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)




PAGE 1-A

PAGE 1-B



RN 616868-72-5 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-[3-(dimethylamino)-2,2-dimethylpropyl]-3-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1
CRN 616868-71-4
CMF C41 H45 N7 O2




PAGE 1-B



CM 2
CRN 76-05-1
CMF C2 H F3 O2



RN 616868-73-6 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-[3-(dimethylamino)-2,2-dimethylpropyl]-2-phenyl-, (9CI) (CA INDEX NAME)




PAGE 1-B



RN 616868-74-7 CAPLUS

CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-[3-(dimethylamino)-2,2-dimethylpropyl]-2-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1
CRN 616868-73-6
CMF C41 H45 N7 O2




PAGE 1-B



CM 2
CRN 76-05-1
CMF C2 H F3 O2



RN 616868-75-8 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(3-hydroxy-2,2-dimethylpropyl)-3-phenyl-, (9CI) (CA INDEX NAME)




PAGE 1-B

PAGE 1-B



RN 616868-76-9 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(3-hydroxy-2,2-dimethylpropyl)-3-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1
CRN 616868-75-8
CMF C39 H40 N6 O3




PAGE 1-B



CM 2
CRN 76-05-1
CMF C2 H F3 O2



RN 616868-77-0 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(3-hydroxy-2,2-dimethylpropyl)-2-phenyl-, (9CI) (CA INDEX NAME)




PAGE 1-B



RN 616868-78-1 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(3-hydroxy-2,2-dimethylpropyl)-2-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1
CRN 616868-77-0
CMF C39 H40 N6 O3



PAGE 1-B




CM 2
CRN 76-05-1
CMF C2 H F3 O2



RN 616868-79-2 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[4-(aminosulfonyl)phenyl]methyl]-2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl- (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B




RN 616868-80-5 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[4-(aminosulfonyl)phenyl]methyl]-2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 616868-79-2
CMF C41 H37 N7 O4 S

PAGE 1-A



PAGE 1-B




CM 2

CRN 76-05-1
CMF C2 H F3 O2



RN 616869-39-7 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(1H-benzimidazol-2-yl)-1-piperidinyl]methyl]phenyl]-N-(2-hydroxyethyl)-7-methyl-2-phenyl- (9CI) (CA INDEX NAME)



RN 616869-40-0 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(1H-benzimidazol-2-yl)-1-piperidinyl]methyl]phenyl]-N-(2-hydroxyethyl)-7-methyl-2-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1

CRN 616869-39-7
CMF C37 H36 N6 O2

PAGE 1-B




CM 2

CRN 76-05-1
CMF C2 H F3 O2



RN 616868-81-6 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[4-(aminosulfonyl)phenyl]methyl]-3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



RN 616868-82-7 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[4-(aminosulfonyl)phenyl]methyl]-3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 616868-81-6
CMF C41 H37 N7 O4 S




CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 616869-41-1 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(1H-benzimidazol-2-yl)-1-piperidinyl]methyl]phenyl]-N-(2-hydroxyethyl)-7-methyl-3-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)



RN 616869-42-2 CAPLUS
CN 6-Quinoxalinecarboxamide, 2-[4-[[4-(1H-benzimidazol-2-yl)-1-piperidinyl]methyl]phenyl]-N-(2-hydroxyethyl)-7-methyl-3-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1

CRN 616869-41-1
CMF C37 H36 N6 O2



CM 2


CRN 76-05-1

CMF C2 H F3 O2




RN 616869-89-7 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-3-[4-[[4-(2-methyl-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl- (9CI) (CA INDEX NAME)


PAGE 1-A



PAGE 1-B




RN 616869-90-0 CAPLUS
CN 6-Quinoxalinecarboxamide, N-(2-hydroxyethyl)-3-[4-[[4-(2-methyl-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl- (9CI) (CA INDEX NAME)



RN 616869-92-2 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[3-(1H-imidazol-1-yl)propyl]-3-[4-[[4-(2-methyl-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 616869-95-5 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-3-[4-[[4-(5-fluoro-1H-benzimidazol-2-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl- (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B




RN 616870-60-1 CAPLUS
CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 612847-33-3
CMF C40 H38 N8 O2

PAGE 1-A



PAGE 1-B



CM 2

CRN 76-05-1
CMF C2 H F3 O2




RN 616870-81-6 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-3-[4-[[4-(2-methyl-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 616869-89-7
CMF C41 H45 N7 O

PAGE 1-A



PAGE 1-B



CM 2


CRN 76-05-1
CMF C2 H F3 O2



RN 616870-82-7 CAPLUS
CN 6-Quinoxalinecarboxamide, N-(2-hydroxyethyl)-3-[4-[[4-(2-methyl-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1

CRN 616869-90-0
CMF C37 H36 N6 O2



CM 2

CRN 76-05-1
CMF C2 H F3 O2





RN 616870-84-9 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[3-(1H-imidazol-1-yl)propyl]-3-[4-[[4-(2-methyl-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 616869-92-2
CMF C41 H40 N8 O

PAGE 1-A







CM 2

CRN 76-05-1
CMF C2 H F3 O2

RN 616870-87-2 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-3-[4-[[4-(5-fluoro-1H-benzimidazol-2-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 616869-95-5
CMF C40 H42 F N7 O

CM 2

CRN 76-05-1
CMF C2 H F3 O2

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 38 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2003:618232 CAPLUS

DOCUMENT NUMBER: 139:323527

TITLE: Preparation of triazolo[4,3-b]pyridazines and 2,3-diarylquinoxalines for the treatment of cancer

INVENTOR(S): Barnett, Stanley F.; Defeo-Jones, Deborah; Haskell, Kathleen M.; Huber, Hans E.; Nahas, Deborah D.;

Lindley, Craig W.; Zhao, Zhijian; Hartman, George D.

PATENT ASSIGNOR(S): Merck & Co., Inc., USA

SOURCE: PCT Int. Appl., 170 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent



LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003084473	A2	20031016	WO 2003-US10632	20030404
WO 2003084473	A3	20040212		
W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GR, GU, HM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MG, MK, MN, MM, MX, MY, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CP, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003226301	A1	20031020	AU 2003-226301	20030404
PRIORITY APPLN. INFO.:				
			US 2002-370827P	P 20020408
			US 2002-417202P	P 20021009
			WO 2003-US10632	W 20030404

GI



AB Triazolo[4,3-b]pyridazines I [R1 = (un)substituted Ph, furyl, thienyl, pyridinyl; R2 = substituted NH2, OH; R3 = H, R4 = (un)substituted cycloalkyl, aryl; R3R4 = (un)substituted CH:CH:CH] and quinoxalines II [R5, R6 = (un)substituted Ph; R7 = H, alkyl, halogen, OH, alkoxy] were prepared for use as inhibitors of one or two of the isoforms of Akt, a serine/threonine protein kinase, acting particularly on the pleckstrin

homol. domain of Akt. Thus, 3,6-dichloropyridazine was converted to its 4-cyclobutyl derivative which was cyclized with BzNHNH2 and aminated to give I [R1 = Ph, R2 = NHCH2CMe2CH2NMe2, R3 = H, R4 = cyclobutyl]. This compound had IC50 for inhibition of Akt1 of 1.4 μM.

612847-34-4P 612847-42-4P 612847-43-5P

612848-56-3P 612848-57-4P 612848-59-6P

612848-61-0P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USBS (Uses)

(Preparation of triazolo[4,3-b]pyridazines and 2,3-diarylquinoxalines for the treatment of cancer)

RN 612847-34-4 CAPLUS

CN 6-Quinoxalinecarboxamide, 3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-N-(3-(1H-imidazol-1-yl)propyl)-2-phenyl-, bis(trifluoroacetate) (9CI) (CA INDEX NAME)

CM 1


CRN 612847-33-3
CMF C40 H38 N8 O2

CM 2

CRN 76-05-1
CMF C2 H F3 O2

RN 612847-42-4 CAPLUS


CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)



RN 612847-43-5 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)


CM 1

CRN 612847-43-5
CMF C40 H43 N7 O2

RN 612848-56-3 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-2-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-3-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1

CRN 612847-43-5
CMF C40 H43 N7 O2


CM 2

CRN 76-05-1
CMF C2 H F3 O2

RN 612848-57-4 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[2-(diethylamino)ethyl]-3-[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]methyl]phenyl]-2-phenyl-, trifluoroacetate (9CI) (CA INDEX NAME)

CM 1
CRN 612847-42-4
CMF C40 H43 N7 O2




CM 2
CRN 76-05-1
CMF C2 H F3 O2



RN 612848-59-6 CAPLUS
CN D-arabino-Hexose, 2-deoxy-2-[[[2-[[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinylmethyl]phenyl]-3-phenyl-6-quinoxalinyloxy]carbonyl]amino]-2-oxo-1H-benzimidazol-1-yl]methyl]phenyl group, (2S)-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)


CM 1
CRN 612848-58-5
CMF C40 H40 N6 O7

Absolute stereochemistry.



PAGE 1-A

PAGE 1-A




CM 2
CRN 76-05-1
CMF C2 H F3 O2




RN 612848-61-0 CAPLUS
CN D-arabino-Hexose, 2-deoxy-2-[[[2-[[4-[[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinylmethyl]phenyl]-3-phenyl-6-quinoxalinyloxy]carbonyl]amino]-2-oxo-1H-benzimidazol-1-yl]methyl]phenyl group, (2S)-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1
CRN 612848-60-9
CMF C40 H40 N6 O7

Absolute stereochemistry.



PAGE 1-B



CM 2

CRN 76-05-1
CMF C2 H F3 O2



L13 ANSWER 39 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 2003:677647 CAPLUS

DOCUMENT NUMBER: 140:217939

TITLE: Highly enhanced duplex stability of dipyrro[3,2-a:2',3'-c]phenazine-modified oligonucleotide conjugate

AUTHOR(S): Kitamura, Yusuke; Ihara, Toshihiro; Shirasaka, Yoshinori; Mitsuura, Tomonori; Tazaki, Masato; Jyo, Akimori

CORPORATE SOURCE: Department of Applied Chemistry and Biochemistry, Kumamoto University, Kumamoto, 860-8555, Japan

SOURCE: Nucleic Acids Research Supplement (2003), 3 (3rd International Symposium on Nucleic Acids Chemistry [and] 30th Symposium on Nucleic Acids Chemistry in Japan, 2003), 95-96

PUBLISHER: CODEN: NARSCS

DOCUMENT TYPE: Oxford University Press

LANGUAGE: English

AB Dipyrro[3,2-a:2',3'-c]phenazine (DPPZ) or 1,10-phenanthroline (Phen) was tethered to the 5'-end of a short oligonucleotide (ODN) to generate two ODN conjugates. The conjugates formed stable duplexes with complementary 6 mer (d(TTAAGG)), which is one unit of telomeric repeats of human. The melting temperature of the duplex with DPPZ conjugate was higher than that of the corresponding duplex with unmodified 6 mer by 19.6 °C. This stabilization is enormous compared with those observed in other ODN conjugates reported previously. It would be attributed to the effective interaction of tethered heteroarom. groups with DNA base stack of the duplex.

IT 663942-79-8P

RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (preparation and characterization of highly enhanced duplex stability of dipyrrophenazine-modified oligodeoxyribonucleotide conjugates)

RN 663942-79-8 CAPLUS


CN Guanosine, thymidyl- (3' -5')-thymidyl- (3' -5')-2'-deoxyadenyl- (3' -5')-2'-deoxyguanylyl- (3' -5')-2'-deoxy-5'-O-[[[6-[[[dipyrro[3,2-a:2',3'-c]phenazine-11-yl]carbonyl]amino]hexyl]oxy]hydroxyphenyl]cytidyl- (3' -5')-2'-deoxycytidyl- (3' -5')-2'-deoxycytidyl- (3' -5')-thymidyl- (3' -5')-2'-deoxyadenyl- (3' -5')-2'-deoxyadenosine (1:1) (9CI) (CA INDEX NAME)

CM 1


CRN 663942-78-7
CMF C82 H96 N26 O37 P6


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B








CM 2

CRN 117490-04-7
CMF C60 H75 N24 O35 P5

Absolute stereochemistry.




REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 40 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2003:656742 CAPLUS
 DOCUMENT NUMBER: 139:197375
 TITLE: Preparation of piperidinyl alcohols as chemokine receptor modulators for treatment of diseases such as asthma
 INVENTOR(S): Alcaraz, Lilian; Furber, Mark; Purdie, Mark; Springthorpe, Brian
 PATENT ASSIGNEE(S): AstraZeneca A.B., Swed.
 SOURCE: PCT Int. Appl., 166 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003068743	A1	20030821	WO 2003-58258	20030317
W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RM: OH, OM, OS, LS, MS, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CP, CO, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TO				
CA 2472822	AA	20030821	CA 2003-2472822	20030317
AU 2003206554	A1	20030904	AU 2003-206554	20030317
BR 2003007477	A	20041109	BR 2003-7477	20030217
EP 1478624	A1	20041124	EP 2003-705600	20030217
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
US 2005107428	A1	20050519	US 2003-504936	20030317
CN 1633414	A	20050629	CN 2003-804130	20030317
JP 2005525341	T2	20050825	JP 2003-567874	20030217
ZA 2004006509	A	20050915	ZA 2004-6509	20040816
NO 2004003899	A	20041117	NO 2004-3899	20040917
PRIORITY APPLN. INFO.:			SE 2002-4465	A 20020318
			SE 2002-2673	A 20020909
			WO 2003-58258	N 20030217

CASREACT 139:197375; MARPAT 139:197375
 OTHER SOURCE(S):
 GI



AB The invention provides piperidinyl alcs. (shown as 1; variables defined below; e.g. N-[(2R)-3-[4-(3,4-dichlorophenoxy)piperidin-1-yl]-2-hydroxypropyl]-2-(methylsulfonyl)benzamide) for use as modulators of chemokine receptor (especially CCR3) activity for use in, for example, treating asthma. For 1: X is CH₂, O, S(O)₂ or NR₁₀; Y is a bond, CH₂, NR₁₀, CH₂NH, CH₂NH(CO), CH(OH), CH(NHCO₂R₁₁), CH(NHCO₂R₁₁), CH₂O or CH₂S; Z is C(O), or when Y is a bond Z can also be S(O)₂; R₁ is (un)substituted aryl, (un)substituted heterocyclyl or C₄-6 cycloalkyl fused to a benzene ring; addnl. details are given in the claims. Percent inhibition at 3 nM eotaxin of eotaxin-mediated human eosinophil chemotaxis is tabulated for 16 examples of 1, e.g. 106 % for N-[(2R)-3-[4-(3,4-dichlorophenoxy)piperidin-1-yl]-2-hydroxypropyl]-1-oxo-1,2-dihydroisoquinoline-4-carboxamide. Histamine H₁ receptor binding activity was determined for the same compds., e.g. pK_i = 8.4 for N-[(2R)-3-[4-(3,4-dichlorophenoxy)piperidin-1-yl]-2-hydroxypropyl]-1-oxo-1,2-dihydroisoquinoline-4-carboxamide. 49 Example preps. of intermediates and 234 of 1 are included. For example, to prepare N-[(2R)-3-[4-(3,4-dichlorophenoxy)piperidin-1-yl]-2-hydroxypropyl]-2-(methylsulfonyl)benzamide (0.055 g), a mixture of 2-(methylsulfonyl)benzoic

acid (0.063 g), (2R)-1-amino-3-[4-(3,4-dichlorophenoxy)piperidin-1-yl]propan-2-ol (0.1 g) and N,N-diisopropylethylamine (0.1 mL) in dry DMP (3 mL) was cooled to 0° with stirring; 2-(1H-9-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (0.13 g) was added and the mixture was stirred at 0° for 1-2 h. The invention also provides a process for making 4-(3,4-dichlorophenoxy)piperidine, which is useful as an intermediate for making certain compds. of the invention. The process comprises (a) reacting 4-hydroxypiperidine with a suitable base in a suitable solvent at room temperature; and (b) heating the mixture so produced

and 1,2-dichloro-4-fluorobenzene at 50-90°, or at reflux of the solvent used.


IT 503882-05-7p, N-[(2R)-3-[4-(3,4-dichlorophenoxy)piperidin-1-yl]-2-hydroxypropyl]quinoxaline-6-carboxamide
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; preparation of piperidinyl alcs. as chemokine receptor modulators for treatment of diseases such as asthma)

RN 503882-05-7 CAPLUS

CN 6-Quinoxalinecarboxamide, N-[(2R)-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxypropyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 41 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2003:376819 CAPLUS
 DOCUMENT NUMBER: 138:385173
 TITLE: Preparation of N,N'-substituted-1,3-diamino-2-hydroxypropanes for treating Alzheimer's disease
 INVENTOR(S): Varghese, John; Maillard, Michel; Jagodzinska, Barbara; Beck, James P.; Gallunas, Andrea; Feng, Larry; Seely, Jennifer; Tenbrink, Ruth; Freshos, John; Mickelson, John; Senala, Lakshman; Hom, Roy
 PATENT ASSIGNEE(S): Sanofi Pharmaceuticals, Inc., USA; Pharmacia & Upjohn Company
 SOURCE: PCT Int. Appl., 1243 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003040096	A2	20030515	WO 2002-US36072	20021108
WO 2003040096	A3	20040506		
W: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				

RW: GH, GM, KS, LS, MW, MZ, SD, SL, SZ, TZ, UO, ZM, ZW, AM, AZ, BY, KQ, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GO, GM, ML, MR, NE, SN, TD, TG

CA 2466284 A1 20030515 CA 2002-2466284 20021108
WO 2003040096 A2 20030515 WO 2002-2436072 20021108

M: AR, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW

RW: GH, GM, KS, LS, MW, MZ, SD, SL, SZ, TZ, UO, ZM, ZW, AM, AZ, BY, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GO, GM, ML, MR, NE, SN, TD, TG


US 2004171881 A1 20040902 US 2002-291318 20021108
EP 1453789 A2 20040908 EP 2002-793909 20021108

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IS, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK

BR 2002014035 A 20050426 BR 2002-14035 20021108
JP 2005520791 T2 20050714 JP 2003-542142 20021108
CN 1759095 A 20060412 CN 2002-826786 20021108
ZA 2004003578 A 20051010 ZA 2004-3578 20040511
NO 2004002359 A 20040806 NO 2004-2359 20040607

PRIORITY APPLN. INFO.: US 2001-337122P P 20011108
US 2001-344086P P 20011228
US 2002-345635P P 20020103
WO 2002-US36072 W 20021108

OTHER SOURCE(S): MARPAT 138:385173
GI



AB The title compds. [I; R1 = (un)substituted alkyl, alkenyl, alkynyl, etc.; R2 = H, alkyl, haloalkyl, alkenyl, etc.; R3 = H, alkyl, haloalkyl, alkenyl, etc.; or R2 and R3 are taken together with the carbon to which they are attached to form a carbocycle of 3-7 carbon atoms, optionally where one carbon atom is replaced by a heteroatom selected from the group consisting of O, S, SO2, (un)substituted NH; R4 = alkyl, haloalkyl,


hydroxyalkyl, etc.; R5 = R6X (wherein X = CO, SO2, (un)substituted CH2; R6 = (un)substituted Ph, naphthyl, indanyl, etc.); R25 = H, alkyl, alkoxy, etc.] which have activity as inhibitors of β -secretase and are therefore useful in treating a variety of disorders such as Alzheimer's disease, were prepared. E.g., a multi-step synthesis of (1S,2R)-II, starting from (2S)-2-[(tert-butoxycarbonyl)amino]-3-(3,5-difluorophenyl)propanoic acid, was given. The compds. I showed IC50 of < 20 μ M in cell free inhibition assay utilizing a synthetic APP substrate. This is a Part 1 of 1-2 series.

IT 527723-62-89
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(Preparation of N,N'-substituted-1,3-diamino-2-hydroxypropanes for treating Alzheimer's disease)

RN 527725-62-8 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[(1S,2R)-1-[(3,5-difluorophenyl)methyl]-3-[[[3-ethylphenyl)methyl]amino]-2-hydroxypropyl]-1,2,3,4-tetrahydro-2,3-dioxo-(5C)] (CA INDEX NAME)

Absolute stereochemistry.



L13 ANSWER 42 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2003:282533 CAPLUS
DOCUMENT NUMBER: 138:304304
TITLE: Preparation of difluoroalkene derivatives as pest control agents containing the same, and intermediate thereof

INVENTOR(S): Abe, Tetsumiya; Tamai, Ryuji; Ito, Minoru; Tamaru, Masatoshi; Yano, Hiroyuki; Takahashi, Satoru; Muramatsu, Norimichi
PATENT ASSIGNER(S): Kumiai Chemical Industry Co., Ltd., Japan; Ihara Chemical Industry Co., Ltd.

SOURCE: PCT Int. Appl., 195 pp.
CODEN: PIXX22
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003029211	A1	20030410	WO 2002-JP10142	20020930
M: AR, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,				

UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: GH, GM, KS, LS, MW, MZ, SD, SL, SZ, TZ, UO, ZM, ZW, AM, AZ, BY, KQ, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GO, GM, ML, MR, NE, SN, TD, TG

EP 1439169 A1 20040721 EP 2002-775265 20020930

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IS, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK

US 2004248872 A1 20041209 US 2004-491128 20040329
JP 2001-295687 A 20010928
JP 2001-142329 A 20020517
WO 2002-JP10142 W 20020930

PRIORITY APPLN. INFO.: US 2002-345635P P 20020103
WO 2002-US36072 W 20021108


OTHER SOURCE(S): MARPAT 138:304304

AB The difluoroalkenyl heterocyclecarboxylate, -thiocarboxylate, or dithiocarboxylate derivative, represented by the general formula Q-C(L1)-L2-(CH2)n-C(F)2; CF2 or pharmacol. acceptable salts thereof (wherein L1 and L2 are the same or different and each represents oxygen or sulfur; n is an integer of 2 to 8; and Q represents an optionally substituted 5- to 12-membered heterocyclic group having any desired heteroatom selected among nitrogen, oxygen, and sulfur wherein the heteroatom in the heterocyclic ring is a nitrogen, it may be oxidized to N-oxide), which are useful as insecticides, acaricides, and nematocides, are prepared. These compds. are sufficiently effective in controlling various pests even when used in a small dose and are highly safe for crops, natural enemies to the pests, and animals. Thus, 4-phenyl-1,2,3-thiadiazole-5-carboxylic acid 0.23, 6,6-difluoro-5-methyl-5-hexenol 0.17, and 4-dimethylaminopyridine 0.13 g were dissolved in 4 mL CH2Cl2, treated with 0.29 g 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride at room temperature, and stirred for 20 h to give 6,6-difluoro-5-methyl-5-hexenyl 4-phenyl-1,2,3-thiadiazole-5-carboxylate (I). I and 4,4-difluoro-3-methyl-3-butenyl 6-butoxy-2-methylpyrimidine-4-carboxylate at 500 ppm controlled 290% 4th instar larvae of Nilaparvata lugens.

IT 509100-31-6P
RL: AGR (Agricultural use); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)

(Preparation of difluoroalkenyl heterocyclecarboxylate and -thiocarboxylates as pest control agents such as insecticides, acaricides, and nematocides)

RN 509100-31-6 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 4,4-difluoro-3-methyl-3-butenyl ester (9C1) (CA INDEX NAME)



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 43 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2002:889219 CAPLUS
DOCUMENT NUMBER: 137:379407
TITLE: Colorimetric sensor compositions and methods
INVENTOR(S): Sessler, Jonathan; Andrioletti, Bruno; Try, Andrew
PATENT ASSIGNER(S): USA
SOURCE: U.S., 30 pp.
CODEN: USXXAM

DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6482949	B1	20021119	US 2000-579040	20000526
US 2003162960	A1	20030828	US 2002-222028	20020816
PRIORITY APPLN. INFO.:			US 1999-136467P	P 19990528
			US 2000-579040	A3 20000526



OTHER SOURCE(S): MARPAT 137:379407

AB The present invention provides novel compounds, exemplified by pyrrolic nitrogens used as anion and neutral species recognition elements with an aromatic core as a signal group. Described are methods for the synthesis of various pyrrolic aryl compds. as well as various applications for these compds. Methods of use include the binding and detection of specific analytes in a mixture and, in some examples, the separation of the analyte from the mixture. Addnl. methods of use include the transport of therapeutic agents and the sensing of components, degradants, and impurities in foodstuffs.

IT 475476-81-4P
RL: SPN (Synthetic preparation); PREP (Preparation)

(colorimetric sensor compds. and methods based on pyrrolic-aryl compds. for anion and neutral species recognition and determination)

RN 475476-81-4 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2,3-di-1H-pyrrol-2-yl-, octyl ester (9C1) (CA INDEX NAME)



REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 44 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2002:808500 CAPLUS
DOCUMENT NUMBER: 138:34814
TITLE: Synthesis of the DNA-[Ru(tpy)(dppz)(CH3CN)]2+ conjugates and their photo cross-linking studies with the complementary DNA strand

AUTHOR(S): Ossipov, Dimitri; Oghil, Suresh; Chattopadhyaya, Jyoti
CORPORATE SOURCE: Biomedical Center, Department of Bioorganic Chemistry, University of Uppsala, Uppsala, S-751 23, Swed.
SOURCE: Journal of the American Chemical Society (2002), 124(45), 13416-13433

PUBLISHER: CODEN: JACBAT; ISSN: 0002-7863
DOCUMENT TYPE: American Chemical Society
LANGUAGE: English
OTHER SOURCE(S): CASREACT 138:34814


AB We here report our studies on the conjugation of photoreactive Ru²⁺-complex to oligonucleotides (ODNs), which give a stable duplex with the complementary target DNA strand. These functionalized DNA duplexes bearing photoreactive Ru²⁺-complex can be specifically photolyzed to give the reactive aqua derivative, [Ru(tpy)(dppz)(H₂O)]²⁺-ODN (tpy = 2,2':6',2''-terpyridine; dppz = dipyrro[3,2-a:2',3'-c]phenazine), in situ, which successfully cross-links to give photoproduct(s) in the duplex form with the target complementary DNA strand. Thus, the stable precursor of the aquaruthenium complex, the monofunctional polypyridyl ruthenium complex [Ru(tpy)(dppz)(CH₃CN)]²⁺, has been site-specifically tethered to ODN, for the first time, by both solid-phase synthesis and postsynthetic modifications. (i) In the first approach, pure 3'-[Ru(tpy)(dppz)(CH₃CN)]²⁺-ODN conjugate has been obtained in 42% overall yield (from the monomer blocks) by the automated solid-phase synthesis on a support labeled with [Ru(tpy)(dppz)Cl]⁺ complex with subsequent liberation of the crude conjugate from the support under mild conditions and displacement of the Cl⁻ ligand by acetonitrile in the coordination sphere of the Ru²⁺ label. (ii) In the second approach, the single-modified (3'- or 5'- or middle-modified) or 3',5'-bis-modified Ru²⁺-ODN conjugates were prepared in 28-50% yield by an amide bond formation between an active ester of the metal complex and the ODNs conjugated with an amino linker. The pure conjugates were characterized unambiguously by UV-visible (UV-vis) absorption spectroscopy, enzymic digestion followed by HPLC quantitation, PAGE (PAGE), and mass spectrometry (MALDI-TOF as well as by ESI). [Ru(tpy)(dppz)(CH₃CN)]²⁺-ODNs form highly stabilized ODN-DNA duplexes compared to the unlabeled counterpart (ΔT_m varies from 8.4 to 23.6°) as a result of intercalation of the dppz moiety; they undergo clean and selective photolysis of the CH₃CN ligand to give the corresponding aqua complex, [Ru(tpy)(dppz)(H₂O)]²⁺-ODNs (in the aqueous medium), which is evidenced from the change of their UV-vis absorption properties and the detection of the naked Ru²⁺-ODN ions generated in the course of the matrix-assisted laser desorption ionization time-of-flight (MALDI-TOF) mass spectrometric anal. Thus, when [Ru(tpy)(dppz)(CH₃CN)]²⁺-ODN conjugate was hybridized to the complementary guanine (G)-rich target strand (T), and photolyzed in a buffer (pH 6.8), the corresponding aqua complex formed in situ immediately reacted with the G residue of the opposite strand, giving the cross-linked product. The highest yield (34%) of the photo cross-linked product obtained was with the ODN carrying two reactive Ru²⁺ centers at both 3'- and 5'-ends. For ODNs carrying only one Ru²⁺ complex, the yield of the cross-linked adduct in the corresponding duplex is found to decrease in the following order: 3'-Ru²⁺-ODN (22%) > 5'-Ru²⁺-ODN (9%) > middle-Ru²⁺-ODN (7%). It was also found that the photo cross-coupling efficiency of the tethered Ru²⁺ complex with the target T strand decreased as the stabilization of the resulting duplex increased: 3'-Ru²⁺-ODN (VI-T) (ΔT_m = 7°) < 5'-Ru²⁺-ODN (V-T) (ΔT_m = 16°) < middle-Ru²⁺-ODN (VII-T) (ΔT_m = 24.3°, Table 2). This shows that, with the rigidly packed structure, as in the duplex with middle-Ru²⁺-ODN, the metal center flexibility is considerably reduced, and consequently the accessibility of target G residue by the aquaruthenium moiety becomes severely restricted, which results in a poor yield in the cross-coupling reaction. The cross-linked product was characterized by PAGE, followed by MALDI-TOF MS.

IT 478818-94-9P 478819-02-2P 478819-70-4P
478819-77-1P 478819-80-6P
RL: BSU (Biological study, unclassified); SPN (Synthetic preparation);
BIOL (Biological study); PREP (Preparation)
(DNA-[Ru(tpy)(dppz)(CH₃CN)]²⁺ conjugates and their photo crosslinking studies with complementary DNA strand shows enhanced thermal and nuclease stability)

RN 478818-94-9 CAPLUS
CN Ruthenium(2+), (acetonitrile)(N-ethylpyrido[3,2-a:2',3'-c]phenazine-11-carboxamide-κN4,κN5)(2,2':6',2''-terpyridine-κN1,κN1',κN1'')-, (OC-6-43)-, bis[hexafluorophosphate(1-)] (9CI) (CA INDEX NAME)


CM 1

CRN 478818-93-8
CMF C38 H29 N9 O Ru
CCI CCS



CM 2


CRN 16919-18-9
CMF F6 P
CCI CCS



RN 478819-02-2 CAPLUS
CN Ruthenium(2+), aqua(N-ethylpyrido[3,2-a:2',3'-c]phenazine-11-carboxamide-κN4,κN5)(2,2':6',2''-terpyridine-κN1,κN1',κN1'')-, (OC-6-43)-, bis[hexafluorophosphate(1-)] (9CI) (CA INDEX NAME)


CM 1

CRN 478819-01-1
CMF C36 H28 N8 O2 Ru
CCI CCS



CM 2


CRN 16919-18-9
CMF F6 P
CCI CCS




RN 478819-70-4 CAPLUS
CN Ruthenium(2+), [N-[2-[2-[2-(2,3-dihydroxypropoxy)ethoxy]ethoxy]ethyl]dipyrro[3,2-a:2',3'-c]phenazine-11-carboxamide-κN4,κN5](pyridine)(2,2':6',2''-terpyridine-κN1,κN1',κN1'')-, (OC-6-32)-, bis[hexafluorophosphate(1-)] (9CI) (CA INDEX NAME)

CM 1

CRN 478819-69-1
CMF C48 H45 N9 O6 Ru
CCI CCS




PAGE 1-A



CM 2


CRN 16919-18-9
CMF F6 P
CCI CCS




PAGE 1-B

RN 478819-77-1 CAPLUS
 CN Ruthenium(2+), (acetonitrile) [N-[2-[2-(2,3-dihydroxypropoxy)ethoxy]ethoxy]ethyl]dipyrido[3,2-a:2',3'-c]phenazine-11-carboxamide-κN4,κN5[(2,2':6',2'':terpyridine-κN1,κN1',κN1'')], (OC-6-43)-, bis[hexafluorophosphate(1-)] (9CI) (CA INDEX NAME)
 CM 1
 CRN 478819-76-0
 CMF C45 H43 N9 O6 Ru
 CCI CCS


PAGE 1-A



PAGE 1-B




CM 2
 CRN 16919-18-9
 CMF F6 P
 CCI CCS




RN 478819-80-6 CAPLUS
 CN Ruthenium(2+), aqua [N-[2-[2-(2,3-dihydroxypropoxy)ethoxy]ethoxy]ethyl]dipyrido[3,2-a:2',3'-c]phenazine-11-carboxamide-κN4,κN5[(2,2':6',2'':terpyridine-κN1,κN1',κN1'')], (OC-6-43)-, bis[hexafluorophosphate(1-)] (9CI) (CA INDEX NAME)

1''), (OC-6-43)-, bis[hexafluorophosphate(1-)] (9CI) (CA INDEX NAME)
 CM 1
 CRN 478819-79-3
 CMF C43 H42 N8 O7 Ru
 CCI CCS


PAGE 1-A



PAGE 1-B




CM 2
 CRN 16919-18-9
 CMF F6 P
 CCI CCS




IT 478415-63-1P 478415-64-4P 478819-41-9P
 478819-51-1P 478819-57-7P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (DNA-[Ru(tpy)(dppz)(CH3CN)]₂⁺ conjugates and their photo crosslinking studies with complementary DNA strand shows enhanced thermal and nuclease stability)
 RN 478415-63-3 CAPLUS
 CN Dipyrido[3,2-a:2',3'-c]phenazine-11-carboxamide, N-[2-[2-(2,3-dihydroxypropoxy)ethoxy]ethoxy]ethyl]- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 478415-64-4 CAPLUS
 CN Dipyrido[3,2-a:2',3'-c]phenazine-11-carboxamide, N-[11-[bis(4-methoxyphenyl)phenylmethoxy]-14-bis(4-methoxyphenyl)-14-phenyl-3,6,9,13-tetraoxatetradec-1-yl]- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 478819-41-9 CAPLUS
 CN Ruthenium(1+), chloro [N-[2-[2-(2,3-dihydroxypropoxy)ethoxy]ethoxy]ethyl]dipyrido[3,2-a:2',3'-c]phenazine-11-carboxamide-κN4,κN5[(2,2':6',2'':terpyridine-κN1,κN1',κN1'')], (OC-6-43)-, hexafluorophosphate(1-) (9CI) (CA INDEX NAME)

CM 1
 CRN 478819-40-8
 CMF C43 H40 Cl N8 O6 Ru
 CCI CCS

PAGE 1-A




PAGE 1-B



CM 2

CRN 16919-18-9
CMF P6 P
CCI CCS




RN 478819-51-1 CAPLUS
CN Ruthenium(1+), chloro[N-[11-hydroxy-14,14-bis(4-methoxyphenyl)-14-phenyl-3,6,9,13-tetraoxatetradec-1-yl]dipyrido[3,2-a:2',3'-c]phenazine-11-carboxamide-κN4,κN5][2,2':6',2''-terpyridine-κN1,κN1'',κN1'''], (OC-6-43)-, hexafluorophosphate(1-)
(9CI) (CA INDEX NAME)


CM 1

CRN 478819-50-0
CMF C64 H58 Cl N8 O8 Ru
CCI CCS

PAGE 1-A




PAGE 1-B




CM 2

CRN 16919-18-9
CMF P6 P
CCI CCS




RN 478819-57-7 CAPLUS
CN Ruthenium, chloro[mono[2-bis(4-methoxyphenyl)phenylmethoxy]-1-[12-(dipyrido[1,2-a:2',3'-c]phenazin-11-yl)-κN4,κN5]-12-oxo-2,5,8-trioxo-11-azadodec-1-yl]ethyl] butanedioate[2,2':6',2''-terpyridine-κN1,κN1'',κN1'''], (OC-6-43)- (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B




REFERENCE COUNT: 71 THERE ARE 71 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 45 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2002:558411 CAPLUS


DOCUMENT NUMBER: 137:257239
TITLE: SAR by MS: A Ligand Based Technique for Drug Lead Discovery Against Structured RNA Targets
AUTHOR(S): Swayze, Eric B.; Jefferson, Elizabeth A.; Sannes-Lowery, Kristin A.; Blyn, Lawrence B.; Risen, Lisa M.; Arakawa, Satoshi; Osgood, Stephen A.; Hofstadler, Steven A.; Griffey, Richard H.
CORPORATE SOURCE: Isis Therapeutics, A Division of Isis Pharmaceuticals Inc., Carlsbad, CA, 92008, USA
SOURCE: Journal of Medicinal Chemistry (2002), 45(18), 3816-3819
CODEN: JMCNAR; ISSN: 0022-2623
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 137:257239

AB A technique for lead discovery vs. RNA targets utilizing mass spectrometry (MS) screening methods is described. The structure-activity relationships (SAR) derived from assaying weak binding motifs allows the pharmacophores discovered to be elaborated via "SAR by MS" to higher affinity ligands. Application of this strategy to a subdomain of the 23S rRNA afforded a new class of compds. with functional activity.
IT 462119-54-6 462119-55-7 462119-56-8
RL: CST (Combinatorial study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); CMBI (Combinatorial study); USES (Uses)
(SAR by MS: ligand-based technique for drug lead discovery against structured RNA targets)


RN 462119-54-6 CAPLUS
CN 6-Quinoxalinecarboxamide, 1,2,3,4-tetrahydro-2,3-dioxo-N-propyl- (9CI) (CA INDEX NAME)



RN 462119-55-7 CAPLUS
CN 6-Quinoxalinecarboxamide, 1,2,3,4-tetrahydro-2,3-dioxo-N-2-propenyl- (9CI) (CA INDEX NAME)




RN 462119-56-8 CAPLUS
CN 6-Quinoxalinecarboxamide, 1,2,3,4-tetrahydro-2,3-dioxo-N-(phenylmethyl)- (9CI) (CA INDEX NAME)



IT 462119-69-3P
RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
(SAR by MS: ligand-based technique for drug lead discovery against structured RNA targets)

RN 462119-69-3 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[3-[5-[[[1(R)-3-amino-1-(1-piperazinylcarbonyl)propyl]amino]carbonyl]-2-furanyl]propyl]-1,2,3,4-tetrahydro-2,3-dioxo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 46 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2002:501576 CAPLUS


DOCUMENT NUMBER: 137:208794
TITLE: Electron-deficient columnar plastic crystals
AUTHOR(S): Bock, Harald; Babeau, Annick; Seguy, Isabelle; Jolinat, Pascale; Destruel, Pierre
CORPORATE SOURCE: Centre de Recherche Paul Pascal, CNRS, Pessac, 33600, Fr.
SOURCE: ChemPhysChem (2002), 3(6), 532-535
CODEN: CPCHPT; ISSN: 1439-4235
PUBLISHER: Wiley-VCH Verlag GmbH
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The compound 5,6,11,12,17,18-hexazetrinsaphthylene-2,8,15-tricarboxylic acid (I) was synthesized from hexaketocyclohexane octahydrate and 3,4-diaminobenzoic acid which were refluxed in glacial acetic acid. The synthesis of the corresponding esters of I (R = Et, Pr, Bu, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, 1-ethylpropyl, 2-methylpropyl, 2-ethylbutyl, 2-ethoxyethyl, 2-isopropoxyethyl, rac-3-methoxybutyl, and di(ethoxymethyl)methyl) is also described. Furthermore, the compound 3',5',3'',5'',3''',5'''-hexapentylloxycarbonyl-2,4,6-triphenyl-1,3,5-triazine was prepared by transesterification of the hexamethyl homolog which was refluxed with K2CO3, 1-pentanol, and 1-bromopentane. The compds. were characterized by XRD, MS, IR-NMR spectroscopy, DSC, optical-absorption spectroscopy, and cyclic voltammetry. They form columnar mesophases with a widely differing degree of order going from highly fluidic nematic via hexagonal liquid crystallinity to plastic crystallinity. The electron


transport in the strongly electron-deficient compound I is discussed.
 IT 444579-19-5P
 RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)
 (Preparation and electronic and optical properties of)
 RN 444579-19-5 CAPLUS
 CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tributyl ester (9CI) (CA INDEX NAME)




IT 444579-17-3P 444579-18-4P 444579-20-8P
 444579-21-9P 444579-22-0P 444579-23-1P
 444579-24-2P 444579-25-3P 444579-26-4P
 444579-27-5P 444579-28-6P 444579-29-7P
 444579-30-0P 444579-31-1P
 RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)
 (Preparation and liquid-crystalline transition temps. of)
 RN 444579-17-3 CAPLUS
 CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, triethyl ester (9CI) (CA INDEX NAME)




RN 444579-18-4 CAPLUS
 CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tripropyl ester (9CI) (CA INDEX NAME)




RN 444579-20-8 CAPLUS
 CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tripropyl ester (9CI) (CA INDEX NAME)




RN 444579-21-9 CAPLUS
 CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, trihexyl ester (9CI) (CA INDEX NAME)




RN 444579-22-0 CAPLUS
 CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, triheptyl ester (9CI) (CA INDEX NAME)




RN 444579-23-1 CAPLUS
 CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, trioctyl ester (9CI) (CA INDEX NAME)




RN 444579-24-2 CAPLUS
 CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, trinonyl ester (9CI) (CA INDEX NAME)




RN 444579-25-3 CAPLUS
 CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris(1-ethylpropyl) ester (9CI) (CA INDEX NAME)




RN 444579-26-4 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris(1-methylpropyl) ester (9CI) (CA INDEX NAME)




RN 444579-27-5 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris(2-ethylbutyl) ester (9CI) (CA INDEX NAME)




RN 444579-28-6 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris(2-ethoxyethyl) ester (9CI) (CA INDEX NAME)




RN 444579-29-7 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris[2-(1-methylethoxy)ethyl] ester (9CI) (CA INDEX NAME)



RN 444579-30-0 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris(3-methoxybutyl) ester (9CI) (CA INDEX NAME)



RN 444579-31-1 CAPLUS
CN Diquinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, tris[2-ethoxy-1-(ethoxymethyl)ethyl] ester (9CI) (CA INDEX NAME)




REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 47 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2002:481308 CAPLUS
DOCUMENT NUMBER: 137:194994
TITLE: New Inhibitors of Bacterial Protein Synthesis from a Combinatorial Library of Macrocycles
AUTHOR(S): Jefferson, Elizabeth A.; Arakawa, Satoshi; Blyn, Lawrence B.; Miyaji, Alycia; Osgood, Stephen A.; Ranken, Raymond; Risen, Lisa M.; Swayze, Eric E.
CORPORATE SOURCE: Ibis Therapeutics, A Division of Isis Pharmaceuticals Inc., Carlsbad, CA, 92008, USA
SOURCE: Journal of Medicinal Chemistry (2002), 45(16), 3430-3439
CODEN: JMCMAR; ISSN: 0022-2623
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
AB A mixture-based combinatorial library of 14-membered macrocycles was synthesized to target rRNA and uncover a new class of antibacterial agents. High-throughput screening identified a macrocyclic mixture that inhibited cell-free-coupled transcription/translation in *Escherichia coli*-derived extracts, with an IC50 value in the 25-50 µM range. In a follow-up library of 64 single macrocycles, 9 gave IC50 values ranging from 12 to 50 µM in the cell-free protein synthesis inhibition assay. Some of the macrocycles were screened in a translation inhibition assay, and IC50 values generally paralleled those obtained in the uncoupled transcription/translation assay. Additional analogs were prepared in a preliminary structure-activity relation study, and more potent macrocycles were identified with low micromolar activity (IC50 values = 2-3 µM). Some of these macrocycles displayed antibacterial activity against lipopolysaccharide mutant *E. coli* bacterial cells (IC50 values = 12-50 µM).
IT 452338-59-9 452338-60-2 452338-61-3
452338-62-4 452338-63-5
EL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USKS (Uses)
(new inhibitors of bacterial protein synthesis from a combinatorial library of macrocycles)
RN 452338-59-9 CAPLUS
CN 6-Quinoxalino[2,3-a:2',3'-c]phenazine-2,8,15-tricarboxylic acid, N-[2-[[[3,4,5,6,7,8,9,10,11,12-decahydro-5,8,12-trioxo-2H-1,4,7,11-


benzoxatriazacyclotetradecin-14-yl]amino]-2-oxoethyl]-1,2,3,4-tetrahydro-2,3-dioxo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A



PAGE 1-B



PAGE 1-C




RN 452338-60-2 CAPLUS


CN 6-Quinoxalinecarboxamide, N-[(3S,6S,9S)-9-amino-3-[(aminiminomethyl)amino]propyl]-6-[(4-aminophenyl)methyl]-3,4,5,6,7,8,9,10,11,12-decahydro-5,8,12-trioxo-2H-1,4,7,11-benzoxatriazacyclotetradecin-14-yl]amino]-2-oxoethyl]-1,2,3,4-tetrahydro-2,3-dioxo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A



PAGE 1-B



PAGE 1-C




RN 452338-61-3 CAPLUS


CN 6-Quinoxalinecarboxamide, N-[(3S,6S,9S)-9-amino-3-[(aminiminomethyl)amino]propyl]-6-[(4-aminophenyl)methyl]-3,4,5,6,7,8,9,10,11,12-decahydro-5,8,12-trioxo-2H-1,4,7,11-benzoxatriazacyclotetradecin-14-yl]amino]-2-oxoethyl]-1,2,3,4-tetrahydro-2,3-dioxo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A



PAGE 1-B



PAGE 1-C




RN 452338-62-4 CAPLUS


CN 6-Quinoxalinecarboxamide, N-[(3S,6S,9S)-9-amino-3-[(aminiminomethyl)amino]propyl]-6-[(4-aminophenyl)methyl]-3,4,5,6,7,8,9,10,11,12-decahydro-5,8,12-trioxo-2H-1,4,7,11-benzoxatriazacyclotetradecin-14-yl]amino]-4-oxobutyl]-1,2,3,4-tetrahydro-2,3-dioxo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A



PAGE 1-B



PAGE 1-C




RN 452338-63-5 CAPLUS


CN 6-Quinoxalinecarboxamide, N-[(3S,6S,9S)-9-amino-3-[(aminiminomethyl)amino]propyl]-6-[(4-aminophenyl)methyl]-3,4,5,6,7,8,9,10,11,12-decahydro-5,8,12-trioxo-2H-1,4,7,11-benzoxatriazacyclotetradecin-14-yl]amino]-6-oxohexyl]-1,2,3,4-tetrahydro-2,3-dioxo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A



PAGE 1-B



PAGE 1-C



REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 48 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2002:184970 CAPLUS
 DOCUMENT NUMBER: 136:221460
 TITLE: Improvements relating to water treatment
 INVENTOR(S): Walker, Gavin Michael
 PATENT ASSIGNEE(S): The Queen's University of Belfast, UK
 SOURCE: PCT Int. Appl., 17 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English


FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002020152	A1	20020314	WO 2001-GB3994	20010906
W: AR, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GR, GM, HR, HU, ID, IL, IN, IS, JP, KS, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2001096052	A5	20020322	AU 2001-86052	20010906
EP 1322415	A1	20030702	EP 2001-965407	20010906
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
US 2004020859	A1	20040205	US 2003-363563	20030602
PRIORITY APPLN. INFO.: GB 2000-22049 A 20000908 WO 2001-GB3994 W 20010906				

AB A process for obtaining an adsorbent/flocculant material comprises heating dolomite to around 800°. The heated dolomite is washed with a suitable material able to increase its surface porosity, such as by removing any magnesium oxide from the surface pores. One particular washing substance is borax buffer. Use of the adsorbent/flocculant material of the invention to adsorb one or more substances such as nitrates and phosphates from a material such as water, is also described.

IT 206058-73-3, Levafix Brilliant Blue S-B
 RL: REM (Removal or disposal); PROC (Process)
 (water treatment using treated dolomite as adsorbent/flocculant)
 RN 206058-73-3 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[[4-[[[(2,3-dichloro-6-quinoxalyl)carbonyl]amino]methyl]-3-sulfonyl]methyl]amino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A





●2 Na

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 49 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2002:157741 CAPLUS
 DOCUMENT NUMBER: 136:200190
 TITLE: Benzimidazoles and analogues and their use as neutrophil inhibitors
 INVENTOR(S): Bush, Rodney Dean; Hershberger, Paul Mitchell; Young, Judith Anne; Kasibhatla, Shavani
 PATENT ASSIGNEE(S): The Procter & Gamble Company, USA
 SOURCE: PCT Int. Appl., 56 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002016327	A1	20020328	WO 2001-US25224	20010810
WO 2002016327	C1	20020801		
W: AR, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GR, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2001081246	A5	20020304	AU 2001-81246	20010810
US 2004006104	A1	20040108	US 2003-368261	20030318
PRIORITY APPLN. INFO.: US 2000-227201P P 20000823 WO 2001-US25224 A1 20010810				
OTHER SOURCE(S): MARPAT 136:200190				



AB Title compds. I [X, Y = heteroatoms wherein at least X or Y is (un)substituted nitrogen; Z = C, two C atoms or a heteroatom; R = alkyl, aromatic ring, carbocyclic aliphatic ring, halo, haloalkyl, heteroalkyl, heterocyclyl, H, OH, NH2, SH, OCH3; R1-2 = alkyl, aromatic, carbocyclic aliphatic, halo(alkyl), heteroalkyl, heterocyclic aliphatic, H; L = C10-A-NR3, NR3-A-C10, R3N-A-C10-A-NR3; A = alkyl, bond; R3 = alkyl, aromatic ring, carbocyclic aliphatic ring, haloalkyl, heteroalkyl, heterocyclyl, H; B = alkyl, haloalkyl, heteroalkyl, bond; G = nil or a substituent that links R4-5 into a cyclic ring structure; if G is nil, R4 = alkyl-carboxy, aryl-carboxy, etc.; R5 = H, alkyl, aromatic ring, carbocyclic aliphatic ring, halo(alkyl), heteroalkyl, heterocyclic aliphatic ring, etc.; R6 = alkyl, aromatic ring, carbocyclic aliphatic ring, halo, haloalkyl, heteroalkyl, lower heteroalkyl, etc.] were prepared. For instance, 5-benzimidazolecarboxylic acid was coupled to L-phenylalanine benzyl ester (DMF, EDAC, HOBT, Et3N) and the resulting amide debenzylated (MeOH, H2-Pd/C) to give II. Comps. I are useful for the treatment and prevention of diseases and conditions associated with undesirable or abnormal inflammatory responses, such as ischemia-reperfusion injury.


IT 401791-70-6P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 RN 401791-70-6 CAPLUS
 CN L-Phenylalanine, N-(6-quinoxalylcarbonyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 401791-58-0P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (intermediate; benzimidazoles and analogs and use as neutrophil inhibitors)
 RN 401791-58-0 CAPLUS


Absolute stereochemistry



L13 ANSWER 50 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2002:107059 CAPLUS
DOCUMENT NUMBER: 136:151182
TITLE: Antimicrobial biaryl compounds
INVENTOR(S): Jefferson, Elizabeth Ann; Swazey, Eric
PATENT ASSIGNEE(S): Iria Pharmaceutical, Inc., USA
SOURCE: PCT Int. Appl., 44 pp.
CODEN: PIXKd2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002009648	A2	200202027	WO 2001-US24067	20010801
WO 2002009648	A3	20020627		
N: AS, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, DE, DK, DM, DZ, EC, EG, ES, FI, GB, GR, GU, HK, HR, HU, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MK, MN, MM, KN, MX, NZ, NL, PL, PT, RU, RO, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RM: OH, GM, KR, LS, MN, NZ, SD, SL, SZ, TZ, UG, UZ, WT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, GU, HK, HR, HU, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MK, MN, MM, KN, MX, NZ, NL, PL, PT, RU, RO, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
DE, DK, ES, FI, FR, GB, GR, GU, HK, HR, HU, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MK, MN, MM, KN, MX, NZ, NL, PL, PT, RU, RO, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
BJ, CP, CO, CI, CN, CA, GN, OM, QG, MM, KE, NE, NG, SD, TG				
US 6849660	B1	20050201	US 2000-630122	20000801
CA 2418121	AA	20020207	CA 2001-2418121	20010801
AU 2001080944	A5	20020213	AU 2001-80944	20010801
EP 1350528	A2	20030502	EP 2001-959380	20010801
R: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, GU, HK, HR, HU, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MK, MN, MM, KN, MX, NZ, NL, PL, PT, RU, RO, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
JP 2004451941	L2	20040702	JP 2002-515203	20010801
PRIORITY/APPL. INFO.:			US 2000-630122	A 20000801
			WO 2001-US24067	N 20010801
OTHER SOURCE(S):				
MARPAT 136:151182				
0				

OTHER SOURCE(S): MARPAT 136:151182
GI




AB Biaryl I (X = CH, O, S, N; Y = CH₂, NH; Z, n = 0, 1; one of R1 and R2 = (un)substituted CONR2H, COOCH₃, CH=CH₂, SO₂NH₂ and the other is H or R3; one of R5 and R6 = NHCO₂R, NHSO₂R², NH₂(O)R² and the other is H, R4, O = amino acid or peptide residue; R² = H, halogen, (un)substituted NH₂, NO₂, R³ = H, halogen, (un)substituted amino, carboxyl, alkyl, alkenyl, alkynyl; R⁷ = H, amino, (un)substituted alkyl, alkenyl, alkynyl; 5-16 member carbocyclic or heterocyclic) were prepd for use as antiarrhythmic agents. Thus, polymer-supported piperazine was acylated with 5-Bromo-2-thiophenecarboxylic acid, coupled with 3-HMBC6448(GN) 2, and acylated with 2,3-bis(4-phenylthienyl)carboxylic acid to give the biaryl II. In a coupled bacterial transcription-translation assay II had an IC₅₀ of 25 µM.

IT 395648-38-1P
RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); THU
(Therapeutic use); BIOL (Biological study); PREP (Preparation); USES

(preparation of acylaminobiarylcarboxamides as bactericides)
RN 395648-38-1 CAPLUS



6-Quinoxalinecarboxamide, 1,2,3,4-tetrahydro-2,3-dioxo-N-[2-oxo-2-[(3-{5-(1-piperazinylcarbonyl)-2-thienyl}phenyl)amino]ethyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 51 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2001:863510 CAPLUS
DOCUMENT NUMBER: 136:5913
TITLE: Preparation of substituted N-[[aminoiminomethyl or
aminoethyl]phenyl]propyl amides as Factor Xa
inhibitors
INVENTOR(S): Klein, Scott I.; Guertin, Kevin R.; Spada, Alfred P.;
Pauls, Heinz W.; Gong, Yong; Mcgary, Daniel G.
PATENT ASSIGNEE(S): Aventis Pharmaceuticals Products Inc., USA
SOURCE: U.S., 131 pp., Cont.-in-part of U.S. Ser. No. 884,405.
CODEN: USXXAM
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 5

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6323227	B1	20011127	US 1999-259258	19990226
US 6080762	A	20000627	US 1997-884405	19970627
WO 9900515	A1	19990107	WO 1998-013550	19980626
WM: AL, AM, AT, AU, AZ, BA, BE, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GR, HU, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MX, MY, NZ, PL, PT, RU, RO, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW	LS, MM, BD, SZ, UO, ZN, AT, BS, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, SE, SF, BJ, CF, CO, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
PRIORITY APPLN. NO. :			US 1996-9485P	P 19960102
			US 1997-884405	A2 19970627
			WO 1998-013550	A1 19980626
			WO 1999-013570	A1 19990627


OTHER SOURCE(S) : MARPAT 136:5913
GI



AB Title compds. [R = H, OH, NH₂; R₁ = R₂ = H; or R₁R₂ = -NH-; R₃ = H, CO₂R₆, CO₂H, CONH₂, CH₂OR₇, CH₂NH₂; R₄ = H, alkyl, alkyloxy, thioetherocyclyl, CH₂(CH₂)nAr, CH(CH₃)nAr, CH₂Ar; R₅ = alk(en)ymyl, cycloalk(en)yl, heterocyclyl(aryl), aryl, heteroaryl, fused systems, etc.; R₆ = H, lower alkyl; R₇ = H, lower alkyl, aralkyl, heteroaryl, aryl, heteroaryl; R₈ = H, alkyl, aralkyl, heteroaryl, aryl, heteroaryl, COOCH₃, OCH₃, lower alkyl, C₂N₂, Y¹Y²-B; R₉ = alkyl, aralkyl, heteroaryl, aryl, Y¹, Y² = H, alkyl; O = R⁷, R⁷S, Y¹Y₂N; Y¹, Y² = H, alkyl, aryl, aralkyl; or one of Y¹ and Y² = aryl or heteroaryl and the other is as given; Ar = aryl or heteroaryl; the products are useful as intermediates for the synthesis of drugs, prodrugs, N-oxides, hydrates, and solvates are useful as Factor Xa inhibitors. For example, 4-(pyridin-3-yl)benzoic acid was amidated with tert-butyl 3-amino-propionate-HCl via an acid chloride, and the resulting pyrrolidine amide was converted to a sequence of (1) 4-(pyridin-3-yl)-N-tert-butyl-L-glutamate, (2) 4-(pyridin-3-yl)-L-glutamate, (3) L-glutamic acid, (4) L-glutamine, (5) L-glutamic acid, (6) L-glutamic acid, (7) L-glutamic acid, (8) L-glutamic acid, (9) L-glutamic acid, (10) L-glutamic acid, (11) L-glutamic acid, (12) L-glutamic acid, (13) L-glutamic acid, (14) L-glutamic acid, (15) L-glutamic acid, (16) L-glutamic acid, (17) L-glutamic acid, (18) L-glutamic acid, (19) L-glutamic acid, (20) L-glutamic acid, (21) L-glutamic acid, (22) L-glutamic acid, (23) L-glutamic acid, (24) L-glutamic acid, (25) L-glutamic acid, (26) L-glutamic acid, (27) L-glutamic acid, (28) L-glutamic acid, (29) L-glutamic acid, (30) L-glutamic acid, (31) L-glutamic acid, (32) L-glutamic acid, (33) L-glutamic acid, (34) L-glutamic acid, (35) L-glutamic acid, (36) L-glutamic acid, (37) L-glutamic acid, (38) L-glutamic acid, (39) L-glutamic acid, (40) L-glutamic acid, (41) L-glutamic acid, (42) L-glutamic acid, (43) L-glutamic acid, (44) L-glutamic acid, (45) L-glutamic acid, (46) L-glutamic acid, (47) L-glutamic acid, (48) L-glutamic acid, (49) L-glutamic acid, (50) L-glutamic acid, (51) L-glutamic acid, (52) L-glutamic acid, (53) L-glutamic acid, (54) L-glutamic acid, (55) L-glutamic acid, (56) L-glutamic acid, (57) L-glutamic acid, (58) L-glutamic acid, (59) L-glutamic acid, (60) L-glutamic acid, (61) L-glutamic acid, (62) L-glutamic acid, (63) L-glutamic acid, (64) L-glutamic acid, (65) L-glutamic acid, (66) L-glutamic acid, (67) L-glutamic acid, (68) L-glutamic acid, (69) L-glutamic acid, (70) L-glutamic acid, (71) L-glutamic acid, (72) L-glutamic acid, (73) L-glutamic acid, (74) L-glutamic acid, (75) L-glutamic acid, (76) L-glutamic acid, (77) L-glutamic acid, (78) L-glutamic acid, (79) L-glutamic acid, (80) L-glutamic acid, (81) L-glutamic acid, (82) L-glutamic acid, (83) L-glutamic acid, (84) L-glutamic acid, (85) L-glutamic acid, (86) L-glutamic acid, (87) L-glutamic acid, (88) L-glutamic acid, (89) L-glutamic acid, (90) L-glutamic acid, (91) L-glutamic acid, (92) L-glutamic acid, (93) L-glutamic acid, (94) L-glutamic acid, (95) L-glutamic acid, (96) L-glutamic acid, (97) L-glutamic acid, (98) L-glutamic acid, (99) L-glutamic acid, (100) L-glutamic acid, (101) L-glutamic acid, (102) L-glutamic acid, (103) L-glutamic acid, (104) L-glutamic acid, (105) L-glutamic acid, (106) L-glutamic acid, (107) L-glutamic acid, (108) L-glutamic acid, (109) L-glutamic acid, (110) L-glutamic acid, (111) L-glutamic acid, (112) L-glutamic acid, (113) L-glutamic acid, (114) L-glutamic acid, (115) L-glutamic acid, (116) L-glutamic acid, (117) L-glutamic acid, (118) L-glutamic acid, (119) L-glutamic acid, (120) L-glutamic acid, (121) L-glutamic acid, (122) L-glutamic acid, (123) L-glutamic acid, (124) L-glutamic acid, (125) L-glutamic acid, (126) L-glutamic acid, (127) L-glutamic acid, (128) L-glutamic acid, (129) L-glutamic acid, (130) L-glutamic acid, (131) L-glutamic acid, (132) L-glutamic acid, (133) L-glutamic acid, (134) L-glutamic acid, (135) L-glutamic acid, (136) L-glutamic acid, (137) L-glutamic acid, (138) L-glutamic acid, (139) L-glutamic acid, (140) L-glutamic acid, (141) L-glutamic acid, (142) L-glutamic acid, (143) L-glutamic acid, (144) L-glutamic acid, (145) L-glutamic acid, (146) L-glutamic acid, (147) L-glutamic acid, (148) L-glutamic acid, (149) L-glutamic acid, (150) L-glutamic acid, (151) L-glutamic acid, (152) L-glutamic acid, (153) L-glutamic acid, (154) L-glutamic acid, (155) L-glutamic acid, (156) L-glutamic acid, (157) L-glutamic acid, (158) L-glutamic acid, (159) L-glutamic acid, (160) L-glutamic acid, (161) L-glutamic acid, (162) L-glutamic acid, (163) L-glutamic acid, (164) L-glutamic acid, (165) L-glutamic acid, (166) L-glutamic acid, (167) L-glutamic acid, (168) L-glutamic acid, (169) L-glutamic acid, (170) L-glutamic acid, (171) L-glutamic acid, (172) L-glutamic acid, (173) L-glutamic acid, (174) L-glutamic acid, (175) L-glutamic acid, (176) L-glutamic acid, (177) L-glutamic acid, (178) L-glutamic acid, (179) L-glutamic acid, (180) L-glutamic acid, (181) L-glutamic acid, (182) L-glutamic acid, (183) L-glutamic acid, (184) L-glutamic acid, (185) L-glutamic acid, (186) L-glutamic acid, (187) L-glutamic acid, (188) L-glutamic acid, (189) L-glutamic acid, (190) L-glutamic acid, (191) L-glutamic acid, (192) L-glutamic acid, (193) L-glutamic acid, (194) L-glutamic acid, (195) L-glutamic acid, (196) L-glutamic acid, (197) L-glutamic acid, (198) L-glutamic acid, (199) L-glutamic acid, (200) L-glutamic acid, (201) L-glutamic acid, (202) L-glutamic acid, (203) L-glutamic acid, (204) L-glutamic acid, (205) L-glutamic acid, (206) L-glutamic acid, (207) L-glutamic acid, (208) L-glutamic acid, (209) L-glutamic acid, (210) L-glutamic acid, (211) L-glutamic acid, (212) L-glutamic acid, (213) L-glutamic acid, (214) L-glutamic acid, (215) L-glutamic acid, (216) L-glutamic acid, (217) L-glutamic acid, (218) L-glutamic acid, (219) L-glutamic acid, (220) L-glutamic acid, (221) L-glutamic acid, (222) L-glutamic acid, (223) L-glutamic acid, (224) L-glutamic acid, (225) L-glutamic acid, (226) L-glutamic acid, (227) L-glutamic acid, (228) L-glutamic acid, (229) L-glutamic acid, (230) L-glutamic acid, (231) L-glutamic acid, (232) L-glutamic acid, (233) L-glutamic acid, (234) L-glutamic acid, (235) L-glutamic acid, (236) L-glutamic acid, (237) L-glutamic acid, (238) L-glutamic acid, (239) L-glutamic acid, (240) L-glutamic acid, (241) L-glutamic acid, (242) L-glutamic acid, (243) L-glutamic acid, (244) L-glutamic acid, (245) L-glutamic acid, (246) L-glutamic acid, (247) L-glutamic acid, (248) L-glutamic acid, (249) L-glutamic acid, (250) L-glutamic acid, (251) L-glutamic acid, (252) L-glutamic acid, (253) L-glutamic acid, (254) L-glutamic acid, (255) L-glutamic acid, (256) L-glutamic acid, (257) L-glutamic acid, (258) L-glutamic acid, (259) L-glutamic acid, (260) L-glutamic acid, (261) L-glutamic acid, (262) L-glutamic acid, (263) L-glutamic acid, (264) L-glutamic acid, (265) L-glutamic acid, (266) L-glutamic acid, (267) L-glutamic acid, (268) L-glutamic acid, (269) L-glutamic acid, (270) L-glutamic acid, (271) L-glutamic acid, (272) L-glutamic acid, (273) L-glutamic acid, (274) L-glutamic acid, (275) L-glutamic acid, (276) L-glutamic acid, (277) L-glutamic acid, (278) L-glutamic acid, (279) L-glutamic acid, (280) L-glutamic acid, (281) L-glutamic acid, (282) L-glutamic acid, (283) L-glutamic acid, (284) L-glutamic acid, (285) L-glutamic acid, (286) L-glutamic acid, (287) L-glutamic acid, (288) L-glutamic acid, (289) L-glutamic acid, (290) L-glutamic acid, (291) L-glutamic acid, (292) L-glutamic acid, (293) L-glutamic acid, (294) L-glutamic acid, (295) L-glutamic acid, (296) L-glutamic acid, (297) L-glutamic acid, (298) L-glutamic acid, (299) L-glutamic acid, (300) L-glutamic acid, (301) L-glutamic acid, (302) L-glutamic acid, (303) L-glutamic acid, (304) L-glutamic acid, (305) L-glutamic acid, (306) L-glutamic acid, (307) L-glutamic acid, (308) L-glutamic acid, (309) L-glutamic acid, (310) L-glutamic acid, (311) L-glutamic acid, (312) L-glutamic acid, (313) L-glutamic acid, (314) L-glutamic acid, (315) L-glutamic acid, (316) L-glutamic acid, (317) L-glutamic acid, (318) L-glutamic acid, (319) L-glutamic acid, (320) L-glutamic acid, (321) L-glutamic acid, (322) L-glutamic acid, (323) L-glutamic acid, (

of the MEM group, and conversion to the Me ester, (3) Pd-catalyzed cyanation of the iodide, and (4) Pinna reaction and ammonolysis of the nitrile, to give title compound II. Three example compds. showed KI values of 19.0-94.0 nM in a Factor Xa assay, 46 nM to 1.72 μ M in a trypsin assay, and 477 nM to 2.71 μ M in a thrombin assay.

IT	<p>85667; 860 477 nm to 2.71 μm in a threonine assay.</p> <p>21567 20-69</p> <p>RL: RAC (Biological activity) or effector, except adeney; BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PRSP (Preparation); USES (Uses) (preparation of substituted (aminomethyl)- or (aminomethyl)phenylpropyl amides as Factor Xc inhibitors)</p>
RN	219673-02-6 CAPLUS
CN	<p>Benzenesulfonamide, 3-(aminomethyl)-, 1-[1-(6-guineazyl)-guinea-benzylamino]ethyl-, ester (9CI) (CA INDEX NAME)</p>



REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT.


L13 ANSWER 52 OF 181 CAPLUS COPYRIGHT 2006 ACS on STM
 ACCESSION NUMBER: 2001:587526 CAPLUS
 DOCUMENT NUMBER: 135:1318478
 TITLE: Traceless, Self-Cleaving Solid- and Solution-Phase Parallel Synthesis of 3,4,7-Trisubstituted 3,4-Dihydroquinoxalin-2-ones
 AUTHOR(S): Laborde, Edgardo; Peterson, Brian T.; Robinson, Louise
 CORPORATE SOURCE: Telik Inc., South San Francisco, CA, 94080, USA
 SOURCE: Journal of Combinatorial Chemistry 2001, 3(6), 572-577
 CODEN: JCCHEP; ISSN: 1520-4766
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 135:1318478

OTHER SOURCE(S) CASREACT 135319474
 AS This article describes the synthesis of 3,4-dihydroquinazolin-2-one for the parallel synthesis of
 3,4-dihydroquinazolin-2-one containing three points of diversity. The
 synthesis begins with com. available resin-bound α -amino acid
 as the source of the diversity element and employs a reaction of
 and solution-phase chemistry to introduce the other two. The key step is
 an intramol. cyclization and simultaneous traceless cleavage from the
 solid support to give a disubstituted 3,4-dihydroquinazolin-2-one. The
 resulting substituted resin is solubilized in solution and the amino
 nitrogen using a scavenger resin to dispose of excess reagent. All the
 reactions in the sequence take place at room temperature without the need to

strong acids or to maintain an inert atmospheric, thereby preserving the chiral integrity of the starting α -amino acid and facilitating the generation of libraries in a high-throughput parallel format.

IT 367941-10-4P
RL: SPN (Synthetic preparation); PREP (Preparation)
(traceless self-cleaving solid- and solution-phase parallel synthesis of
3,4,7-trisubstituted 3,4-dihydroquinoxalin-2-one)s

	3,4,7-trisubstituted 3,4-dihydroquinoxalin-2-ones)
RN	367941-10-4 CAPLUS
CN	6-Quinoxalinecarboxamide, N-[(3-fluorophenyl)methyl]-3,4-dihydro-2-(1-methylethyl)-3-oxo- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 53 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2001:466005 CAPLUS
 DOCUMENT NUMBER: 135:319476
 TITLE: Cold pad-batch dyeing of Lyocell
 AUTHOR(S): Siedow, K.
 CORPORATE SOURCE: DyStar Textilfarben GmbH y Co., Frankfurt am Main, Germany
 SOURCE: Revista de Quimica Textil (2001), 151, 42, 44-46, 48-53
 CODEN: ROTED3; ISSN: 0300-3418
 PUBLISHER: Asociacion Espanola de Quimicos y Coloristas Textiles
 DOCUMENT TYPE: Journal
 LANGUAGE: Spanish

AB Methods to reduce crease and rub marks on Lyocell fabrics and suitable cold pad-batch dyeing processes, i.e., soda and silicate, are described. Finishing methods and reagents for woven fabrics with a peach skin effect (fleece) and with a smooth (non-fibrillated) surface are outlined. Operation conditions for cold pad-batch dyeing of Lyocell were determined and dyed Lyocell was compared to viscose fabrics in terms of color intensity, dye affinity, tailing test, fixing yield, washing-out, etc. Types of cold pad-batch process include soda process for Levafix dyes, water glass methods for Remazol dyes; recommendations are given for dye selection for each type.

IT 206058-73-3, Levafix Brilliant Blue S-B
 RL: NUU (Other use, unclassified); USSS (Uses)
 (dye selection and parameters for cold pad-batch dyeing and finishing of Lyocell)

RN 206058-73-3 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[[4-[[[(2,3-dichloro-6-quinoxalyl)carbonyl]amino]methyl]-3-sulfonyl]methyl]amino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)



PAGE 1-A




PAGE 2-A


L13 ANSWER 54 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2001:214540 CAPLUS
 DOCUMENT NUMBER: 135:30391
 TITLE: Synthesis of [Ru(phen)2dppz]2+-tethered oligo-DNA and studies on the metallointercalation mode into the DNA duplex
 AUTHOR(S): Ossipov, Dimitri; Pradeepkumar, P. I.; Holmer, Melcer; Chattopadhyaya, Jyoti
 CORPORATE SOURCE: Department of Bioorganic Chemistry Biomedical Center, University of Uppsala, Uppsala, Sweden.
 SOURCE: Journal of the American Chemical Society (2001), 123(15), 3551-3562
 CODEN: JACSAT; ISSN: 0002-7863
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 135:30391
 AB To explore the binding properties of [Ru(phen)2dppz]2+ complex (phen = 1,10-phenanthroline, dppz = dipyrido[3,2-a:2',3'-c]phenazine) in a sequence-specific manner in DNA duplex, it was tethered through the dppz ligand to a central position as well as both at the 3'- and 5'-ends of oligodeoxyribonucleotide (ODN). The middle [Ru(phen)2dppz]2+-ODN tethered was resolved and isolated as four pure diastereomers, while the 3'- or 5'-[Ru(phen)2dppz]2+-ODNs were inseparable on RP-HPLC. Thermal stability

of the (Ru2+-ODN)·DNA duplexes is found to increase considerably ($\Delta T_m = 12.6-23.4^\circ$), depending upon the site of the covalent attachment of the tethered [Ru(phen)2dppz]2+ complex, or the chirality of the [Ru(phen)2dppz]2+-linker tethered at the middle of the ODN, compared to the unlabeled counterpart. Gross differences in CD between the [Ru(phen)2dppz]2+-tethered and the native DNA duplexes showed that the global duplex conformation of the former has considerably altered from the B-type, but is still recognized by DNase I. The thermal melting studies, CD measurements, as well as DNase I digestion data, are interpreted as a result of intercalation of the dppz moiety, which is realized by threading of the Ru(phen)2 complex part through the DNA duplex core. DNase I footprinting with four diastereomerically pure middle [(Ru(phen)2dppz]2+-ODN)·DNA duplexes furthermore showed that the tethered [Ru(phen)2dppz]2+-linker chirality dictates the stereochem. accessibility of various phosphodiester moieties (around the intercalation site) toward the cleavage reaction by the enzyme. The diastereomerically pure ruthenium-modified duplexes, with the well-defined π -stack, will be useful to explore stereochem.-dependent energy- and electron-transfer chemical to understand oxidative damage to the DNA double helix as well as the long-range energy- and electron-transfer processes with DNA as a reactant.

IT 342906-42-7p 342906-43-8p 342906-45-0p
 342906-46-1p
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (synthesis of [Ru(phen)2dppz]2+-tethered oligodeoxyribonucleotides)
 RN 342906-42-7 CAPLUS
 CN Ruthenium(2+), [N-[[11-hydroxy-14,14-bis(4-methoxyphenyl)-14-phenyl-3,6,9,13-tetraoxatetradec-1-yl]dipyrido[3,2-a:2',3'-c]phenazine-11-carboxamide- κ N4, κ N5]bis(1,10-phenanthroline- κ N1, κ N10)-, (OC-6-33)- (9CI) (CA INDEX NAME)




PAGE 1-A




PAGE 1-B

RN 342906-43-8 CAPLUS
 CN Ruthenium(2+), [1-[[bis(4-methoxyphenyl)phenylmethoxy]methyl]-13-(dipyrido[3,2-a:2',3'-c]phenazin-11-yl)- κ N4, κ N5]-13-oxo-3,6,9-trioxo-12-azatridec-1-yl] P-(2-cyanoethyl)-N,N-bis(1-methylethyl)phosphonamidite]bis(1,10-phenanthroline- κ N1, κ N10)-, (OC-6-33)- (9CI) (CA INDEX NAME)


PAGE 1-A



PAGE 1-B



RN 342906-45-0 CAPLUS
 CN Ruthenium(1+), [mono[1-[[bis(4-methoxyphenyl)phenylmethoxy]methyl]-13-(dipyrido[3,2-a:2',3'-c]phenazin-11-yl)- κ N4, κ N5]-13-oxo-3,6,9-trioxo-12-azatridec-1-yl] butanedioate]bis(1,10-phenanthroline- κ N1, κ N10)-, (OC-6-33)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 48 THERE ARE 48 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 55 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2001:64269 CAPLUS
 DOCUMENT NUMBER: 134:128208
 TITLE: Detection of biomolecules by sensitizer-linked
 substrates for biomolecules
 INVENTOR(S): Gray, Harry B.; Crane, Brian R.; Winkler, Jay R.;
 Dmochowski, Ivan Julian; Wilker, Jonathan J.; Dunn,
 Alexander Robert
 PATENT ASSIGNER(S): California Institute of Technology, USA
 SOURCE: PCT Int. Appl. 174 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

[illegible]


US 2000-192703P P 20000328

AB Methods and compns. for detecting and characterizing target biomols. using sensitizer-linked substrate mols. are disclosed. High throughput screening assays and therapeutic applications of the inventions are also included. Substrate-sensitizer compds. adamantane (or ethylbenzene or imidazole) linked via a C9-C13 chain to [Ru(bpy)₃]²⁺ or similar sensitizer were prepared, studied and tested with cytochrome P 450.

IT 321835-64-7P 321835-65-8P
RL: ARG (Analytical reagent use); BPR (Biological process); BSU
(Biological study, unclassified); RCT (Reactant); SPN (Synthetic
preparation); ANST (Analytical study); BIOL (Biological study); PREP
(Preparation); PROC (Process); RACT (Reactant or reagent); USBS (Uses)
(detection of biomole. by sensitizer-linked substrates for biomole.)

RN 321835-64-7 CAPLUS
 CN Ruthenium(2+), [N-[6-oxo-6-(tricyclo[3.3.1.1^{3,7}]dec-2-ylamino)hexyl]dipyrido[3,2-a:2',3'-c]phenazine-11-carboxamide-κN4,κN5]bis(1,10-phenanthroline-κN1,κN10)-, (OC-6-33)- (9CI) (CA INDEX NAME)

quinoxaliny]carbonyl]amino]methyl]-3-sulfo]phenyl]methyl]amino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)



●2 No


REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 57 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STM
 ACCESSION NUMBER: 2000:729220 CAPLUS
 DOCUMENT NUMBER: 134:18537
 TITLE: Dimerization of Cibacron Blue F 3GA and other dyes:
 influence of salts and temperature
 AUTHOR(S): Alberghina, Gaetano; Bianchini, Roberto; Fichera,
 Maria; Fischella, Salvatore
 CORPORATE SOURCE: Dipartimento di Scienze Chimiche, Universita di
 Catania, Catania, 95125, Italy
 SOURCE: Dyes and Pigments (2000), 46(3), 129-137
 CODEN: DYPIDX; ISSN: 0143-7208
 PUBLISHER: Kluwer Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English


AB The monomer-dimer equilibrium of Cibacron Blue F 3QA (CB) and five other dyes (Levafix Brilliant Blue EB, Reactive Scarlet 017, Methyl orange, Basic Blue 3, and Chicago Blue Sky) have been investigated in water and in the presence of KH₂PO₄. Aggregation of CB has been also examined in the presence of NaH₂PO₄, LiCl, and KCl. When a new iterative approach, based

RN 342906-46-1 CAPLUS
CN Ruthenium(2+), [N-[2-[2-[2-(2,3-dihydroxypropoxy)ethoxy]ethoxy]ethyl]dipyrido[3,2-a:2',3'-c]phenazine-11-carboxamide- kN4, kN5]bis[1,10-phenanthroline- kN1, kN10]-, (OC-6-33)- (9CI) (CA INDEX NAME)

PAGE 1-A



RN 321835-65-8 CAPLUS
CN Ruthenium(2+), [N-[2-oxo-2-(tricyclo[3.3.1.1^{3,7}]dec-2-ylamino)ethyl]dipyrido[3,2-a:2',3'-c]phenazine-11-carboxamide-κN4,κN5]bis(1,10-phenanthroline-κN1,κN10)-, (OC-6-33)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 56 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 2000:739213 CAPLUS
DOCUMENT NUMBER: 134:72834
TITLE: Lycopell: cold pad-batch dyeing process. Part 2
AUTHOR(S): Siedow, K.
CORPORATE SOURCE: DyStar Textilfabren GMBH & Co., Frankfurt Am Main,
Germany
SOURCE: Tinctoria (2000), 97(8), 31-36
CODEN: TINCAM; ISSN: 0040-7984
Edition: Ariminum
Journal
Language: Italian

LANGUAGE: Italian

AB Operation conditions for cold pad-batch dyeing of Lyocell were determined and dyed lyocell was compared to viscose fabrics in terms of color intensity and dye affinity. Sulfonated disperse dyes and disperse dyes were used. Types of cold pad-batch process include soda process for Levafix dyes, waterless process for methods for Remazol dyes; recommendations are given for dye selection for each type.


IT 206058-73-3, Levafix Brilliant Blue R-B
RL: NUU (Other use, unclassified); USES (Uses)
(soda and silicate cold pad-batch dyeing of Lyocell with Levafix and
Remazol dyes and quality parameters of dyed fabrics)

2-ANTRACENESULFONIC ACID, 1-AMINO-4-[[[4-[[[2,3-DICHLORO-6-

on non-linear least-square fitting procedure was applied. it was found that the dimerization conste. depend on the extension of organic mole. and the number of sulfonic groups. In the case of CB, cations had a greater effect on the equilibrium than anions. Anal. of the calculated spectra for monomer and dimer of Basic Blue 3 after deconvolution allowed us to specify the geometry of the dimer.

IT 206058-73-3, Levafix Brilliant Blue BB
 RL: PRP (Properties); TEM (Technical or engineered material use); USES (Uses)
 (Influence of salts and temperature on aqueous associative dimerization of dyes)
 RN 206058-73-3 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[[4-[[[2,3-dichloro-6-quinoxaliny]carbonyl]amino]methyl]-3-sulfonyl]methyl]amino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A



REFERENCE COUNT: 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 58 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2000:700553 CAPLUS
 DOCUMENT NUMBER: 134:4913
 TITLE: Synthesis of some sulfonamide derivatives with potential antibacterial activity

AUTHOR(S): El-Din, Nabaweya Sharaf
 CORPORATE SOURCE: Faculty of Pharmacy, University of Tanta, Egypt
 SOURCE: Chemistry of Heterocyclic Compounds (New York) (Translation of Khimiya Geterotsiklicheskikh Soedinenii) (2000), 36(4), 449-454
 CODEN: CHCCAL; ISSN: 0009-3122
 PUBLISHER: Consultants Bureau
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 134:4913
 AB Some new quinoxaline-6-sulfonamide and phthalazine-6-sulfonamide deriva. were synthesized in 61-68% yields by treating the corresponding quinoxaline- and phthalazine-6-sulfonyl chlorides with the appropriate amine (PrNH2, 1-amino-2-propanol, glycine, p-H2NC6H4CO2H, morpholine, piperazine). The majority of the prepared compds. showed antibacterial activity.
 IT 112170-26-0P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation and antibacterial activity of)
 RN 112170-26-0 CAPLUS
 CN Glycine N-[[1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny]sulfonyl]- (9CI) (CA INDEX NAME)





REFERENCE COUNT: 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 59 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2000:433346 CAPLUS
 DOCUMENT NUMBER: 133:73861
 TITLE: Preparation of α-amidinobenzyl-β-(aroylamino)alkanoates and analogs as factor Xa inhibitors
 INVENTOR(S): Klein, Scott I.; Guertin, Kevin R.; Spada, Alfred P.
 PATENT ASSIGNOR(S): Aventis Pharmaceuticals Products, Inc., USA
 SOURCE: U.S., 118 pp., Cont.-in-part of U.S. 9724118.
 CODEN: USXKAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 5
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6080767	A	20000627	US 1997-884405	19970627
WO 9724118	A1	19970710	WO 1996-US20770	19961223
W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KO, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SF, BJ, CP, CO, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2264556	AA	19990107	CA 1998-2264556	19980626

WO 9900356 A1 19990107 WO 1998-US13550 19980626
 W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KO, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW
 RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SF, BJ, CP, CO, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG
 AU 9881771 A1 19990119 AU 1998-81771 19980626
 AU 741173 B2 20011122
 EP 931060 A1 19990728 EP 1998-931728 19980626
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO
 BR 9806060 A 19990831 BR 1998-6060 19980626
 JP 2001500532 T2 20010116 JP 1999-505870 19980626
 AP 1061 A 20020424 AP 1999-1467 19980626
 W: GH, GM, KE, LS, MW, SD, SZ, UG, ZW
 ZA 9805664 A 19990113 ZA 1998-5664 19980629
 NO 9900854 A 19990423 NO 1999-854 19990223
 NO 314758 B1 20030519
 US 6123227 B1 20011127 US 1999-259528 19990226
 US 6277865 B1 20010821 US 1999-273618 19990222
 US 1996-9485P P 19960102
 WO 1996-US20770 A2 19961223
 US 1997-884405 A 19970627
 US 1998-79002P P 19980321
 WO 1998-US13550 W 19980626
 OTHER SOURCE(S): MARPAT 133:73861
 GI



AB H2NC1R12CH2CH2CH2CH2CH2NRCORS [R1,R2 = H; R1R2 = NR9; R3 = H, COR6, CO2R6, COR(R6)2, CH2OR7, CH2SR7; R4 = H, (hydroxy)alkyl, aminoalkyl, (CH2CH2)nR, (CH)CHnR, CH2R; R = (un)substituted (hetero)aryl; R5 = (ar)alk(en)yl, heterocyclyl, (hetero)aryl, etc.; R6,R8 = H or alkyl; R7 = H, alkyl, acyl, (hetero)aryl, etc.; R9 = H, OH, alkoxy(carbonyl), alkanoyl, etc.; Z = phenylene; n = 0-2] were prepared as factor Xa inhibitors (no date). Thus, 4-(ND)C6H4CH:CHCO2Me was cyclcondensed with 4-(MeO)C6H4N:CHCH:CHPh (preparation (pre) given) to give, after N-deprotection, β-lactam I. The latter was N-acylated by 4-PHC6H4COCl and the product hydrolyzed to give, after amination/esterification, title compound II.

IT 219673-02-6P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of α-amidinobenzyl-β-(aroylamino)alkanoates and analogs as factor Xa inhibitors)
 RN 219673-02-6 CAPLUS
 CN Benzenepropanoic acid, 3-(aminoinomethyl)-, n-[[1-[[6-quinoxaliny]carbonyl]amino]ethyl]-, methyl ester (9CI) (CA INDEX NAME)




REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L13 ANSWER 60 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 2000:267974 CAPLUS
 DOCUMENT NUMBER: 133:4414
 TITLE: Synthesis and Assembly of Self-Complementary Cavitands
 AUTHOR(S): Renslo, Adam R.; Tucci, Fabio C.; Rudkevich, Dmitry M.; Rebek, Julius, Jr.
 CORPORATE SOURCE: Skaggs Institute for Chemical Biology and the Department of Chemistry, Scripps Research Institute, La Jolla, CA, 92037, USA
 SOURCE: Journal of the American Chemical Society (2000), 122(19), 4573-4582
 CODEN: JACSAT; ISSN: 0002-7863
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Cavitands with self-complementary shapes were prepared by the covalent attachment of adamantane guest moieties to the upper rim of the host structures. Relatives of the "self-folding" cavitands, these new structures possess a seam of intramolecular hydrogen bonds that stabilize the folded conformation. Their self-complementary shapes result in the formation of noncovalent dimers of considerable kinetic and thermodyn. stability (-ΔG295 = 4.5-6.5 kcal/mol in p-xylene-d10). The dimerization of the cavitands is reversible and subject to control by solvent and temperature. The dimerization process is enthalpically favored and entropy opposed and occurs with significant enthalpy-entropy compensation.
 IT 108229-81-8P 242129-41-5P 270563-44-5P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (arylation; synthesis and inclusion dimerization of self-complementary cavitands bearing adamantyl recognition sites)


RN 108229-81-8 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, phenylmethyl ester (9CI) (CA INDEX NAME)



RN 242129-41-5 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(tricyclo[3.3.1.1.3,7]dec-1-ylmethyl)- (9CI) (CA INDEX NAME)



RN 270563-44-5 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, tricyclo[3.3.1.1.3,7]dec-1-ylmethyl ester (9CI) (CA INDEX NAME)




IT 270563-41-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(hydrogenolysis, synthesis and inclusion dimerization of self-complementary cavittands bearing adamantyl recognition sites)

RN 270563-41-2 CAPLUS
CN 13,23:14,22-Dimetheno-15H,17H,19H,21H-benzo[2',3']benzo[2'',3''] [1,7]benzo dioxonino[3''',2''':9'',10''] [1,4]benzodioxonino[6'',5'':9'',10''] [1,4]benzo dioxonino[6'',5'':9,10] [1,4]benzodioxonino[2,3-b]quinoxaline-27-carboxylic acid, 2,3,9,10,35,36-hexakis[(1-oxooctyl)amino]-15,17,19,21-tetraundecyl-, phenylmethyl ester, (15R,17S,19R,21S)-rel- (9CI) (CA INDEX NAME)

PAGE 1-A

Me- (CH₂)₆-


PAGE 1-B




PAGE 1-C

Me- (CH₂)₆-Me

PAGE 2-A



PAGE 2-B



PAGE 3-B

(CH₂)₁₀-Me

IT 270563-45-6P
RL: PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)
(no dimerization; synthesis and inclusion dimerization of self-complementary cavittands bearing adamantyl recognition sites)

RN 270563-45-6 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2-chloro-3-[(24R,26R,28R,34S)-31-hydroxy-2,3,9,10,16,17-hexakis[(1-oxooctyl)amino]-24,26,28,34-tetraundecyl-22,30-methano-24H,26H,28H-tetrabenzo[b,b',e,e'] [1,7]benzodioxonino[3,2-j:5,6-j']bis[1,4]benzodioxonin-21-yl]oxy]-, phenylmethyl ester, rel- (9CI) (CA INDEX NAME)


*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

IT 242143-99-3P
RL: PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)
(prepared and characterized in present paper and mis-assigned in earlier paper; synthesis and inclusion dimerization of self-complementary cavittands bearing adamantyl recognition sites)



RN 242143-99-3 CAPLUS
CN 13,23:14,22-Dimetheno-15H,17H,19H,21H-benzo[2',3']benzo[2'',3''] [1,7]benzo dioxonino[3''',2''':9'',10''] [1,4]benzodioxonino[6'',5'':9'',10''] [1,4]benzo dioxonino[6'',5'':9,10] [1,4]benzodioxonino[2,3-b]quinoxaline-27-carboxamide, 2,3,9,10,35,36-hexakis[(1-oxooctyl)amino]-N-(tricyclo[3.3.1.1.3,7]dec-1-ylmethyl)-15,17,19,21-tetraundecyl-, stereoisomer (9CI) (CA INDEX NAME)

PAGE 1-A

Me- (CH₂)₆-



—(CH₂)₆—Me



IT 270563-88-7
 RL: PMU (Formation, unclassified); PRP (Properties); FORM (Formation, nonpreparative)
 (synthesis and inclusion dimerization of self-complementary cavitate bearing adamantyl recognition sites)
 RN 270563-88-7 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2-chloro-3-[[[31-hydroxy-2,3,9,10,16,17-hexakis[(1-oxooctyl)amino]-24,26,28,34-tetraundecyl-22,30-methano-24H,26H,28H-tetrabenzo[b,b',e,e'] [1,7]benzodioxonino[3,2-j:5,6-j']bis[1,4]benzodioxonin-21-yl]oxy]-N-tricyclo[3.3.1.1^{3,7}]dec-1-yl-, stereoisomer, compd. with stereoisomer of 2-chloro-3-[[[31-hydroxy-2,3,9,10,16,17-hexakis[(1-oxooctyl)amino]-24,26,28,34-tetraundecyl-22,30-methano-24H,26H,28H-tetrabenzo[b,b',e,e'] [1,7]benzodioxonino[3,2-j:5,6-j']bis[1,4]benzodioxonin-21-yl]oxy]-N-tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]-6-quinoxalinecarboxamide (1:1) (9CI) (CA INDEX NAME)
 CN 1
 CRN 270563-64-9
 CMF C158 H228 C1 N9 O15
 *** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
 CN 2
 CRN 270563-63-8
 CMF C157 H226 C1 N9 O15
 *** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
 IT 270563-64-9P 270563-65-0P

RL: PREP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)
 (synthesis and inclusion dimerization of self-complementary cavitate bearing adamantyl recognition sites)

RN 270563-64-9 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2-chloro-3-[[[31-hydroxy-2,3,9,10,16,17-hexakis[(1-oxooctyl)amino]-24,26,28,34-tetraundecyl-22,30-methano-24H,26H,28H-tetrabenzo[b,b',e,e'] [1,7]benzodioxonino[3,2-j:5,6-j']bis[1,4]benzodioxonin-21-yl]oxy]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-, stereoisomer (9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

RN 270563-65-0 CAPLUS

CN 6-Quinoxalinecarboxylic acid, 2-chloro-3-[[[31-hydroxy-2,3,9,10,16,17-hexakis[(1-oxooctyl)amino]-24,26,28,34-tetraundecyl-22,30-methano-24H,26H,28H-tetrabenzo[b,b',e,e'] [1,7]benzodioxonino[3,2-j:5,6-j']bis[1,4]benzodioxonin-21-yl]oxy]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl ester, stereoisomer (9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

REFERENCE COUNT: 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 61 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STM

ACCESSION NUMBER: 2000:122112 CAPLUS

DOCUMENT NUMBER: 132:202768

TITLE:

Sequence-Recognition and Cleavage of DNA by a

Netropsin-phenazine-di-N-oxide Conjugate

Hellisey, Philippe; Giorgi-Renault, Sylviane; Colson,

Pierre; Houssier, Claude; Bailly, Christian

Laboratoire de Chimie Therapeutique Faculte des

Sciences Pharmaceutiques et Biologiques, UMR

CMRS-Universite Rene Descartes no. 8638, Paris, 75270,

FR.

Bioconjugate Chemistry (2000), 11(2), 219-227

CODEN: BCCHES; ISSN: 1043-1802

American Chemical Society

Journal

English

AB The synthesis, DNA-binding and cleaving properties, and cytotoxic

activities of R-128, a hybrid mol. in which a bis-pyrrolecarboxamide-

amide element related to the antibiotic netropsin is covalently tethered to

a phenazine-di-N-oxide chromophore was reported. The affinity and mode of

interaction of the conjugate with DNA were investigated by a

combination of absorption spectroscopy, CD, and elec. linear dichroism.

This hybrid mol. binds to AT-rich sequences of DNA via a bimodal process

involving minor groove binding of the netropsin moiety and intercalation

of the phenazine moiety. The bidentate mode of binding was evidenced by

linear dichroism using calf thymus DNA and poly(dA-dT)·(dA-dT). In

contrast, the drug fails to bind to poly(dG-dC)·poly(dG-dC),

because of the obstructive effect of the guanine 2-amino group exposed in

the minor groove of this polynucleotide. DNase I footprinting studies

indicated that the conjugate interacts preferentially with AT-rich

sequences, but the cleavage of DNA in the presence of a reducing agent can

occur at different sequences not restricted to the AT sites. The main

cleavage sites were detected with a periodicity of about 10 base pairs

corresponding to approx. one turn of the double helix. This suggests that

the cleavage may be dictated by the structure of the double helix rather

than the primary nucleotide sequence. The conjugate which is moderately

toxic to cancer cells complements the tool box of reagents which can be

utilized to produce DNA strand scission. The DNA cleaving properties of

R-128 entreat further exploration into the use of phenazine-di-N-oxides as

tools for investigating DNA structure.

IT 260416-09-9P, R 128



RL: BAC (Biological activity or effector, except adverse); BSU (Biological

study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)


(R 128; sequence-recognition and cleavage of DNA by netropsin-phenazine-di-N-oxide conjugate R-128)

RN 260416-09-9 CAPLUS


CN 2-Phenazinecarboxamide, N-[6-[[[5-[[[5-[[[3-amino-3-iminopropyl]amino]carbonyl]-1-methyl-1H-pyrrol-3-yl]amino]carbonyl]-1-methyl-1H-pyrrol-3-yl]amino]-6-oxohexyl]-, 5,10-dioxido, monohydrochloride (9CI) (CA INDEX NAME)




IT 260389-77-3P 260389-78-4P 260389-79-5P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (sequence-recognition and cleavage of DNA by netropsin-phenazine-di-N-oxide conjugate R-128)
 RN 260389-77-3 CAPLUS
 CN Hexanoic acid, 6-[[[2-phenazinylcarbonyl]amino]-, methyl ester (9CI) (CA INDEX NAME)



RN 260389-78-4 CAPLUS
 CN Hexanoic acid, 6-[[[5,10-dioxido-2-phenazinylcarbonyl]amino]-, methyl ester (9CI) (CA INDEX NAME)



RN 260389-79-5 CAPLUS
CN Hexanoic acid, 6-[[5,10-dioxido-2-phenaziny]carbonyl]amino]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 62 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2000:41754 CAPLUS

DOCUMENT NUMBER: 132:208087

TITLE: Dipyrro[3,2-a:2',3'-c]phenazine-tethered oligo-DNA: synthesis and thermal stability of their DNA · DNA and DNA · RNA duplexes and DNA · DNA · DNA triplexes

AUTHOR(S): Ossipov, Dimitri; Zamaratski, Edouard; Chattopadhyaya, Jyoti

CORPORATE SOURCE: Department of Bioorganic Chemistry, Biomedical Center, University of Uppsala, Swed.

SOURCE: Helvetica Chimica Acta (1999), 82(12), 2186-2200

CODEN: HCACAV; ISSN: 0018-019X

PUBLISHER: Verlag Helvetica Chimica Acta

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Dipyrro[3,2-a:2',3'-c]phenazine (dppz) derivs. were conjugated to 9-mer and 18-mer DNA (ODN) at a site without nucleobase, either at the 5'- or 3'-end or at an internucleotide position, via linkers of 7, 12, or 18 atoms lengths. These dppz-linked ODNs were synthesized using novel backbone glycerol phosphoramidites: glycerol, serving as artificial nucleoside without nucleobase, was modified to amines which were suitable for the subsequent key reaction with dppz-carboxylic acid. The products of these reactions were then transformed to the standard phosphoramidite derivs. or used for loading on a CPG support. The dppz-modified ODNs were subsequently assembled in the usual manner using automated solid-phase DNA synthesis. The 9-mer ODN-dppz conjugates were tested for their ability to form stable duplexes with target DNA or RNA strands (D11 or R11) while the 18-mer ODN-dppz conjugates were tested for their ability to form stable triplexes with a DNA target duplex D24 · D24. The presence of the conjugated dppz derivative increases the stability of DNA · DNA and DNA · RNA duplexes, typically by a ΔT_m of 7.3-10.9° and 4.5-7.4°, resp., when the dppz is tethered at the 5'- or 3'-terminal. The dppz derivs. also stabilize triplexes when attached to


the 5'- or 3'-end, with a ΔT_m varying from 3.8-11.1°. The insertion of a dppz building block at the center of a 9-mer results in a considerably poorer stability of the corresponding DNA · DNA duplexes (ΔT_m = 0.5 to 4.2°) and DNA · RNA duplexes (ΔT_m = -1.5 to 0.9°), while the replacement of one interior nucleotide by a dppz building unit in the corresponding 8-mer ODN does not reveal the formation of any duplex at all. Different types of modifications in the middle of the 18-mer ODN, in general, do not lead to any triplex formation, except when the dppz derivative is tethered to the ODN through a 12-atom-long linker.

IT 259796-25-3P 259796-26-4P 259796-27-5P
259796-37-7P 259796-38-8P 259796-40-2DP, CPG
bound 259796-43-5DP, CPG bound 259796-44-6DP, CPG
bound

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(synthesis and thermal stability of dipyrro[phenazine]-tethered oligo-DNA and their DNA/RNA duplexes and triplexes)


RN 259796-25-3 CAPLUS

CN Dipyrro[3,2-a:2',3'-c]phenazine-11-carboxamide, N-[3-[(2,2-dimethyl-1,3-dioxolan-4-yl)methoxy]propyl]- (9CI) (CA INDEX NAME)




RN 259796-26-4 CAPLUS

CN Dipyrro[3,2-a:2',3'-c]phenazine-11-carboxamide, N-[11-hydroxy-14,14-bis(4-methoxyphenyl)-14-phenyl-3,6,9,13-tetraoxatetradec-1-yl]- (9CI) (CA INDEX NAME)




PAGE 1-A

PAGE 1-B




RN 259796-27-5 CAPLUS
CN Dipyrro[3,2-a:2',3'-c]phenazine-11-carboxamide, N-[17-hydroxy-20,20-bis(4-methoxyphenyl)-20-phenyl-3,6,9,12,15,19-hexaoxaicos-1-yl]- (9CI) (CA INDEX NAME)


PAGE 1-A



PAGE 1-B




RN 259796-37-7 CAPLUS
CN Dipyrro[3,2-a:2',3'-c]phenazine-11-carboxamide, N-[3-(2,3-dihydroxypropoxy)propyl]- (9CI) (CA INDEX NAME)




RN 259796-38-8 CAPLUS

CN Dipyrro[3,2-a:2',3'-c]phenazine-11-carboxamide, N-[3-[(bis(4-methoxyphenyl)phenylmethoxy]-2-hydroxypropoxy)propyl]- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 259796-40-2 CAPLUS



CN Butanedioic acid, mono[1-[(bis(4-methoxyphenyl)phenylmethoxy)methyl]-2-[3-[(dipyrro[3,2-a:2',3'-c]phenazine-11-ylcarbonyl)amino]propoxy]ethyl] ester (9CI) (CA INDEX NAME)





PAGE 1-A




RN 259796-43-5 CAPLUS
CN Butanedioic acid, mono[1-[[bis(4-methoxyphenyl)phenylmethoxy)methyl]-13-dipyrro[3,2-a:2',3'-c]phenazin-11-yl-13-oxo-3,6,9-trioxo-12-azatridec-1-yl] ester (9CI) (CA INDEX NAME)





RN 259796-44-6 CAPLUS
CN Butanedioic acid, mono[1-[[bis(4-methoxyphenyl)phenylmethoxy)methyl]-19-dipyrro[3,2-a:2',3'-c]phenazin-11-yl-19-oxo-3,6,9,12,15-pentaoxa-18-azanonadec-1-yl] ester (9CI) (CA INDEX NAME)





IT 259796-24-2P 259796-39-9P 259796-40-2P
259796-41-3P 259796-42-4P 259796-43-5P
259796-44-6P 259796-45-7P 259796-46-8P
259796-47-9P
RL: SPN (Synthetic preparation); PREP (Preparation)
(synthesis and thermal stability of dipyrro[phenazine]-tethered
oligo-DNA and their DNA/RNA duplexes and triplexes)
RN 259796-24-2 CAPLUS
CN Dipyrro[3,2-a:2',3'-c]phenazine-11-carboxamide, N-(2-hydroxyethyl)-N-methyl- (9CI) (CA INDEX NAME)





RN 259796-39-9 CAPLUS
CN Phosphoramidous acid, bis(1-methylethyl)-, 1-[[bis(4-methoxyphenyl)phenylmethoxy)methyl]-2-[3-[[dipyrro[3,2-a:2',3'-c]phenazin-11-yl-13-oxo-3,6,9-trioxo-12-azatridec-1-yl] 2-cyanoethyl ester (9CI) (CA INDEX NAME)





RN 259796-40-2 CAPLUS
CN Butanedioic acid, mono[1-[[bis(4-methoxyphenyl)phenylmethoxy)methyl]-2-[3-[[dipyrro[3,2-a:2',3'-c]phenazin-11-yl-13-oxo-3,6,9-trioxo-12-azatridec-1-yl] 2-cyanoethyl ester (9CI) (CA INDEX NAME)





RN 259796-41-3 CAPLUS
CN Phosphoramidous acid, bis(1-methylethyl)-, 1-[[bis(4-methoxyphenyl)phenylmethoxy)methyl]-13-dipyrro[3,2-a:2',3'-c]phenazin-11-yl-13-oxo-3,6,9-trioxo-12-azatridec-1-yl 2-cyanoethyl ester (9CI) (CA INDEX NAME)





RN 259796-42-4 CAPLUS
CN Phosphoramidous acid, bis(1-methylethyl)-, 1-[[bis(4-methoxyphenyl)phenylmethoxy)methyl]-19-dipyrro[3,2-a:2',3'-c]phenazin-11-yl-19-oxo-3,6,9,12,15-pentaoxa-18-azanonadec-1-yl 2-cyanoethyl ester (9CI) (CA INDEX NAME)



RN 259796-43-5 CAPLUS
CN Butanedioic acid, mono-1-[[bis(4-methoxyphenyl)phenylmethoxy]methyl]-13-dipyrrolyl[3,2,5'-2',3'-c]phenazin-11-yl-13-oxo-3,6,9-trioxo-12-azatridec-1-yl] ester (9CI) (CA INDEX NAME)



RN 259796-44-6 CAPLUS
CN Butanedioic acid, mono[1-[[bis(4-methoxyphenyl)phenylmethoxy)methyl]-19-dipyrido[3,2-a:2',3'-c]phenazin-11-yl-19-oxo-3,6,9,12,15-pentaaza-18-azanadec-1-yl] ester (9CI) (CA INDEX NAME)








```

RN      259796-45-7  CAPLUS
CN      Thymidine, thymidyl-yl-(3'-5')-2'-deoxycytidyl-yl-(3'-5')-2'-
        deoxycytidyl-yl-(3'-5')-2'-deoxyadenylyloxy-[2-{3-[(dipyrro-3,2-
        a:2',3'-c]phenazin-11-ylcarbonyl)amino]propoxymethyl]-1,2-
        ethanediol)oxyphosphinico-(3'-5')-2'-deoxyadenylyl-(3'-5')-
        2'-deoxycytidyl-yl-(3'-5')-2'-deoxyadenylyl-(3'-5')- (9CI)
        (CA INDEX NAME)

```





Absolute stereochemistry.

 HO 






RN 259796-46-8 CAPLUS
 CN Thymidine, thymidyl- (3' → 5')-2'-deoxycytidyl- (3' → 5')-2'-
 deoxycytidyl- (3' → 5')-2'-deoxyadenylyloxy [2- (1,2-dipyrrodo [3,2-
 a:2',3'-c] phenazin-11-yl-10-oxo-2,5,8,11,14-pentaaza-17-azaoctadec-1-yl)-1,2-
 ethanediyl]oxyphosphinico- (3' → 5')-2'-deoxyadenylyl- (3' → 5')-
 2'-deoxycytidyl- (3' → 5')-2'-deoxyadenylyl- (3' → 5')- (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.




RN 259796-47-9 CAPLUS
 CN Thymidine, thymidyl- (3' → 5')-2'-deoxycytidyl- (3' → 5')-2'-
 deoxycytidyl- (3' → 5')-2'-deoxyadenylyloxy [2- (1,2-dipyrrodo [3,2-
 a:2',3'-c] phenazin-11-yl-10-oxo-2,5,8,11,14-pentaaza-17-azaoctadec-1-yl)-1,2-
 ethanediyl]oxyphosphinico- (3' → 5')-2'-deoxyadenylyl- (3' → 5')-
 2'-deoxycytidyl- (3' → 5')-2'-deoxyadenylyl- (3' → 5')- (9CI)
 (CA INDEX NAME)


Absolute stereochemistry.




PAGE 3-A



PAGE 3-B



PAGE 3-C




PAGE 4-A




thioformamide, formamide acetal or thioformamide acetal, in the presence of a halogenating agent. Examples of suitable halogenating agents include but are not limited to thionyl chloride, phosgene, and phosgene derivative. Reactants containing more than one addnl. N-C-C-N group may also be used to prepare comds. with two or more imidazolium groups, by the procedures of the present invention. Certain comds. of the invention prepared from reactants with multiple N-C-C-N groups may have both unreacted N-C-C-N moieties and substituted imidazolium groups. E.g., 1,2-bis(dimethylamino)-2,3-dimethyl-6-(2-pyridinyl)pyrido[1',2':3,4]imidazo[1,5-a]quinoxalin-11-ium perchlorate (90%) was prepared by reaction of 7,8-dimethyl-2,3-di-2-pyridylquinoxaline with Me₂NCSH/BOC12, followed by treatment with aqueous NaClO₄.


IT 246518-08-1P 246518-09-2P 246518-10-5P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of imidazolium cations)
 RN 246518-08-1 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-di-2-pyridinyl-, butyl ester (9CI) (CA INDEX NAME)



RN 246518-09-2 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-di-2-pyridinyl-, octyl ester (9CI) (CA INDEX NAME)




RN 246518-10-5 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-di-2-pyridinyl-, decyl ester (9CI) (CA INDEX NAME)



REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS

REFERENCE COUNT: 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 63 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1999:191213 CAPLUS
 DOCUMENT NUMBER: 132:64244
 TITLE: Synthesis of some sulfonamide derivatives with potential antibacterial activity
 AUTHOR(S): El-Din, Nabaweya Sheraf
 CORPORATE SOURCE: Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Tanta, Tanta, Egypt
 SOURCE: Oriental Journal of Chemistry (1999), 15(2), 223-228
 CODEN: OJCHEG; ISSN: 0970-020X
 PUBLISHER: Oriental Scientific Publishing Co.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Some new quinoxaline-6-sulfonamide and phthalazine-6-sulfonamide derivatives were synthesized. Most of the products showed antibacterial activity.
 IT 112170-26-0P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation and antibacterial activity of)
 RN 112170-26-0 CAPLUS
 CN Glycine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 64 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1999:670141 CAPLUS
 DOCUMENT NUMBER: 131:286517
 TITLE: Imidazolium cations and processes for their preparation
 INVENTOR(S): Donovan, Robert J.; Morgan, Robert J.
 PATENT ASSIGNEE(S): The Rockefeller University, USA
 SOURCE: U.S., 23 pp., Cont.-in-part of U.S. 5,874,587.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5969150	A	19991019	US 1998-124546	19980729
US 5874587	A	19990223	US 1996-673687	19960625
PRIORITY APPL. INFO.:		CASREACT 131:286517; MARPAT 131:286517		
OTHER SOURCE(S):		AB The present invention relates to novel imidazolium comds. and improved processes for the preparation of imidazolium cations with one or more imidazolium moieties optionally substituted with the same or different substituents, which are prepared from a reactant with at least one N-C-C-N group, by reacting with with an N-substituted or N,N-disubstituted		

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 65 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1999:625564 CAPLUS
 DOCUMENT NUMBER: 131:346148
 TITLE: Dimeric analogues of non-cationic tricyclic aromatic carboxamides are a new class of cytotoxic agents
 AUTHOR(S): Spicer, Julie A.; Gamage, Swarna A.; Atwell, Graham J.; Finley, Graeme J.; Baguley, Bruce C.; Denny, William A.
 CORPORATE SOURCE: Auckland Cancer Society Research Centre, Faculty of Medicine and Health Science, The University of Auckland, Auckland, N. Z.
 SOURCE: Anti-Cancer Drug Design (1999), 14(3), 281-289
 CODEN: ACDDRA; ISSN: 0266-9536
 PUBLISHER: Oxford University Press
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB A series of tricyclic aromatic carboxamides, and their corresponding dimeric analogs, were prepared and their growth-inhibitory properties were evaluated in a series of cell lines. The dimeric comds. were prepared by reaction of the appropriate acids with carbonyl-1,1'-dimidazole, isolating the resulting imidazoles, and reacting these with a stoichiometric amount of the diamine. The monomeric carboxamides containing a (CH₂)₂NMe₂ side chain had widely differing inhibitory potencies, with the known nitronaphthalimide (mitonafide) and acridine-4-carboxamide (DACA) being the most potent. The corresponding bis analogs, linked by a (CH₂)₃NMe₂(CH₂)₃ chain, were generally more potent, with the largest increases (dimer/monomer ratio 20- to 30-fold) seen for the nitronaphthalimides and the phenazines. Based on the intrinsic cytotoxicity of the monomers and the highest degree of increase in cytotoxicity on dimerization, the most interesting chromophores appear to be the acridine-4-carboxamide and phenazine-1-carboxamide. Both of these comds. showed significant growth delays (apprx. 6 days) in an in vivo colon 38 tumor model in mice.
 IT 250684-07-2P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USBS (Uses)
 (dimeric analogs of non-cationic tricyclic aromatic carboxamides are a new class of cytotoxic agents in relation to structure)
 RN 250684-07-2 CAPLUS
 CN 2-Phenazinecarboxamide, N,N'-[(methylimino)di-3,1-propanediyl]bis- (9CI) (CA INDEX NAME)




IT 250684-04-9
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USBS (Uses)
 (dimeric analogs of non-cationic tricyclic aromatic carboxamides are a new class of cytotoxic agents in relation to structure)
 RN 250684-04-9 CAPLUS
 CN 2-Phenazinecarboxamide, N-[2-(dimethylamino)ethyl]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 66 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1999:455125 CAPLUS
 DOCUMENT NUMBER: 131:199390
 TITLE: Self-Complementary Cavitands
 AUTHOR(S): Renslo, Adam R.; Rudkevich, Dmitry M.; Rebek, Julius Jr.
 CORPORATE SOURCE: Skaggs Institute for Chemical Biology and The Department of Chemistry, The Scripps Research Institute, La Jolla, CA, 92037, USA
 SOURCE: Journal of the American Chemical Society (1999), 121(32), 7459-7460
 CODEN: JACSAT; ISSN: 0002-7863
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Vase-like cavitands with single adamantyl groups covalently bound on the upper rim via variable-length spacers were prepared and their inclusion processes studied in competing and noncompeting solvents. NMR and computer modeling suggested that in noncompeting p-xylene the adamantane balls were included quant. within the highly shielding environment of the cavitand binding socket in self-complementary dimeric assemblies. In CDCl₃ the assembly is much weaker, and variable temperature binding studies provided $\Delta H = -10.6$ kcal/mol and $\Delta S = -24.5$ eu, i.e., the binding is enthalpically favorable and entropically unfavorable.
 IT 242129-41-5
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (coupling with cavitand diol; preparation of cavitands with covalently bound adamantyl groups on their upper rims and their self-complementary cavity-guest binding in dimeric assemblies)
 RN 242129-41-5 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- (9CI) (CA INDEX NAME)



IT 242144-39-4
 RL: PMU (Formation, unclassified); PEP (Physical, engineering or chemical process); PRP (Properties); FORM (Formation, nonpreparative); PROC (Process)
 (preparation of cavitands with covalently bound adamantyl groups on their upper rims and their self-complementary cavity-guest binding in dimeric assemblies)
 RN 242144-39-4 CAPLUS
 CN 13,23:14,22-Dimetheno-15H,17H,19H,21H-benzo[2',3']benzo[2'',3''] [1,7]benzodioxonino[3''',2''':9'',10''] [1,4]benzodioxonino[6'',5'':9'',10''] [1,4]benzodioxonino[6'',5':9,10] [1,4]benzodioxonino[2,3-b]quinoxaline-27-carboxamide,


2,3,9,10,35,36-hexakis[(1-oxooctyl)amino]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-15,17,19,21-tetraundecyl-, stereoisomer, compd. with stereoisomer of 2,3,9,10,35,36-hexakis[(1-oxooctyl)amino]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-yl-15,17,19,21-tetraundecyl-13,23:14,22-dimetheno-15H,17H,19H,21H-benzo[2',3']benzo[2'',3''] [1,7]benzodioxonino[3''',2''':9'',10''] [1,4]benzodioxonino[6'',5'':9'',10''] [1,4]benzodioxonino[6'',5':9,10] [1,4]benzodioxonino[2,3-b]quinoxaline-27-carboxamide (1:1) (9CI) (CA INDEX NAME)

CM 1
 CRN 242143-99-3
 CNF C158 H227 N9 O15

PAGE 1-A

Me-(CH₂)₆-

PAGE 1-B




PAGE 1-C

CM 2
 CRN 242143-97-1
 CNF C157 H225 N9 O15

PAGE 1-A


-(CH₂)₆-Me

PAGE 2-A




Me-(CH₂)₆-



PAGE 2-B






PAGE 3-B

(CH₂)₁₀-Me




$-(CH_2)_6-Me$

 $(CH_2)_{10}-Me$

IT 242143-99-3P
 RL: PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)
 (self- and heterodimerization; preparation of cavitands with covalently bound adamantyl groups on their upper rims and their self-complementary cavity-guest binding in dimeric assemblies)
 RN 242143-99-3 CAPLUS
 CN 13,23:14,22-Dimetheno-15H,17H,19H,21H-benzo[2',3']benzo[2'',3''] [1,7]benzo dioxonino[3''',2''':9'',10''] [1,4]benzodioxonino[6'',5'':9',10'] [1,4]benzo dioxonino[6',5':9,10] [1,4]benzodioxonino[2,3-b]quinoxaline-27-carboxamide, 2,3,9,10,35,36-hexakis[(1-oxooctyl)amino]-N-(tricyclo[3.3.1.1.3.7]dec-1-ylmethyl)-15,17,19,21-tetraundecyl-, stereoisomer (9CI) (CA INDEX NAME)

 $Me-(CH_2)_6-$

 $-(CH_2)_6-Me$

 $(CH_2)_{10}-Me$

REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L13 ANSWER 67 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1999:447645 CAPLUS
 DOCUMENT NUMBER: 131:310427
 TITLE: Probing the Role of Polyphenol Oxidation in Mediating Insect-Pathogen Interactions. Galloyl-Derived Electrophilic Traps for the Lymantria dispar Nuclear Polyhedrosis Virus Matrix Protein Polyhedrin
 AUTHOR(S): Feldman, Ken S.; Sambandan, Aruna; Bowers, Katherine S.; Appel, Heidi M.
 CORPORATE SOURCE: Department of Chemistry and Pesticide Research Laboratory, The Pennsylvania State University, University Park, PA, 16802, USA
 SOURCE: Journal of Organic Chemistry (1999), 64(16), 5794-5803
 CODEN: JOCEAH; ISBN: 0022-3263
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 131:310427
 GI:



AB Galloyl-derived orthoquinone probes have been designed, synthesized, and utilized in an ongoing study of insect-pathogen interactions. A stable galloyl-derived orthoquinone O-Me ether modified with both acidic and fluorescent appendages (I) was successful in trapping the model nucleophile cysteine, a test protein bearing a single cysteine residue, and the viral occlusion body matrix protein polyhedrin from *Lymantria dispar* nuclear polyhedrosis virus (LdNPV), a pathogen of the gypsy moth caterpillar (GMC). This latter observation may be related to the mol. mechanism by which gallotannins decrease LdNPV infectivity in GMC's. Sufficient site isolation was not achieved with a polymer-bound reactive galloyl hydroxyorthoquinone electrophile to permit similar nucleophile trapping to compete with oligomerization.

IT 246048-12-4P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (trapping cysteine and cysteine-containing proteins by galloyl-derived orthoquinone ether as model for mediating insect-pathogen interactions)
 RN 246048-12-4 CAPLUS
 CN L-Serine, N-[[5-(dimethylamino)-1-naphthalenyl]sulfonyl]-, 4-methoxy-2-phenazinecarboxylate (ester) (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 58 THERE ARE 58 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L13 ANSWER 68 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1999:147950 CAPLUS
 DOCUMENT NUMBER: 130:206994

TITLE: Pyrazine derivatives formed by the reaction of deoxyglucosone with diamino derivatives, antibodies recognizing the product and application in diabetes diagnosis

INVENTOR(S): Uchida, Yoshiaki; Kuran, Yoshihiro; Ito, Satoru
 PATENT ASSIGNER(S): Fujirebio, Inc., Japan
 SOURCE: Ger. Offen., 22 pp.
 CODEN: GMXXBX
 DOCUMENT TYPE: Patent

LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19837664	A1	19990225	DE 1998-19837664	19980819
US 6291198	B1	20010918	US 1998-134388	19980814
JP 11181000	A2	19990706	JP 1998-249122	19980819
JP 3508563	B2	20040322		
JP 3359187	A1	19990324	GB 1998-18343	19980820
PRIORITY APPLN. INFO.:			JP 1997-240348	A 19970821




AB The invention concerns a monoclonal antibody that recognizes pyrazine deriva. (I) that are formed when a diamino derivative (II) is reacting with a 1,2-dicarbonyl derivative, e.g. deoxyglucosone, an in vivo intermediate involved in the nonenzymic glycation of proteins causing diabetic complications. Monoclonal antibodies are raised using I type immunogens and are immobilized after purification. II compds. are labeled at R3, thus when binding to the antibody they can be detected. In I and II the groups are the following: R1 and R2 = H, Me, trihydroxypropyl, dihydroxypropyl, hydroxymethyl; R3 = a spacer group plus a reactive label to form covalent bonds, e.g. carboxyl, hydroxyl, sulphydryl, amino, maleimide, aldehyde, halogen, biotin, etc.; A = pyridine, benzene, furan. Typical 1,2-dicarbonyl compds. that react with II are deoxyglucosone and methylglyoxal. The invention also concerns a test kit that contains the antibody, the carrier to immobilize the antibody and a labeled diamino compound II. Diamino deriva. were synthesized, reacted with deoxyglucosone and coupled to keyhole limpet hemocyanin to immunize mice; monoclonal antibodies were isolated and used in immunoassays. Synthesized diamino deriva. were also labeled with biotin and used as reagents to determine deoxyglucosone using the antibodies immobilized on ELISA plates.


IT 220928-38-1P 220928-39-2P
 RL: ARG (Analytical reagent use); SPN (Synthetic preparation); ANST (Analytical study); PREP (Preparation); USBS (Uses)
 (antigen; pyrazine deriva. formed by the reaction of deoxyglucosone with diamino deriva., antibodies recognizing the product and application in diabetes diagnosis)
 RN 220928-38-1 CAPLUS
 CN 6-Quinoxalinecarboxamide, N-[4-[[2-[[1-[2-[4-[5-[[3aS,4S,6aR]-hexahydro-2-oxo-1H-thieno[3,4-d]imidazol-4-yl]-1-oxopentyl]-1-piperazinyl]ethyl]-2,5-dioxo-3-pyrrolidinyl]thio]ethyl]amino]-4-oxobutyl]-3-[(2S,3R)-2,3,4-trihydroxybutyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




PAGE 1-B




RN 220928-39-2 CAPLUS
 CN 6-Quinoxalinecarboxamide, N-[4-[[2-[[1-[2-[4-[5-[[3aS,4S,6aR]-hexahydro-2-oxo-1H-thieno[3,4-d]imidazol-4-yl]-1-oxopentyl]-1-piperazinyl]ethyl]-2,5-dioxo-3-pyrrolidinyl]thio]ethyl]amino]-4-oxobutyl]-2-[(2S,3R)-2,3,4-trihydroxybutyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




PAGE 1-B




IT 220882-55-3P 220882-67-7P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)
 (immunogen, coupling with keyhole limpet hemocyanin for immunization, reaction with BSA to form antigen for antibody isolation; pyrazine deriva. formed by the reaction of deoxyglucosone with diamino deriva.)
 RN 220882-55-3 CAPLUS
 CN 6-Quinoxalinecarboxamide, N-[4-oxo-4-[[2-(2-pyridinyldithio)ethyl]amino]butyl]-3-[(2S,3R)-2,3,4-trihydroxybutyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 220882-67-7 CAPLUS
 CN 6-Quinoxalinecarboxamide, N-[4-oxo-4-[[2-(2-pyridinyldithio)ethyl]amino]butyl]-2-[(2,3,4-trihydroxybutyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L13 ANSWER 69 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1999:34887 CAPLUS
 DOCUMENT NUMBER: 130:110161
 TITLE: Preparation of substituted N-[(aminomethyl)phenyl]propyl amides as Factor Xa inhibitors
 INVENTOR(S): Klein, Scott I.; Guertin, Kevin R.; Spada, Alfred P.; Pauls, Heinz W.; Gong, Yong; McGarry, Daniel G.
 PATENT ASSIGNEE(S): Rhone-Poulenc Rorer Pharmaceuticals Inc., USA
 SOURCE: PCT Int. Appl., 252 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 5
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9900356	A1	19990107	WO 1998-US13550	19980626
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GR, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, MY, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW			
RM:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GN, GU, HK, HR, HU, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, MY, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW			
US 6080767	A	20000627	US 1997-884405	19970627
CA 2264556	AA	19990107	CA 1998-2264556	19980626
AU 9881771	A1	19990119	AU 1998-61771	19980626
US 741173	B2	20011122		
EP 931060	A1	19990728	EP 1998-931728	19980626
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, SI, FI, RO			
BR 9806060	A	19990831	BR 1998-0606	19980626
JP 2001500532	T2	20010116	JP 1999-505870	19980626
AP 1061	A	20020424	AP 1999-1467	19980626
W:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW			
NO 9900554	A	19990423	NO 1999-854	19990223
NO 314758	B1	20030519		
US 6323227	B1	20011127	US 1999-259528	19990226
PRIORITY APPLN. INFO.:			US 1997-884405	A2 19970627
			US 1996-9485P	P 19960102
			WO 1996-US20770	A2 19961223
			WO 1998-US13550	W 19980626
OTHER SOURCE(S):		MARPAT 130:110161		



REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L13 ANSWER 70 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1998:664773 CAPLUS
 DOCUMENT NUMBER: 130:13971
 TITLE: New synthesis of alloxazine derivatives
 AUTHOR(S): Krasnov, K. A.
 CORPORATE SOURCE: St. Petersburg State Chemical and Pharmaceutical Academy, St. Petersburg, 197376, Russia
 SOURCE: Russian Journal of Organic Chemistry (Translation of Zhurnal Organicheskoi Khimii) (1998), 34(1), 115-119
 CODEN: RJOCQJ; ISSN: 1070-4280
 PUBLISHER: MAIK Nauka/Interperiodica Publishing
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 130:13971

AB Nitrosation of 6-anilino-1,3-dimethyluracile yields 1,3-dimethylalloxazine derivate, and the corresponding 5-oxides. Reduction of the N-oxides results in formation of the alloxazines. This reaction opens the way to difficultly accessible alloxazine derivate.

IT 215865-98-8P 215866-02-7P
 RL: SPN (Synthetic preparation); PREP (Preparation) (new synthesis of alloxazine derivate.)


RN 215865-98-8 CAPLUS

CN Benzo[gl]pteridine-7-carboxylic acid, 1,2,3,4-tetrahydro-1,3-dimethyl-2,4-dioxo-, ethyl ester (9CI) (CA INDEX NAME)




RN 215866-02-7 CAPLUS


CN Benzo[gl]pteridine-7-carboxylic acid, 1,2,3,4-tetrahydro-1,3-dimethyl-2,4-dioxo-, ethyl ester, 5-oxide (9CI) (CA INDEX NAME)



G1



I



II

AB Title compds. I [R = H, OH, NH2; R1 = R2 = H, or R1R2 = NR9; R3 = H, CO2R6, COR6, CON(R6)2, CH2OR7, CH2SR7; R4 = H, alkyl, alkyl-O, thioheterocyclyl, (CH2CH2)nAr, (CH:CH)nAr, CH2Ar; R5 = alk(en/yn)yl, cycloalk(en)yl, heterocycl(en)yl, aryl, heteroaryl, fused systems, etc.; R6 = H, lower alkyl; R7 = H, lower alkyl, aralkyl, lower acyl, aryl, heteroaryl; R8 = H, lower alkyl; R9 = H, R10O2C, R10O, HO, cyano, R10CO, OHC, lower alkyl, OCN, Y1Y2'W, R10 = alkyl, aralkyl, heteroalkyl; Y1', Y2' = H, alkyl, O = R7O, R7S, Y1Y2N; Y1, Y2 = H, alkyl, aryl, aralkyl; or one of Y1 and Y2 = acyl or aryl and the other is as given; Ar = aryl or heteroaryl; n = 0-2] and their pharmaceutically acceptable salts, prodrugs, N-oxides, hydrates, and solvates, are useful as Factor Xa inhibitors. For example, 4-(pyridin-3-yl)benzoic acid was amidated with tert-Bu 3-aminopropionate-HCl via the acid chloride, and the resulting β-acylamino ester underwent a sequence of (1) α-alkylation with 5-iodo-2-[(2-methoxyethoxy)methoxy]benzyl bromide, (2) acidic deprotection of the MEM group, and conversion to the Me ester, (3) Pd-catalyzed cyanation of the iodide, and (4) Pinner reaction and ammonolysis of the nitrile, to give title compound II. Three example compds. showed Ki values of 19.0-94.0 nM in a Factor Xa assay, 46 nM to 1.72 μM in a trypsin assay, and 477 nM to 2.71 μM in a thrombin assay.

IT 215673-02-6P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); B10L (Biological study); PREP (Preparation); USES (Uses) (preparation of substituted [(aminomethyl)phenyl]- or [(aminomethyl)phenyl]propyl amides as Factor Xa inhibitors)

RN 215673-02-6 CAPLUS

CN Benzenepropanoic acid, 3-(aminomethyl)- α-[1-[(6-quinoxalinyloxy)amino]ethyl]-, methyl ester (9CI) (CA INDEX NAME)

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 71 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1998:608600 CAPLUS
 DOCUMENT NUMBER: 129:230740

TITLE: Heteroaryl-hexanoic acid amide derivatives, their preparation and their use as selective inhibitors of MIP-1 α binding to its CCR1 receptor

INVENTOR(S): Brown, Matthew Frank; Kath, John Charles; Poss, Christopher Stanley

PATENT ASSIGNEE(S): Pfizer Inc., USA

SOURCE: PCT Int. Appl., 106 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 8838167	A1	19980903	WO 1998-US1568	19980205
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW			
RM:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, QA, GN, ML, MR, NE, SN, TD, TO			
CA 2282834	AA	19980903	CA 1998-2282834	19980205
CA 2282834	C	20041005		
AU 9861354	A1	19980918	AU 1998-61354	19980205
AU 745687	B2	20020328		
EP 964443	A1	19991229	EP 1998-906013	19980205
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, SI, LT, LV, FI, RO			
TR 9902056	T2	20000121	TR 1999-9902056	19980205
BR 9807858	A	20000222	BR 1998-7858	19980205
JP 2000513740	T2	20001017	JP 1998-537644	19980205
JP 3771591	B2	20060426		
IL 131163	A1	20050619	IL 1998-131163	19980205
ZA 9801602	A	19990921	ZA 1998-1602	19980226
AP 1056	A	20020405	AP 1998-1200	19980226
W:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, QA, GN, ML, MR, NE, SN, TD, TO			
BQ 103688	A	20001130	BQ 1999-103688	19990824
NO 9904101	A	19990825	NO 1999-4101	19990825
NO 313877	B1	20021216		
US 6403587	B1	20020611	US 2000-380269	20000518
US 2002198207	A1	20021226	US 2002-198207	20030522
PRIORITY APPLN. INFO.:			US 1997-391699	P 19970226
			WO 1998-US1568	W 19980205
			US 2000-380269	A3 20000518
OTHER SOURCE(S):		MARPAT 129:230740		

G1




I

AB I [R1 = optionally substituted (C2-C9)heteroaryl; R2 = optionally substituted phenyl-(CH2)m-, naphthyl-(CH2)m-, (C3-C10)cycloalkyl-(CH2)m-, (C1-C6)alkyl or (C2-C9)heteroaryl-(CH2)m-; m = integer from zero to four; R3 = H, optionally substituted (C1-C10)alkyl, (C3-C10)cycloalkyl-(CH2)n-, (C2-C9)heteroaryl-(CH2)n-, (C2-C9)heteroaryl-(CH2)n-, aryl-(CH2)n-, n = integer from zero to six; R4 and the carbon to which it is attached form an optionally substituted and/or fused five to seven membered carbocyclic ring; R5 = H, (C1-C6)alkyl, hydroxy, (C1-C6)alkoxy, hydroxy-(C1-C6)alkyl, (C1-C6)alkoxyCO, (C3-C10)cycloalkyl-(CH2)p-, optionally substituted (C2-C9)heterocycloalkyl-(CH2)p-, (C2-C9)heteroaryl-(CH2)p-, phenyl-(CH2)p- or naphthyl-(CH2)p-, p = integer from zero to four; R4 and R5 together with the nitrogen atom to which they are attached form an optionally substituted (C2-C9)heterocycloalkyl group; R6 = H, (C1-C6)alkyl, amino] were prepared. The present compounds are potent and selective inhibitors of MIP-1 α binding to its receptor CCR1, and are thus useful to treat inflammation and other immune disorders. E.g., quinoxaline-2-carboxylic acid 1[(S)-benzyl-4(R)-benzylcarbamoyl-7-fluoro-2(S)-hydroxy-7-methyloctyl]amide was prepared

IT 212789-54-3P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of heteroaryl-substituted hexanamide and their use as selective inhibitors of MIP-1 α binding to its CCR1 receptor)

RN 212789-54-3 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[[(1S,2S,4R)-4-(aminocarbonyl)-7-fluoro-2-hydroxy-7-methyl-1-phenylmethyloctyl]- (9CI) (CA INDEX NAME)]

Absolute stereochemistry.



REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 72 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1998:402403 CAPLUS
DOCUMENT NUMBER: 129:81964
TITLE: Preparation and use of ketobenzamides as calpain inhibitors
INVENTOR(S): Lubisch, Wilfried; Moller, Achim; Treiber, Hans-Jorg
PATENT ASSIGNEE(S): BASF A.-G., Germany; Lubisch, Wilfried; Moller, Achim; Treiber, Hans-Jorg
SOURCE: PCT Int. Appl., 84 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9825883	A1	19980618	WO 1997-EP6655	19971128
W: AL, AU, BG, BR, BY, CA, CN, CZ, GE, HU, ID, IL, JP, KR, KZ, LT, LV, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, UA, US, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				


RM: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE
CA 9274464 AA 19980618 CA 1997-2274464 19971128
AU 927523 A1 19980703 AU 1998-57523 19971128
AU 721620 B2 20000713
EP 944582 A1 19980929
EP 944582 B1 20003702 EP 1997-953714 19971128
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, SI, FI, RO
CN 1245486 A 20000223 CN 1997-181748 19971128
NZ 335981 A 20000401 NZ 1997-335981 19971128
BR 9713704 A 20000509 BR 1997-13704 19971128
JP 2001506614 T2 20010522 JP 1998-526186 19971128
RU 1905999 C2 20021010 RU 1999-115765 19971128
SK 282680 B6 20021106 SK 1999-745 19971128
AT 244216 E 20030715 AT 1997-953714 19971128
SE 2202663 T3 20040401 SE 1997-953714 19971128
HR 970680 B1 20020831 HR 1997-970680 19971210
ZA 9711141 A 19990611 ZA 1997-11141 19971211
TW 536530 B 20030611 TW 1997-8618865 19971211
US 6103720 A 20000815 US 1999-319511 19990608
NO 9902821 A 19990611 NO 1999-2821 19990610
KR 2000057495 A 20000915 KR 1999-705172 19990610
BG 63182 B1 20011231 BG 1999-103485 19990611
PRIORITY APPLN. INFO.: DE 1996-19651316 A 19961211
NO 1997-EP6655 W 19971128

OTHER SOURCE(S): MARPAT 129:81964
AB The invention concerns ketobenzamides of formula R1X(R2)n-C6H3-CONHCH(R3)COOCH(R4) where R1 = Ph, naphthyl, (substituted)heterocycle; R2 = Cl, Br, F, NO2, NH2, NHR5, CO2H, (substituted)alkyl, -alkenyl, -alkynyl, R5 = CO-alkyl, COPh, CO-C10H7, SO2-alkyl, CO-alkoxy, ureido, alkoxy; R3 = (substituted)alkyl; X = (substituted)(functionalized)chain from 0-10 atoms, or R2-substituted-C6H3; R4 = OH, (substituted)alkoxy, (substituted)NH2, heterocyclic ring, useful as calpain inhibitors. The invention further concerns their preparation. The novel compounds are suitable for combating diseases. Thus, 3(S)-1-amino-2-hydroxy-4-phenylbutyric acid Me ester was condensed with 2-phenylbenzoic acid to give (S)-I [R1 = Ph; X = null; n = 0; R3 = CH2Ph; R4 = OMe(II)]. In vitro calpain-inhibition tests, II had KI of <10 μ M.

IT 209174-24-3P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(preparation and use of ketobenzamides as calpain inhibitors)

RN 209174-24-3 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[4-[[[(1S)-3-amino-2,3-dioxo-1-(phenylmethyl)propyl]amino]carbonyl]phenyl]methyl]- (9CI) (CA INDEX NAME)]


Absolute stereochemistry.



IT 209174-23-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and use of ketobenzamides as calpain inhibitors)

RN 209174-23-2 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[[4-[[[(1S)-3-amino-2-hydroxy-3-oxo-1-(phenylmethyl)propyl]amino]carbonyl]phenyl]methyl]- (9CI) (CA INDEX NAME)]

Absolute stereochemistry.




REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 73 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1998:402013 CAPLUS
DOCUMENT NUMBER: 129:146424
TITLE: Reactive dyes-human serum albumin (HSA) binding by frontal analysis affinity chromatography
AUTHOR(S): Alberghina, Gaetano; Babulano, Giuliano A.; Fieschella, Salvatore; Renda, Emanuela; Bianchini, Roberto; Forte, Claudia
CORPORATE SOURCE: Dipartimento di Scienze Chimiche, Universita di Catania, Catania, I-95125, Italy
SOURCE: Gazzetta Chimica Italiana (1997), 127(12), 803-808
CODEN: GCIT9A; ISSN: 0016-5603
Societa Chimica Italiana
PUBLISHER: Journal
DOCUMENT TYPE: English
LANGUAGE: English

AB The binding with human serum albumin (HSA) of Cibacron Blue F 3GA, Levafix Brilliant Blue SB, the structure of which has been carefully investigated, Reactive Orange 046, and Reactive Red 022 dyes, immobilized on sepharose CL 6B, has been quant. investigated by frontal affinity chromatog. This technique allowed the calcn. of the association const. of the ligand-HSA complexes, on the basis of the dependence of the HSA elution volume on its initial concentration. The values obtained were high and comparable for Cibacron and Levafix dyes (5.4 and 5.7-106 M-1, resp.) and significantly smaller for Reactive Orange 046 and Reactive Red 022 (0.23 and 0.4-106 M-1). A possible interpretation of these data, based on the different structures of the bonded dyes, is proposed.

IT 206058-73-3
RL: ARG (Analytical reagent use); PEP (Physical, engineering or chemical process); ANST (Analytical study); PROC (Process); USES (Uses)
(Levafix Brilliant Blue S-8; reactive dyes-human serum albumin binding by frontal anal. affinity chromatog.)

RN 206058-73-3 CAPLUS
CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxalyl)carbonyl]amino]methyl]-3-sulfophenyl]methyl]amino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)]



PAGE 1-A

PAGE 2-A

REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 74 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1998:26491 CAPLUS
DOCUMENT NUMBER: 128:308499
TITLE: Bis(acridinecarboxamide) and bis(phenazinecarboxamide) as antitumor agents
INVENTOR(S): Denny, William Alexander; Gamage, Swarnalatha Akuritaye; Spicer, Julie Ann; Baguley, Bruce Charles; Finley, Graeme John
PATENT ASSIGNEE(S): Xenova Ltd., UK; Denny, William Alexander; Gamage, Swarnalatha Akuritaye; Spicer, Julie Ann; Baguley, Bruce Charles; Finley, Graeme John
SOURCE: PCT Int. Appl., 100 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9817650	A1	19980430	WO 1997-GB2886	19971017
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE,				

DK, SE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW

RM: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CP, CO, CI, CM, GA, GN, ML, MR, NE, SN, TD, TO


CA 2268411 AA 19980430 CA 1997-2268411 19971017
 AU 9747137 A1 19980515 AU 1997-47137 19971017
 AU 971724 B2 20000330 19971017
 ZA 9709331 A 19980521 ZA 1997-9331 19971017
 ZA 9709328 A 19980706 ZA 1997-9328 19971017
 EP 934278 A1 19990811 EP 1997-909456 19971017
 EP 934278 B1 20020904 19971017

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI

GB 2334032 A1 19990811 GB 1999-8192 19971017
 GB 2334032 B2 20001108 19971017
 BR 9711948 A 19990824 BR 1997-11948 19971017
 CN 1240430 A 20000105 CN 1997-180614 19971017
 CN 116285 B 20030730 19971017
 NZ 335055 A 20000929 NZ 1997-335055 19971017
 JP 2001503399 T2 20010313 JP 1998-519109 19971017
 RU 2179972 C2 20020227 RU 1999-109978 19971017
 AT 223381 E 20000915 AT 1997-909456 19971017
 PT 934278 T 20030311 PT 1997-909456 19971017
 ES 2183142 T3 20030316 ES 1997-909456 19971017
 CZ 295302 B6 20050713 CZ 1999-1271 19971017
 TW 432060 B 20010501 TW 1997-86115404 19971018
 BG 103329 A 20001130 BG 1999-103329 19990413
 BG 64555 B1 20050729 19990413
 NO 9901833 A 19990603 NO 1999-1833 19990416
 NO 313381 B1 20020923 19990416
 KR 2000049252 A 20000725 KR 1999-703357 19990416
 US 6114332 A 20000905 US 1999-284623 19990618
 HK 1018773 A1 20010302 HK 1999-103666 19990826
 A 1996-21795 A 19961018
 W 19971017

PRIORITY APPLN. INFO.: CASREACT 128:308499; MARPAT 128:308499


OTHER SOURCE(S):
 GI



AB Compds. I [R1-R4 = H, C1-4 alkyl, OH, etc.; or R1 and R2 together form a methylenedioxy group; R5, R6 = H, C1-4 alkyl; X = CH, N; Z = (CH2)n, (CH2)no(CH2)n, (CH2)nNR7(CH2)n, (CH2)nNR7(CH2)nNR7(CH2)n, (CH2)nN(CH2)n, (CH2)n, R7 = H, C1-4 alkyl; n = 1-4; with the exception of compds. wherein each X is N, each of R1-R6 is H, the carboxamide moiety is attached to position 1 of each phenazine ring and Z is (CH2)2NH(CH2)2, (CH2)3NH(CH2)3, (CH2)3N(CH2CH2)2N(CH2)3, (CH2)2NH(CH2)2NH(CH2)2 or (CH2)3NH(CH2)2NH(CH2)3] or a pharmaceutically acceptable acid addition salt or N-oxide thereof; have activity as an antitumor and antibacterial agent. Thus, bis[[(5-methylacridine-4-carboxamido)propyl]methylamine] was prepared and showed an IC50 value of 11 nM on a wild-type human leukemia line (Jurkat; JLC).

IT 206531-48-8P
 RI: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USSS (Uses)
 Bis[acridinecarboxamide] and bis[phenazinecarboxamide] as antitumor and antibacterial agents

RM 206531-48-8 CAPLUS
 CN 2-Phenazinecarboxamide, N,N'-[(methylimino)di-3,1-propanediyl]bis-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 75 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1998:266807 CAPLUS
 DOCUMENT NUMBER: 128:295818

TITLE: An improved computational approach to the determination of thermodynamic and spectral complexation parameters from overlapping bands. Applications to the case of the Cibacron dimer and to the multiple HSA-Levafix association

AUTHOR(S): Ambrosetti, Roberto; Ricci, Domenico; Bianchini, Roberto

CORPORATE SOURCE: CNR, Istituto di Chimica Quantistica ed Energetica Molecolare, Pisa, I-56126, Italy

SOURCE: Gazzetta Chimica Italiana (1997), 127(10), 567-575
 CODEN: GCITAP; ISSN: 0016-5603

PUBLISHER: Societa Chimica Italiana
 DOCUMENT TYPE: Journal
 LANGUAGE: English


AB An algorithm for the simultaneous evaluation of the thermodyn. parameters related to multiple equilibrium and of the spectra of dye complexes is described. The algorithm can accept as input data any stoichiometry for complex species and relies on the simultaneous fitting of large sets of data obtained at different concns., temps., and wavelengths. Data from different measuring techniques, such as UV-visible absorption or CD, may be included in a single fit. Details on an easily modifiable, yet computationally efficient implementation of the algorithm on a standard PC are given. Results are presented for the dimer aggregation of the dye Cibacron Blue F3GA and for the association of human serum albumin with a Levafix reactive anthraquinone dye.

IT 206058-73-3D, complexes with human serum albumin
 RI: PRP (Properties)
 (algorithm for determination of spectral and thermodyn. parameters of)

RM 206058-73-3 CAPLUS
 CN 2-Anthracesulfonic acid, 1-amino-4-[[[4-[[[2,3-dichloro-6-quinolalyl]carbonyl]amino]methyl]-3-sulfonyl]methyl]amino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)

AB 2-Anthracesulfonic acid, 1-amino-4-[[[4-[[[2,3-dichloro-6-quinolalyl]carbonyl]amino]methyl]-3-sulfonyl]methyl]amino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A

● 2 Na

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 76 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1998:210752 CAPLUS
 DOCUMENT NUMBER: 128:257445

TITLE: Preparation of indolylbenzoquinoloxalinones and related compounds as protein kinase C inhibitors.

INVENTOR(S): Bergstrand, Hakan; Karabelas, Koetas; Sjö, Peter

PATENT ASSIGNEE(S): Astra Aktiebolag (Publ), Swed.

SOURCE: PCT Int. Appl., 63 pp.
 CODEN: PIXX02

DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9813368	A1	19980402	WO 1997-581582	19970919

W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RM: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CP, CO, CI, CM, GA, GN, ML, MR, NE, SN, TD, TO


TW 472045 B 20020111 TW 1997-86113549 19970918
 ZA 9708469 A 19980325 ZA 1997-8469 19970919
 CA 2265854 A 19980402 CA 1997-2265854 19970919
 AU 9744775 A1 19980417 AU 1997-44775 19970919
 AU 716279 B2 20000224 19970919
 EP 929551 A1 19990721 EP 1997-943259 19970919

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO

NZ 334531 A 20000929 NZ 1997-334531 19970919
 US 6271231 B1 20010807 US 1997-981266 19971218
 US 2001025043 A1 20010927 US 2001-865231 20010525
 SE 1996-3505 A 19960925
 SE 1997-2747 A 19970718
 WO 1997-581582 W 19970919
 US 1997-981266 A1 19971218

PRIORITY APPLN. INFO.: MARPAT 128:257445

OTHER SOURCE(S):
 GI




AB Title compds. [I; A, X, Y, Z = C, N; ≥2 of A, X, Y, Z = C; may be substituted and/or annulated; excluding 3-(1H-indol-3-yl)-1H-quinoloxalin-2-one, 3-(2-methyl-1H-indol-3-yl)-1H-quinoloxalin-2-one, and 3-(1,2-diphenyl-1H-indol-3-yl)-1H-quinoloxalin-2-one], were prepared as protein kinase C inhibitors (no data). Thus, 1,3-phenylenediamine was stirred overnight with [1-(3-(1,3-dioxaisoindol-2-yl)propyl)-1H-indol-3-yl]oxoacetic acid 2,5-dioxopyrrolidin-1-yl ester (preparation given) in THF to give 3-[3-(3-oxo-3,4-dihydroquinoloxalin-2-yl)indol-1-yl]propylammonium acetate. The latter was stirred with MeOH in THF/H2O to give 3-[3-(3-oxo-3,4-dihydroquinoloxalin-2-yl)indol-1-yl]propylammonium acetate.

IT 205376-68-7P
 RI: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USSS (Uses)
 (preparation of indolylbenzoquinoloxalinones and related compds. as protein kinase C inhibitors)

RM 205376-68-7 CAPLUS
 CN 6-Quinoloxalinecarboxylic acid, 2-[1-(3-aminopropyl)-1H-indol-3-yl]-3,4-dihydro-3-oxo-, ethyl ester, monoacetate (9CI) (CA INDEX NAME)

CM 1
 CRN 205376-67-6
 CMF C22 H22 N4 O3



CM 2

CRN 64-19-7
CMP C2 H4 O2



REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 77 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 1999:126254 CAPLUS

DOCUMENT NUMBER: 126 204878

TITLE: Preparation of pyrazinobenzothiazine derivatives and analogs for the treatment of inflammation and autoimmune diseases

INVENTOR(S): Kaneko, Toshihiko; Clark, Richard; Ohi, Norihito; Ozaki, Fumihiro; Kawahara, Tetsuya; Kamada, Atsushi; Okano, Kazuo; Yokohama, Hiromitsu; Muramoto, Kenzo; Arai, Tohru; Onkuro, Masayoshi; Takenaka, Osamu; Sonoda, Jiro

PATENT ASSIGNER(S): Eisai Co., Ltd., Japan
SOURCE: PCT Int. Appl., 1344 pp.
CODEN: PIXX22

DOCUMENT TYPE: Patent

LANGUAGE: Japanese


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9806720	A1	19980219	WO 1997-02787	19970808
W: AU, CA, CN, HU, JP, KR, MX, NO, NZ, RU, US				
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
CA 2262569	AA	19980219	CA 1997-262569	19970808
AU 9737849	A1	19980306	AU 1997-37849	19970808
ZA 9707103	A	19980209	ZA 1997-7103	19970808
EP 934941	A1	19990811	EP 1997-934750	19970808
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI				
US 6518423	B1	20030211	US 1999-230852	19990405
US 2004092737	A1	20040513	US 2002-247310	20020920
PRIORITY APPL. INFO.:			JP 1996-210344	A 19960809
			WO 1997-02787	W 19970808
			US 1999-230852	A3 19990405

OTHER SOURCE(S): MARPAT 126 204878

GI



AB The title compds. I (R1 to R3 are the same or different and each represents hydrogen, optionally substituted lower alkyl, optionally substituted cycloalkyl, etc., provided that when R1 to R3 are all optionally substituted lower alkyl groups, they do not simultaneously represent Me groups; R represents hydrogen, lower alkyl, etc.; E represents N, C, etc.; Z represents O, S, SO, SO2, etc.; and the ring G represents an optionally substituted heteroaryl ring having at least one nitrogen atom) are prepared. I are useful in the treatment and prevention of inflammatory immunol. diseases, autoimmune diseases, rheumatism, collagen disease, asthma, nephritis, ischemic reflow disorders, psoriasis, atopic dermatitis or rejection reactions following organ transplantation. The compound (syn)-[3-(10H-pyrazino[2,3-b][1,4]benzothiazin-8-ylmethyl)-3-azabicyclo[3.3.1]nona-9-yl]acetic acid (II) at 10 mg/kg orally gave 65% inhibition of carrageenin-induced inflammation in rats. II in vitro showed IC50 of 2.3 μM against the expression of ICAM-1.


IT 203651-75-6P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(Preparation of pyrazinobenzothiazine derivs. and analogs for treatment of inflammation and autoimmune diseases)

RN 203651-75-6 CAPLUS

CN Pyrazino[2,3-b]quinoxaline-7-carboxylic acid, 1,4-dihydro-, ethyl ester (9CI) (CA INDEX NAME)



REFERENCE COUNT: 46 THERE ARE 46 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 78 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 1997:679093 CAPLUS

DOCUMENT NUMBER: 127:318971

TITLE: Preparation of benzoxazines for enhancing synaptic response

INVENTOR(S): Rogers, Gary A.; Lynch, Gary S.
PATENT ASSIGNER(S): Regents of the University of California, USA; Cortex Pharmaceuticals, Inc

SOURCE: PCT Int. Appl., 34 pp.
CODEN: PIXX22

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9736907	A1	19971009	WO 1997-US5184	19970328
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GR, HU, IL, IS, JP, KR, KZ, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, AM, AZ, BY, KZ, MD, RU, TJ, TM, TR, UA, UG, UZ, VN, YU, RW: GH, KE, LG, MG, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BP, BJ, CF, CO, CI, CM, GA, GN, ML, MR, NE, SN, TD, TO				
US 5736543	A	19980407	US 1996-624335	19960403
CA 2249654	AA	19971009	CA 1997-249654	19970328
AU 9724274	A1	19971022	AU 1997-24274	19970328
AU 708212	B2	19990729		
EP 891365	A1	19990120	EP 1997-919965	19970328
EP 891365	B1	20021016		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2000053664	T2	20000328	JP 1997-535488	19970328
JP 3170294	B2	20010528		
AT 226206	E	20021115	AT 1997-919965	19970328
ES 2185932	T3	20030501	ES 1997-919965	19970328
US 5562447	A	19991005	US 1998-19883	19980206
AU 9944543	A1	19991021	AU 1999-44543	19990817
AU 721936	B2	20000720		

PRIORITY APPL. INFO.:

OTHER SOURCE(S): MARPAT 127:318971

GI



AB Compds. of formula [I; R1, R2 = independently selected from the group consisting of H and R4O such that at least one of R1 and R2 is R4O; R4 = a member selected from the group consisting of H, alkyl and halo-substituted alkyl; or R1 and R2 together form a single divalent moiety selected from the group consisting of O-R5-O, S-R5-S, O-R5-S, N-CR6CR7N, O-CR8N, N-CR8O; wherein R5 = a member selected from the group consisting of C(R9)2, C(R9)2C(R9)2, CR9CR9; R9 = H, halo, C1-6 alkyl, C1-6 haloalkyl; R3, R6, R7, R8 = a member selected from the group consisting of H, C1-6 alkyl, and C1-6 haloalkyl] are disclosed for use in enhancing synaptic responses mediated by α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic (AMPA) receptors. The compds. are effective in the treatment of subjects suffering from impaired nervous or intellectual functioning due to deficiencies in the number or strength of excitatory synapses or in the number of AMPA receptors. The compds. can also be used for the treatment of non-impaired subjects for enhancing performance in sensory-motor and cognitive tasks which depend on brain networks utilizing AMPA receptors, for improving the performance of subjects with memory deficiencies, for treating schizophrenia, and for improving memory encoding. Thus, 3,4-methylenedioxysuccinic acid was condensed with 4-aminobutyraldehyde di-ethyl acetal using N,N-carbonyldiimidazole in CH2Cl2 at room temperature overnight to quant. give N-(4,4-diethoxybutyl)-3,4-methylenedioxysuccinamide, which was dissolved in CHCl3 and allowed to stand in the presence of camphorsulfonic acid overnight to give I (R1R2 = OCH2O, R3 = H, n = 2). The latter compound and I (R1R2 = OCH2CH2O, R3 = H, n = 2).

n = 2) in vivo showed the threshold dose of 1 and 0.1 mg/kg, resp., for enhancing memory in rats in a learning paradigm that depends on a performance in an 8-arm radial maze described by Staubli et al. (PNAS, 1994), and at 0.1 and >0.03 mM in vitro, resp. increased the amplitude of the field excitatory post-synaptic potential (EPSP) to a value 25% above the base line in slices of rat hippocampus.


IT 197584-96-6P 197584-97-7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(Preparation of benzoxazines with affinity to AMPA receptors for enhancing synaptic response)


RN 197584-96-6 CAPLUS

CN 6-Quinoxalinecarboxamide, N-(4,4-diethoxybutyl)-7-hydroxy- (9CI) (CA INDEX NAME)



RN 197584-97-7 CAPLUS

CN 6-Quinoxalinecarboxamide, N-(5,5-diethoxypentyl)-7-hydroxy- (9CI) (CA INDEX NAME)



L13 ANSWER 79 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 1996:570703 CAPLUS

DOCUMENT NUMBER: 125:259566

TITLE: Electrochemical study of C60-containing diimine and polyazine ligands: towards a fullerene-based photoactive molecular device

AUTHOR(S): Paolucci, Francesco; Marrocchio, Massimo; Ceroni, Paola; Paradisi, Carmen; Roffie, Sergio; Prato, Maurizio

CORPORATE SOURCE: Dep. Chem., Univ. Bologna, Bologna, 40126, Italy

SOURCE: Proceedings - Electrochemical Society (1996), 96-10(Recent Advances in the Chemistry and Physics of Fullerenes and Related Materials, Vol. 3), 157-164

CODEN: PESDOD; ISSN: 0161-6374

PUBLISHER: Electrochemical Society

DOCUMENT TYPE: Journal


LANGUAGE: English

AB The complete electrochem. characterization of the fulleropyrrolidines containing the 2,2'-bipyridine (bpy) and the 2,3-bis(2-pyridyl)quinoxaline (dpq) fragment, resp., is reported. These ligands represent important building blocks for the assembling of supramol. (polynuclear) metal complexes in which a photoinduced intramol. charge separation may in principle take place. The comparison of the cyclic voltammograms of these species with those of suitable model moies. has allowed the localization of the eight reversible reduction processes observed for both species, a fundamental prerequisite for the assignment of the redox sites in the mono- and polynuclear complexes.


IT 182219-47-2

RL: PRP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); PROC (Process); RACT (Reactant or reagent)
 (electrochem. reduction in THF containing tetrabutylammonium hexafluorophosphate: towards fullerene-based photoactive mol. device)
 RN 182219-47-2 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-di-2-pyridinyl-, 13-[4-(1',5'-dihydro-1'-methyl-2'H-[5,6]fullereno-C60-1h-[1,9-c]pyrrol-2-yl)phenyl]-13-oxo-3,6,9,12-tetraoxatridec-1-yl ester (9CI) (CA INDEX NAME)


PAGE 1-A



PAGE 1-B



PAGE 2-A




PAGE 3-A




IT 182219-47-2D, transition metal complexes
 RL: PRP (Physical, engineering or chemical process); PRP (Properties); PROC (Process)
 (photoinduced intramol. charge separation in)
 RN 182219-47-2 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-di-2-pyridinyl-, 13-[4-(1',5'-dihydro-1'-methyl-2'H-[5,6]fullereno-C60-1h-[1,9-c]pyrrol-2-yl)phenyl]-13-oxo-3,6,9,12-tetraoxatridec-1-yl ester (9CI) (CA INDEX NAME)


PAGE 1-A



PAGE 1-B



PAGE 2-A




PAGE 3-A



L13 ANSWER 80 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1996:542076 CAPLUS
 DOCUMENT NUMBER: 125:204309
 TITLE: Electrochemical and ESR spectroscopic study of 2,7-disubstituted phenazines
 AUTHOR(S): Michida, Takeshi; Sayo, Hiroteru
 CORPORATE SOURCE: Fac. Pharmaceutical Sci., Kobe-Gakuin Univ., Kobe, 651-21, Japan
 SOURCE: Chemical & Pharmaceutical Bulletin (1996), 44(8), 1448-1453
 CODEN: CPBUTL; ISSN: 0009-2363
 PUBLISHER: Pharmaceutical Society of Japan
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Cyclic voltammetry (CV) for various 2,7-disubstituted phenazines (1 mM) was carried out in MeCN containing CF3CO2H (1% and 2%) and NaClO4 (0.1M) as a supporting electrolyte under N2. Phenazines showed 2 cathodic peaks (Epc1 and Epc2) and these peaks had counterparts (Epa1 and Epa2, resp.). Plots of the peak potentials against op were linear. The first cathodic wave corresponds to the reduction of singly protonated phenazines followed by proton transfer. The second cathodic wave corresponds to the reduction of the cation radical of dihydrophenazines to produce dihydrophenazine as a final product. ESR spectroscopy of these compds. in MeCN and in MeCN containing 1% CF3CO2H was conducted and computer simulation of the spectra was carried

out. Splitting due to halogen or o-alkyl substituents was observed MO calcn. of anion radicals generated from the phenazines and cation radicals generated from doubly protonated phenazines did not give good agreement with the results of ESR spectrometry.


IT 72848-45-4
RL: PREP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(electrochem. and ESR spectroscopic study of disubstituted phenazines)
RN 72848-45-4 CAPLUS
CN 2,7-Phenazinedicarboxylic acid, diethyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 81 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STM
ACCESSION NUMBER: 1996:170757 CAPLUS
DOCUMENT NUMBER: 124:202310
TITLE: Preparation of 1,2,3,4-tetrahydro-2,3-dioxoquinoxaline-6-sulfonamides as AMPA and kainate receptor antagonists
INVENTOR(S): Rivo, Rndre; Vizi, E. Szilveszter; Makara, Gabor; Reiter, Jozsef; Blasko, Gabor; Simig, Gyula; Gaal, László; Fekete, Marton
PATENT ASSIGNEE(S): Egis Gyógyszergyár Rt., Hung.
SOURCE: PCT Int. Appl., 24 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9531443	A1	19951123	WO 1995-HU15	19950518
W: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GS, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, ME, MG, MN, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TT, UA				
RW: KE, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BW, CP, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
HU 71334	A2	19951128	HU 1994-1522	19940518
HU 217837	B	20000428		
CA 2190532	AA	19951123	CA 1995-2190532	19950518
AU 9524162	A1	19951205	AU 1995-24162	19950518
EP 759910	A1	19970305	EP 1995-918107	19950518
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, NL, SE				
JP 10504019	T2	19980414	JP 1995-529476	19950518
US 5912245	A	19990615	US 1997-737273	19970214
PRIORITY APPL. INFO.:				
OTHER SOURCE(S):				
GI				

IT 170467-25-1P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)
RN 170467-25-1 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2-[(4-(butoxycarbonyl)phenyl)amino]-3-(1-methyl-2-nitroethenyl)-, butyl ester (9CI) (CA INDEX NAME)




L13 ANSWER 83 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STM
ACCESSION NUMBER: 1995:716813 CAPLUS
DOCUMENT NUMBER: 123:112079
TITLE: Preparation of quinoxaline-2-carboxamides as antidiabetics
INVENTOR(S): Komatsu, Makoto; Sato, Hideaki; Taira, Shinichi; Miyake, Masahiro; Magata, Kiyohiko; Yoshida, Hidehiro; Ueyama, Atsunori; Nishi, Takao
PATENT ASSIGNEE(S): Otsuka Pharmaceutical Co. Ltd., Japan
SOURCE: PCT Int. Appl., 507
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9509159	A1	19950406	WO 1994-JP1559	19940922
W: AU, CA, CH, KR, US				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
CA 2150345	AA	19950406	CA 1994-2150345	19940922
AU 9476660	A1	19950418	AU 1994-76660	19940922
AU 674613	B2	19970102		
EP 670831	A1	19950913	EP 1994-927085	19940922
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
CN 1114634	A	19960110	CN 1994-190719	19940922
JP 08012579	A2	19960116	JP 1994-259309	19940928
JP 2759257	B2	19980528		
PRIORITY APPL. INFO.:				
OTHER SOURCE(S):				
GI				




AB Title compds. [I; R = NR3R4; R1 = H, halo, alkyl, alkoxy, etc.; R2 = H,




AB Title compds. [I; R1 = H or NO2; R2, R3 = H, (un)substituted alk(en)yl; NR2R3 = heterocyclyl] were prepared. Thus, I (R1 = NO2, NR2R3 = piperidino) had Ki of 6.3x10-7 and 2.0x10-6M for inhibition of AMPA and kainate binding at rat brain membrane preparation in vitro.

IT 174526-64-8P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of 1,2,3,4-tetrahydro-2,3-dioxoquinoxaline-6-sulfonamides as AMPA and kainate receptor antagonists)
RN 174526-64-8 CAPLUS
CN Glycine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinolalyl)sulfonyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)




L13 ANSWER 82 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STM
ACCESSION NUMBER: 1995:753036 CAPLUS
DOCUMENT NUMBER: 123:339996
TITLE: Synthesis of quinoxaline derivatives from polynitro-3-thiolene 1,1-dioxide
Khlytin, A. L.; Efremova, I. E.; Berestovitskaya, V. M.
CORPORATE SOURCE: Ross. Gos. Pedagog. Univ., Russia
SOURCE: Zhurnal Organicheskoi Khimii (1994), 30(9), 1434-5
CODEN: ZORKAS; ISSN: 0514-7492
PUBLISHER: Nauka
DOCUMENT TYPE: Journal
LANGUAGE: Russian
GI




AB Polynitro-3-thiolene 1,1-dioxide I reacted with para-substituted anilines to give quinoxalines (II; R = Me, Cl, Br, COOBu).

(halo)alkyl, alkoxy, etc.; R3, R4 = H, alkyl, alkanoyl, alkoxy, carbonyl, substituted CH2Ph, heterocyclylalk(en)yl, etc.; m = 0 or 1; n = 0; r = 1 or 2) were prepared. Thus, benzofuroxan was cyclocondensed with MeCOCH2CO2Et and the product converted in 2 steps to I (R2 = Me, m = 1, n = r = 0) (II; R = OSu) which was amidated by 3-aminomethylbenzofuran to give II (3-benzofurylaminoethyl). II (R = NHCH2CH2CH2Me, R5 = 3-benzofuryl) gave 2-deoxyglucose uptake of rat striated muscle L6 cells 249% of controls at 10-6mol (sic).


IT 165735-35-3P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of quinoxaline-2-carboxamides as antidiabetics)
RN 165735-35-3 CAPLUS
CN 2,6-Quinoxalinedicarboxamide, N2-[3-(2-benzofuranyl)-2-butenyl]-N6, N6-diethyl-3-methyl-, 4-oxide (9CI) (CA INDEX NAME)



IT 165736-35-6P 165736-36-7P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation of quinoxaline-2-carboxamides as antidiabetics)
RN 165736-35-6 CAPLUS
CN 2-Quinoxalinecarboxylic acid, 6-[(diethylamino)carbonyl]-3-methyl-, ethyl ester, 1,4-dioxide (9CI) (CA INDEX NAME)




RN 165736-36-7 CAPLUS
CN 2-Quinoxalinecarboxylic acid, 6-[(diethylamino)carbonyl]-3-methyl-, ethyl ester, 4-oxide (9CI) (CA INDEX NAME)



L13 ANSWER 84 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN


ACCESSION NUMBER: 1995:538699 CAPLUS
DOCUMENT NUMBER: 123:265
TITLE: Hypoxia-Selective Agents Derived from Quinoxaline 1,4-Di-N-oxides
AUTHOR(S): Monge, Antonio; Palop, Juan A.; de Ceraín, Adela Lopez; Senador, Virginia; Martínez, Francisco J.; Sainz, Yolanda; Narro, Susana; García, Estrella; de Miguel, Carlos; et al.
CORPORATE SOURCE: Department of Medicinal Chemistry, Universidad de Navarra, Pamplona, 31080, Spain
SOURCE: Journal of Medicinal Chemistry (1995), 38(10), 1786-92
CODEN: JMCMAH; ISSN: 0022-2623
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



AB Hypoxic cells, which are a common feature of solid tumors, but not normal tissues, are resistant to both anticancer drugs and radiation therapy. Thus the identification of drugs with selective toxicity toward hypoxic cells is an important objective in anticancer chemotherapy. The benzotriazine di-N-oxide (SR 4233, Tirapazamine) has been shown to be an efficient and selective cytotoxin for hypoxic cells. Since the bioreductive activation of Tirapazamine is thought to be due to the presence of the 1,4-di-N-oxide moiety, a series of 3-aminoquinoxaline-2-carbonitrile 1,4-di-N-oxides with a range of electron-donating and -withdrawing substituents in the 6- and/or 7- positions has been synthesized and evaluated for toxicity to hypoxic cells. Electrochem. studies of the quinoxaline di-N-oxides and Tirapazamine showed that as the electron-withdrawing nature of the 6(7)-substituent increases, the reduction potential becomes more pos. and the compound is more readily reduced. Apart from the unsubstituted derivative and the 6,7-di-Me derivative I, the quinoxaline di-N-oxides have reduction potentials significantly more pos. than

Tirapazamine (Epc -0.90 V). The most potent cytotoxins to cells in culture were the 6,7-dichloro and 6,7-difluoro derivs. II and III, which were 30-fold more potent than Tirapazamine. The 6(7)-fluoro and 6(7)-chloro compds., IV and V, showed the greatest hypoxia selectivity. Four of the compds., IV, VI, III and II, killed the inner cells of multicellular tumor spheroids in vitro. In vivo Balb/c mice tolerated a dose of these four compds. twice the size of that of Tirapazamine. This study demonstrates that quinoxaline 1,4-di-N-oxides could provide useful hypoxia-selective therapeutic agents.

IT 163777-43-SP
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PRSP (Preparation)
(hypoxia-selective agents derived from quinoxaline di-N-oxides)
RN 163777-45-5 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 3-amino-2-cyano-, ethyl ester, 1,4-dioxide (9CI) (CA INDEX NAME)




L13 ANSWER 85 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 1995:389748 CAPLUS
DOCUMENT NUMBER: 122:14768
TITLE: Preparation of indenquinoxaline derivatives for electrophotographic photoreceptors
INVENTOR(S): Gondaiva, Hideaki; Hamamoto, Isami; Nagasaki, Fumihiko; Takahashi, Hiroshi
PATENT ASSIGNEE(S): Nippon Soda Co. Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 9 pp.
CODEN: JKKXAP
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 06298744	A2	19941025	JP 1993-113861	19930416
PRIORITY APPLN. INFO.:			JP 1993-113861	19930416


GI




AB The title compds. [I; A, B = cyano, NO2, halo, (un)substituted alkyl, alkenyl, alkynyl, aryl, alkoxy, carbonyl, aryloxy, acylamino, alkoxy, alkenyloxy, alkynyloxy, or aryloxy; k, l = 0-4; when k or l = 2, A or B is same or different; Z = O, S, N(W), C(X)(Y); wherein W = (un)substituted alkyl, alkenyl, alkynyl, aryl, or heterocyclyl, cyano, NO2; X, Y = H, (un)substituted alkyl, aryl, alkoxy, carbonyl, aryloxy, carbonyl, alkoxy, acylamino, or aryloxy], having excellent electron-transport ability, are prepared. An electrophotog. photoreceptor comprises a photosensitive layer containing 21 above compds. I as a charge-transport material, formed on a conductive support. Thus, 9.0 g ninhydrin and 7.6 g 3,4-diaminobenzoic acid was dissolved in EtOH and refluxed for 3 h to give a mixture of indenquinoxalinone derivative (II; Z = O, R = CO2H, R1 = H) and regioisomer II (Z = O, R = H, R1 = CO2H) in 98% yield which (7.0 g) was esterified with BuOH in the presence of concentrated H2SO4 in refluxing toluene with removal of H2O through a Dean-Stark apparatus to give 11.9% Bu ester II (Z = O, R = CO2Bu, R1 = H) and 70% regioisomer II (Z = O, R = H, R1 = CO2Bu). The latter regioisomer (0.4 g) was refluxed with malononitrile in the presence of piperidine in MeOH with stirring for 24 h to give 97% title compound II (Z = C(CN)2, R = H, R1 = CO2Bu) (III). An electrophotog. photoreceptor with a charge-transport layer containing III coated on an Al substrate was charged by a corona discharge at +60 kV, left for 30 s in dark, and exposed with a 10-lx halogen lamp to show maximum electrification potential (Vmax) of 420 V, half-reduction exposure dose (E1/2) 15.0 l.s., and residual potential 110 V.

IT 161290-88-6P 161290-91-1P 161290-92-2P
RL: DEV (Device component use); SPN (Synthetic preparation); PRSP (Preparation); USES (Uses)
(Preparation of indenquinoxaline derivs. as charge-transport materials for electrophotog. photoreceptors)


RN 161290-88-6 CAPLUS
CN 11H-Indeno[1,2-b]quinoxaline-7-carboxylic acid, 11-(dicyanomethylene)-, butyl ester (9CI) (CA INDEX NAME)



RN 161290-91-1 CAPLUS
CN 11H-Indeno[1,2-b]quinoxaline-8-carboxylic acid, 11-oxo-, butyl ester (9CI) (CA INDEX NAME)



RN 161290-92-2 CAPLUS
CN 11H-Indeno[1,2-b]quinoxaline-7-carboxylic acid, 11-oxo-, butyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 86 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 1995:267133 CAPLUS
DOCUMENT NUMBER: 123:259652
TITLE: Process for dyeing substrates with dyes containing nucleophilic and electrophilic groups and dyes for Renfrew, Andrew Hunter Morris; Shawcross, Andrew Paul
INVENTOR(S): Zeneca Ltd., UK
SOURCE: Brit. UK Pat. Appl., 84 pp.
CODEN: BAKXDU
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
GB 2272914	A1	19940601	GB 1993-23560	19931115
GB 2272914	B2	19960117		
WO 9412717	A1	19940609	WO 1993-082344	19931115
W: AT, AU, BB, BR, BY, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MK, MN, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, UZ, VN				
RM: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CP, CO, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AU 9454311	A1	19940622	AU 1994-54311	19931115
EP 639237	A1	19950222	EP 1993-924767	19931115
EP 639237	B1	19970604		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
JP 08503981	T2	19960430	JP 1993-512875	19931115
AT 154079	E	19970615	AT 1993-924767	19931115
ES 2102690	T3	19970801	ES 1993-924767	19931115
ZA 9308553	A	19940720	ZA 1993-8553	19931116
CN 1080304	A	19940803	CN 1993-114956	19931120
US 5474580	A	19951212	US 1993-158220	19931129
US 5703215	A	19971230	US 1995-436822	19950530

PRIORITY APPLN. INFO.:

GB 1992-24909 A 19921127
GB 1993-12205 A 19930614
WO 1993-02344 W 19931115
US 1993-158220 A3 19931129

OTHER SOURCE(S):

MARPAT 123:259662

AB A process for the coloration of a substrate, especially a textile, comprises applying to the substrate a mixture comprising an aqueous solvent and a water-soluble dye which contains a nucleophilic group and an electrophilic group and heating or basifying or heating and basifying the dye thereby causing mole. of the dye to join together. In this process the mol. weight of the dye increases, its water-solubility can decrease, and the affinity for textiles may be increased leading to high levels of exhaustion with good fixation and washfastness. Also claimed are polymers and oligomers of the dyes.


IT 168189-19-3P

RL: IMF (Industrial manufacture); NUU (Other use, unclassified); PRP (Physical, engineering or chemical process); PRP (Preparation); PROC (Process); USES (Uses)
(process for dyeing substrates with dyes containing nucleophilic and electrophilic groups and dyes for)

RN 168189-19-3 CAPLUS


CN 6-Quinoxalinecarboxamide, N-[3-[[ethyl(4-[[[2-(sulfoxy)ethyl]sulfonyl]phenyl]azo]phenyl]amino]propyl]-1,2,3,4-tetrahydro-2,3-dithioxo-, monosodium salt (9CI) (CA INDEX NAME)

PAGE 1-A



• Na

PAGE 1-B



L13 ANSWER 87 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER:

1994:405125 CAPLUS

DOCUMENT NUMBER:

121:205125

TITLE:

Preparation of [[carboxyheterocyclyl]carbamoyl]pyrrolidinylthiocarbapenems as antibiotics

INVENTOR(S):

Jung, Frederic Henri; Arnould, Jean Claude

PATENT ASSIGNEE(S):

Zeneca Ltd., UK; Zeneca Pharma S.A.

SOURCE:

Eur. Pat. Appl., 27 pp.

DOCUMENT TYPE:

CODEN: EPXXDW

LANGUAGE:

Patent


FAMILY ACC. NUM. COUNT:

English

PATENT INFORMATION:

1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 561500	A1	19940202	EP 1993-305607	19930716
EP 561500	B1	19980909		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
CA 2099818	AA	19940122	CA 1993-2099818	19930705
AT 170859	S	19980915	AT 1993-305607	19930716
ES 2121585	T3	19981201	ES 1993-305607	19930716
JP 06179674	A3	19940628	JP 1993-177903	19930719
US 5441949	A	19950815	US 1994-307048	19940916
PRIORITY APPLN. INFO.:			EP 1992-402105	A 19920721
OTHER SOURCE(S):			US 1993-66836	B1 19930707
GI				




AB Title compds. [I; R1 = MeCH(OH), MeCHF, CH2OH; R2,R3 = H, alkyl; Z = (iso)quinolinediyl, quinoxalinediyl, quinoxalinediyl, etc.] were prepared Thus, disodium (1R,5S,6S,8R,2'S,4'S)-2-[[2-(8-carboxyquinol-6-ylcarbamoyl)pyrrolidin-4-ylthio]-6-(1-hydroxyethyl)-1-methylcarbamapenem-3-carboxylate, prepared in 5 steps from 6-amino-8-carboxyquinoline (preparation given), had MIC of 0.13 and 0.03 µg/mL against Staphylococcus aureus Oxford and Escherichia coli DCO, resp.

IT 157915-55-4P 157915-56-5P 157915-57-6P

RL: RCT (Reactant); SPN (Synthetic preparation); PRP (Preparation); RACT (Reactant or reagent)
(preparation and reaction of, in preparation of antibiotic)


RN 157915-55-4 CAPLUS

CN 6-Quinoxalinecarboxylic acid, 2,3-dimethyl-8-nitro-, 2-propenyl ester (9CI) (CA INDEX NAME)



RN 157915-56-5 CAPLUS


CN 6-Quinoxalinecarboxylic acid, 8-amino-2,3-dimethyl-, 2-propenyl ester (9CI) (CA INDEX NAME)



RN 157915-57-6 CAPLUS

CN 6-Quinoxalinecarboxylic acid, 8-[[[4-(acetylthio)-1-[[2-(2-propenyl)oxy]carbonyl]-2-pyrrolidinyl]carbonyl]amino]-2,3-dimethyl-, 2-propenyl ester, (2S-cis)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 157915-58-7 CAPLUS

CN 1-Azabicyclo[3.2.0]hept-2-ene-2-carboxylic acid, 6-[[1-hydroxyethyl]-4-methyl-7-oxo-3-[[[1-[[2-(2-propenyl)oxy]carbonyl]-5-[[[2,3-dimethyl-7-[[2-(2-propenyl)oxy]carbonyl]-5-quinoxaliny]amino]carbonyl]-3-pyrrolidinyl]thio]-2-propenyl ester, (4R-[3(2S*,4S*),4 a,5B,6B(R*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L13 ANSWER 88 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER:

1993:490784 CAPLUS

DOCUMENT NUMBER:


119:90784

TITLE:

Polycyclic compounds for cancer diagnosis and therapy

INVENTOR(S): Tai, Seiji; Katayama, Mitsuo; Morishita, Yoshii
PATENT ASSIGNEE(S): Hitachi Chemical Co., Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 15 pp.
CODEN: JKKXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04288022	A2	19921013	JP 1991-49379	19910314
PRIORITY APPLN. INFO.:			JP 1991-49379	19910314
OTHER SOURCE(S):			MARPAT 119:90784	
GI				




AB Polycyclic compds. I [M = H, Al, Si, P, Ge, Cd, Se, Mg, Sn, Zn; R1-4 = H, XCH, OM, W (X = O, N, S, P, Si, CRSR6 (R1-6 = H, alkyl, aryl, aralkyl, etc.); Q = X-W linkage; W = OH, O, SH, S, etc.); k, l, m, n = 0-4; Y = halo, OR7, NR8 (R7-8 = H, (un)substituted alkyl, etc.); p = 0-2; Z1-8 = methylene, N] are reagents for cancer diagnosis or therapy. Thus, Na chloroaluminonaphthalocyaninotriulfonate (1 + 10-5M) was injected into peritoneal cancer cell-bearing mice, and the treated cancer cells were sampled (isolated) and examined at 760 nm. The cancer cells were readily detected. Preparation of the compds. are given.

IT 145964-97-2

RL: RCT (Reactant); RACT (Reactant or reagent)
(reaction of, for polycyclic cyano compound preparation for cancer diagnosis and therapy)

RN 145964-97-2 CAPLUS

CN 6-Quinoxalinecarboxylic acid, 2,3-dicyano-, pentyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 89 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1993:191763 CAPLUS
 DOCUMENT NUMBER: 118:191763
 TITLE: Preparation of azamethine compounds as optical recording media
 INVENTOR(S): Nagasaki, Fumihiko; Hayashi, Yukio
 PATENT ASSIGNEE(S): Nippon Soda Co., Ltd., Japan
 SOURCE: Jpn. Kokai Tokkyo Koho, 16 pp.
 CODEN: JKKXAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04288049	A2	19921013	JP 1991-40710	19910214
PRIORITY APPLN. INFO.:			JP 1991-40710	19910214
OTHER SOURCE(S):		CASREACT 118:191763; MARPAT 118:191763		

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


AB 1,3-Bis(dicyanomethylene)-3-[(hetero)arylimino]-2,3-dihydrocyclopenta[b]quinoxaline and -5,6-benzindenes [I: Y = N, CH; X = O; Q; R1-R4 = H, (un)substituted alkyl, alkoxy, or NH2, halo, NO2, cyano, OR, etc.; R5, R6 = H, (un)substituted alkyl, aryl, cycloalkyl; or R5R6 forms (hetero atom-containing) ring; R7 = H, (un)substituted alkoxy, HO, halo, cyano, acylamino, (un)substituted alkanoyloxy; R8 = H, halo, (un)substituted alkyl, alkoxy; R9-R11 = H, alkyl] are prepared. I showed maximum absorption wavelengths (λ_{max} = 750-900 nm) in a semiconductor oscillation region, excellent solubility in organic solvents, high reflectivity, and excellent stability. Thus, 3.0 g cyclopenta[b]quinoxaline (II; Z = H2), 2.1 g nitrobenzene QNO (R5 = R6 = St, R7 = OMe, R8 = H), and 70 mL Ac2O were stirred at room temperature for 10 h to give II (Z = NO, R5 = R6 =

Et, R7 = OMe, R8 = H) (III) having λ_{max} = 825 nm. A solution of III in CHCl3 was spin-coated on a glass substrate and dried to form a recording medium of approx. 900 Å thickness having λ_{max} = 940 nm and 23% reflectivity at λ_{max} = 830 nm which formed a very clear pit by irradiation with Ga-Al-As semiconductor laser beam.


IT 146677-81-8P 146677-98-7P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of, as intermediate for optical recording material)
 RN 146677-81-8 CAPLUS
 CN 1H-Cyclopenta[b]quinoxaline-6-carboxylic acid, 1,3-bis(dicyanomethylene)-2,3-dihydro-, ethyl ester (9CI) (CA INDEX NAME)



RN 146677-98-7 CAPLUS
 CN 1H-Cyclopenta[b]quinoxaline-6-carboxylic acid, 1,1,3,3-tetrachloro-2,3-dihydro-, ethyl ester (9CI) (CA INDEX NAME)




IT 146677-62-5P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of, as optical recording material)
 RN 146677-62-5 CAPLUS
 CN 1H-Cyclopenta[b]quinoxaline-6-carboxylic acid, 1,3-bis(dicyanomethylene)-2-[[4-(diethylamino)-2-methoxyphenyl]imino]-2,3-dihydro-, ethyl ester (9CI) (CA INDEX NAME)




L13 ANSWER 90 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1992:532895 CAPLUS
 DOCUMENT NUMBER: 117:132895
 TITLE: Tautomerism of 5-dicyanomethylene-9-diethylamino-5,7-dihydrobenzo[e]phenazine and its photooxygenation to an ester in alcohol solution
 AUTHOR(S): Kubo, Yuji; Kuwana, Minoru; Teutsui, Sumica; Yoshida, Katsuhira
 CORPORATE SOURCE: Fac. Sci., Kochi Univ., Kochi, 780, Japan
 SOURCE: Journal of Chemical Research, Synopses (1992), (8), 282-3
 CODEN: JRPSCD; ISSN: 0308-2342
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The title dye exhibits a dicyanomethylenephhenazine-phenazylmalononitrile tautomerism in solution, the latter tautomer of which undergoes photooxygenation in alc. solns. to afford novel alkyl phenazincarboxylates.

IT 143413-98-3P 143413-99-4P
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
 (preparation and spectra of)
 RN 143413-98-3 CAPLUS
 CN Benzo[a]phenazine-5-carboxylic acid, 9-(diethylamino)-, ethyl ester (9CI) (CA INDEX NAME)




RN 143413-99-4 CAPLUS
 CN Benzo[a]phenazine-5-carboxylic acid, 9-(diethylamino)-, butyl ester (9CI) (CA INDEX NAME)




L13 ANSWER 91 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1992:59404 CAPLUS
 DOCUMENT NUMBER: 116:59404
 TITLE: Preparation of 2,3-disubstituted quinoxalines as growth enhancers for animals
 INVENTOR(S): De Jong, Anno; Fuchs, Rainer
 PATENT ASSIGNEE(S): Bayer A.G., Germany
 SOURCE: Eur. Pat. Appl., 19 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 456067	A1	19911113	EP 1991-106891	19910427
R: AT, BS, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
DE 4014937	A1	19911114	DE 1990-4014937	19900510
PRIORITY APPLN. INFO.:			DE 1990-4014937	A 19900510
OTHER SOURCE(S):		MARPAT 116:59404		




AB Title comds. [I: R1, R2 = halo, OH, SH, alkoxy, aryloxy, alkylthio, arylthio; R1R2 = SC(X)S; X = O, S; R3 = H, halo, cyano, CONRSR6, SO2NRSR6, SO2ORS, CH(CN)R7, OR7, CO2R5; R4 = H, alkyl, halo, NO2, CONRSR6; R5 = H, alkyl, (substituted) cycloalkyl, aryl; R6 = R5, SR7; R5R6 = (substituted) heterocyclyl; R7 = (substituted) alkyl, aryl], were prepared. Thus, H2NCHMeCF3, pyridine, and 2,3-dichloroquinoxaline-6-carbonyl chloride were stirred 3 h in MeCN at 80° to give title compound II. II at 0.69 g/day orally in sheep increased propionic acid in rumen liquid from 16.9 mol % (controls) to 21.0%.


IT 26773-25-1P 138452-89-8P 138452-90-1P
 138452-91-2P 138452-96-7P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of, as animal growth enhancer)
 RN 26773-25-1 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-ethyl- (8CI, 9CI) (CA INDEX NAME)




RN 138452-89-8 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2,2,2-trifluoro-1-methylethyl)- (9CI) (CA INDEX NAME)




RN 138452-90-1 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(3,3,3-trifluoropropyl)- (9CI) (CA INDEX NAME)



RN 138452-91-2 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2,2,2-trifluoroethyl)- (9CI)
(CA INDEX NAME)



RN 138452-96-7 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(1-methylethyl)- (9CI) (CA INDEX NAME)




L13 ANSWER 92 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1991:682057 CAPLUS
DOCUMENT NUMBER: 115:282057
TITLE: Manufacture of storage-stable dye solutions
INVENTOR(S): Michne, Martin; Zilliger, Hans Werner; Tegtmeyer, Dietrich
PATENT ASSIGNER(S): Bayer A.-G., Germany
SOURCE: Eur. Pat. Appl., 8 pp.
CODEN: EPXKDW
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 433810	A2	19910626	EP 1990-123612	19901208
EP 433810	A3	19920115		
R: CH, DE, FR, GB, LI				
DE 3942467	A1	19910627	DE 1989-3942467	19891222
US 5096458	A	19920317	US 1990-627068	19901213
JP 04004263	A2	19920108	JP 1990-411097	19901217
			DE 1989-3942467	19891222

PRIORITY APPLN. INFO.:
OTHER SOURCE(S): MARPAT 115:282057
AB Stable aqueous solns. of anionic (preferably reactive) dyes are obtained by pressure filtration of crude dye solns. in which the feed solution is obtained by stirring the optionally dried press cake or a suspension of the crude dye with a solution of Li or ammonium salts of organic or inorg. acids. Thus, 35.18 kg press cake of 1-hydroxy-2-(1,5-disulfo-2-naphthylazo)-6-(2,4-difluoro-5-chloro-4-pyrimidinylamino)-3-naphthalenesulfonic acid Na salt (I) was dissolved in 226.7 kg 4% aqueous LiHCO₃ solution at 45°. The composition was subjected to membrane filtration at 40 bars and 40-45°. The concentrate (94 kg) was treated with dicyandiamide 2, water 3.5, and boric acid 0.5 kg to give a stable dye solution containing 21.2% I at pH 7.5.
IT 137682-84-9

RL: USES (Uses)
(dye, storage-stable aqueous solns. containing)

RN 137682-84-9 CAPLUS
CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxalyl)carbonyl]methyl]amino]methyl]-2-sulfonylphenyl]amino]-9,10-dihydro-9,10-dioxo- (9CI) (CA INDEX NAME)




L13 ANSWER 93 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1991:679818 CAPLUS
DOCUMENT NUMBER: 115:279818
TITLE: Preparation of piperidine derivatives as neurokinin and substance P antagonists
INVENTOR(S): Emonds-Alt, Xavier; Goulaouic, Pierre; Proietto, Vincenzo; Van Broeck, Didier
PATENT ASSIGNER(S): SANOFI, Fr.
SOURCE: Eur. Pat. Appl., 84 pp.
CODEN: EPXKDW
DOCUMENT TYPE: Patent
LANGUAGE: French
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 428434	A2	19910522	EP 1990-403125	19901106
EP 428434	A3	19911009		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
FR 2654100	A1	19910510	FR 1989-14517	19891106
FR 2654100	B1	19920221		
FR 2663329	A1	19911220	FR 1990-7534	19900615
FR 2663329	B1	19921016		
FI 97540	B	19960930	FI 1990-5444	19901102
FI 97540	C	19970110		
CA 2029275	AA	19910507	CA 1990-2029275	19901105
NO 9004802	A	19910507	NO 1990-4802	19901105
NO 177299	B	19950515		
NO 177299	C	19950823		
AU 9065838	A1	19910523	AU 1990-65838	19901105
AU 649973	B2	19940609		
HU 56543	A2	19910930	HU 1990-7027	19901105

US	111292	A1	19960331	IL 1990-111292	19901105
RU	2084453	C1	19970720	RU 1990-4831627	19901105
RU	2114828	C1	19980710	RU 1993-45020	19901105
ZA	9008881	A	19910828	ZA 1990-8881	19901106
JP	03206086	A2	19910909	JP 1990-300929	19901106
PL	165758	B1	19950228	PL 1990-293823	19901106
PL	165854	B1	19950228	PL 1990-293824	19901106
PL	166565	B1	19950630	PL 1990-287644	19901106
PL	166582	B1	19950630	PL 1990-303827	19901106
IL	96241	A1	19960331	IL 1990-96241	19901115
LV	10713	B	19951020	LV 1993-142	19902225
US	5686609	A	19971111	US 1994-208672	19940311
AU	9459245	A1	19940602	AU 1994-59245	19940331
AU	668018	B2	19960418		
NO	9500239	A	19910507	NO 1995-239	19950123
NO	180193	B	19961125		
NO	180193	C	19970308		
NO	9500240	A	19910507	NO 1995-240	19950123
NO	179580	B	19960729		
NO	179580	C	19961106		
US	5618938	A	19970408	US 1995-479634	19950607
FI	9502956	A	19950615	FI 1995-2956	19950615
FI	9502957	A	19950615	FI 1995-2957	19950615
FI	9800227	A	19980202	FI 1998-227	19980202
				FR 1989-14517	19891106
				FR 1990-7534	19900615
				FI 1990-5444	19901102
				NO 1990-4802	19901105
				US 1990-610093	19901105
				IL 1990-96241	19901115
				US 1994-208672	19940311
				FI 1995-2956	19950615


PRIORITY APPLN. INFO.:
OTHER SOURCE(S): MARPAT 115:279818
GI



AB The title compds. I [m = 1-3; Ar, Ar' = thienyl, (substituted) Ph, etc.; X = H; X' = H, OH; or XX' = oxo, dialkylaminoalkoxyimino, etc.; Y = N, CX'; X' = H or X' = carbon-carbon bond; Q = H, alkyl, (CH₂)_q; q = 2 or 3; Rm = piperidino, 4-benzylpiperidino, etc.; R = H, Me, (CH₂)_n; n = 2-6; L = H, amino; T = CO, C(W)NH; W = O, S; Z = H, M, or OM when T = CO; or Z = M when T = C(W)NH; M = H, alkyl, (substituted) phenylalkyl, etc.] were prepared. I are neurokinin and substance P antagonists (no data). Reaction of amine II (Zl = H) with 2,4-dichlorobenzoyl chloride in the presence of Et₃N gave II (Zl = 2,4-dichlorobenzoyl) isolated as its HCl salt. I are also useful as allergy and inflammation inhibitors (no data).
IT 135956-48-8P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of, as neurokinin antagonist)

RN 135956-48-8 CAPLUS
CN 6-Quinoxalinecarboxamide, N-[2-(3,4-dichlorophenyl)-4-[4-(phenylmethyl)-1-piperidinyl]butyl]-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

L13 ANSWER 94 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1991:492302 CAPLUS
DOCUMENT NUMBER: 115:92302
TITLE: Preparation of benzo[a]phenazine derivatives
INVENTOR(S): Shirai, Hiroyoshi; Hanabusa, Kenji; Ooe, Okikazu; Uda, Yoshihiro
PATENT ASSIGNER(S): Taiho Pharmaceutical Co., Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 3 pp.
CODEN: JKXKAP
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 03066698	A2	19910322	JP 1989-204211	19890807
			JP 1989-204211	19890807


PRIORITY APPLN. INFO.: MARPAT 115:92302
OTHER SOURCE(S):
GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The title derivs. I (R = O1-Q3), useful as antitumor agents (no data), are prepared. Thus, 1.90 g I (R = H) was stirred with 1.75 g II in DMF in the presence of powdered K₂CO₃ at 50-60° for 24 h to give 1.14 g I (R = Q1).
IT 135412-59-8P 135412-60-1P 135438-74-3P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use);

BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of, as antitumor agent)
RN 135412-59-8 CAPLUS
CN α -D-Glucopyranose, 1,2-O-(1-methylethylidene)-, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxybenzo[*a*]phenazine-9-carboxylate], monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● HCl

RN 135412-60-1 CAPLUS
CN α -D-Glucopyranose, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxybenzo[*a*]phenazine-9-carboxylate], monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● HCl

RN 135438-74-3 CAPLUS
CN α -D-Glucopyranose, 1,2:3,5-bis-O-(1-methylethylidene)-, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxybenzo[*a*]phenazine-9-carboxylate (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L13 ANSWER 95 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1991:420992 CAPLUS
DOCUMENT NUMBER: 115:20992
TITLE: Substituted tetraquinoxalino[2,3-b:2',3'-g:2'',3''']porphyrazine derivative with near-infrared absorption
INVENTOR(S): Negasaki, Fumihiko; Hatano, Hiromi; Takahashi, Hiroshi
PATENT ASSIGNEE(S): Nippon Soda Co., Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.
CODEN: JKKXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 02289575	A2	19901129	JP 1989-73155	19890324
PRIORITY APPL. INFO.:			JP 1989-32144	A1 19890210
OTHER SOURCE(S):	MARPAT	115:20992		
CI				


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The derivative is I [Z = H; R1-4 = halo, (substituted) alkyl, alkoxy, alkylthio, phenylthio, phenyloxy, and ester, carboxyl, amide, amino; k, l, m, n = 0-4; k + l + m + n \leq 1; M = 2H, metal metal oxide, metal hydroxide, acyl metal, metal alkoxide, metal siloxide, metal halide). The derivative, with high near-IR absorption and solubility to an organic solvent, is useful for an optical recording medium, electrophotog., photoreceptors, redox catalysts, flower preservatives, etc.
IT 134382-42-6 134382-43-7 134382-44-8
134382-45-9 134382-46-0
RL: RCT (Reactant); RACT (Reactant or reagent) (near-IR-absorbing)
RN 134382-42-6 CAPLUS
CN Copper, [tetraethyl 37H,39H-tetraquinoxalino[2,3-b:2',3'-g:2'',3''']porphyrazine-2,11,20,29-tetracarboxylato(2-)-N37,N38,N39,N40]-, (8P-4-1)- (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A




PAGE 3-A



RN 134382-43-7 CAPLUS
CN Tin, dichloro[tetraethyl 37H,39H-tetraquinoxalino[2,3-b:2',3'-g:2'',3''']porphyrazine-2,11,20,29-tetracarboxylato(2-)-N37,N38,N39,N40]-, (OC-6-12)- (9CI) (CA INDEX NAME)

PAGE 1-A





PAGE 2-A




PAGE 3-A

RN 134382-44-8 CAPLUS
CN Vanadium, oxo[tetraethyl 37H,39H-tetraquinoxalino[2,3-b:2',3'-g:2'',3'''-1:2''',3'''-q]porphyrizine-2,11,20,29-tetracarboxylato(2-)-N37,N38,N39,N40]-, (SP-5-12)- (9CI) (CA INDEX NAME)



PAGE 1-A




PAGE 2-A



PAGE 3-A

RN 134382-45-9 CAPLUS
CN Aluminum, chloro[tetraethyl 37H,39H-tetraquinoxalino[2,3-b:2',3'-g:2'',3'''-1:2''',3'''-q]porphyrizine-2,11,20,29-tetracarboxylato(2-)-N37,N38,N39,N40]-, (SP-5-12)- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A




PAGE 3-A


RN 134382-46-0 CAPLUS
CN Zinc, [tetraethyl 37H,39H-tetraquinoxalino[2,3-b:2',3'-g:2'',3'''-1:2''',3'''-q]porphyrizine-2,11,20,29-tetracarboxylato(2-)-N37,N38,N39,N40]-, (SP-4-1)- (9CI) (CA INDEX NAME)



L13 ANSWER 97 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1990:611948 CAPLUS
 DOCUMENT NUMBER: 113:211948
 TITLE: Potential antimitotic agents. Synthesis of some ethyl benzopyrazin-7-ylcarbamates, ethyl pyrido[3,4-b]pyrazin-7-ylcarbamates, and ethyl pyrido[3,4-e]-as-triazin-7-ylcarbamates
 AUTHOR(S): Temple, Carroll, Jr.; Rener, Gregory A.
 CORPORATE SOURCE: South. Res. Inst., Birmingham, AL, 35255-5305, USA
 SOURCE: Journal of Medicinal Chemistry (1990), 33(11), 3044-50
 CODEN: JMCMAH; ISSN: 0022-2623
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 113:211948
 GI




AB Ring analogs and derivs. of antimitotic antitumor 1,2-dihydropyrido[3,4-b]pyrazinylcarbamates, e.g., I, were prepared from benzoic acids, e.g., 4,3,5-trichlorobenzoic acid, and pyridylcarbamates, e.g., II. In vitro evaluation indicated that activity was reduced by removal of the pyridine ring nitrogen of I and destroyed by increasing the basicity of the pyrazine ring of I as in the case of aminopyridotriazinylcarbamates III.
 IT 130145-39-0P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation, borohydride reduction and neoplasia inhibiting activity of)
 RN 130145-39-0 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 8-amino-2-phenyl-, ethyl ester (9CI) (CA INDEX NAME)




L13 ANSWER 98 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1990:179846 CAPLUS
 DOCUMENT NUMBER: 113:179846
 TITLE: Synthesis and antimicrobial activity of some new




L13 ANSWER 96 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1991:104337 CAPLUS
 DOCUMENT NUMBER: 114:104337
 TITLE: Electrochemical investigations on the analysis of reactive dyes with monoazo- and monanthraquinone structures
 AUTHOR(S): Sahn, Uwe; Knittel, Dierk; Schollmeyer, Eckhard
 CORPORATE SOURCE: Deusch. Textilforschungszent. Nord-West e.V., Krefeld, W-4150/1, Germany
 SOURCE: Fresenius' Journal of Analytical Chemistry (1990), 338(7), 824-30
 CODEN: FJACES; ISSN: 0937-0633
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Qual. voltammetric determination of reactive dyes is possible down to concns. of .apprx.10-5 mol/L using d.c. polarog. Using rotating solid electrodes (glassy carbon) a detection limit of .apprx.10-4 mol/L can be obtained. Reduction involves chiefly only the azo- and anthraquinoid groups. Discrimination between an original reactive dye and its hydrolyzate (containing a hydrolytically destroyed reactive group) can be achieved, if the dye contains further cleavable groups such as benzamido substituents in conjugation to the chromophore as is seen with an azo red dye. Cleavage of such a group shifts the reduction potential for .apprx.130 mV to more neg. values.
 IT 132366-73-5
 RL: AMT (Analyte); ANST (Analytical study)
 (determination of, voltammetric, electrochem. reactions in)
 RN 132366-73-5 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxalinyloxy)methyl]methylamino]methyl]-3-sulfonyl]amino]-9,10-dihydro-9,10-dioxo- (9CI) (CA INDEX NAME)



2,3-dichloroquinoxaline-6-sulfonyl amino acid and dipeptide derivatives
 AUTHOR(S): Kora, F. A.; Hussein, M. E.; El-Sayed, R. A.;
 CORPORATE SOURCE: El-Naggar, Ahmed M.
 SOURCE: Fac. Sci., Al-Azhar Univ., Ngar, Egypt
 Polish Journal of Chemistry (1988), 62(7-12), 749-56
 CODEN: PJCHDQ; ISSN: 0137-5083
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 112:179846
 GI




AB Title amino acid derivs. I (X = Gly, Ala, Val, Leu, DL-Leu, Ser, DL-Ser, Phe, Tyr; R = OH) were prepared by sulfonylating the corresponding amino acids with sulfonyl chloride II in the presence of Et3N. Me esters I (X = Gly, DL-Ala, Leu, Ser; R = OMe) were prepared similarly from II and the appropriate amino acid Me ester hydrochlorides. The above Me esters were converted into hydrazides I (X = Gly, DL-Ala, Leu, Ser, R = NHNH2). Dipeptides III (X1-X2 = Phe, DL-Ala, Phe-Leu, Tyr-Gly, Tyr-DL-Ala, Tyr-Leu) were prepared by the DCC method. All the above compds. were active against a number of microorganisms.
 IT 117195-85-4P 117195-86-5P 117195-87-6P
 117195-88-7P 117195-89-8P 117195-90-1P
 117195-91-2P 117195-97-8P 117195-98-9P
 117195-99-0P 117196-00-6P 117196-01-7P
 117196-02-8P 117196-03-9P 117196-04-0P
 117196-05-1P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation and antimicrobial activity of)
 RN 117195-85-4 CAPLUS
 CN Glycine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]- (9CI) (CA INDEX NAME)




RN 117195-86-5 CAPLUS
 CN L-Alanine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117195-87-6 CAPLUS
CN L-Valine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117195-88-7 CAPLUS
CN L-Leucine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117195-89-8 CAPLUS
CN Leucine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]- (9CI) (CA INDEX NAME)




RN 117195-90-1 CAPLUS
CN L-Serine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 117195-91-2 CAPLUS
CN Serine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]- (9CI) (CA INDEX NAME)



RN 117195-97-8 CAPLUS
CN Glycine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]-, hydrazide (9CI) (CA INDEX NAME)




RN 117195-98-9 CAPLUS
CN Alanine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]-, hydrazide (9CI) (CA INDEX NAME)




RN 117195-99-0 CAPLUS
CN L-Leucine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]-, hydrazide (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117196-00-6 CAPLUS
CN L-Serine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]-, hydrazide (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117196-01-7 CAPLUS
CN Alanine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]-L-phenylalanyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117196-02-8 CAPLUS
CN L-Leucine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]-L-phenylalanyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117196-03-9 CAPLUS
CN Glycine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]-L-tyrosyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117196-04-0 CAPLUS
CN Alanine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]-L-tyrosyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117196-05-1 CAPLUS
CN L-Leucine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]-L-tyrosyl-, methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



IT 117195-93-4P 117195-94-5P 117195-95-6P
117195-96-7P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation, hydrazinolysis, and antimicrobial activity of)
RN 117195-93-4 CAPLUS
CN Glycine, N-[(2,3-dichloro-6-quinoxalinyloxy)sulfonyl]-, methyl ester (9CI) (CA INDEX NAME)




RN 117195-94-5 CAPLUS
CN Alanine, N-[(2,3-dichloro-6-quinoxaliny) sulfonyl]-, methyl ester (9CI)
(CA INDEX NAME)




RN 117195-95-6 CAPLUS
CN L-Leucine, N-[(2,3-dichloro-6-quinoxaliny) sulfonyl]-, methyl ester (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




RN 117195-96-7 CAPLUS
CN L-Serine, N-[(2,3-dichloro-6-quinoxaliny) sulfonyl]-, methyl ester (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




IT 117195-92-3P 117222-08-9P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation, peptide coupling reaction, and antimicrobial activity of)
RN 117195-92-3 CAPLUS
CN L-Tyrosine, N-[(2,3-dichloro-6-quinoxaliny) sulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117222-08-9 CAPLUS
CN L-Phenylalanine, N-[(2,3-dichloro-6-quinoxaliny) sulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L13 ANSWER 99 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1990:25680 CAPLUS
DOCUMENT NUMBER: 112:25680
TITLE: Bioavailability-improved anticancer emulsions containing benzo[a]phenazines
INVENTOR(S): Yanaguchi, Hiroshi; Ozawa, Yasuo; Kano, Akira; Hayashi, Hidefumi; Shoji, Minoru; Aihara, Hirokazu; Kotomo, Susumu; Nakaie, Shiro
PATENT ASSIGNEE(S): Taisho Pharmaceutical Co., Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 3 pp.
CODEN: JKKXAP
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 01143834	A2	19890606	JP 1987-304047	19871201
JP 06062418	B4	19940817		
PRIORITY APPLN. INFO.:			JP 1987-304047	19871201
OTHER SOURCE(S):		MARPAT 112:25680		
GI				




AB Anticancer emulsions contain benzo[a]phenazines (I: R = alkyl) with average particle size 40-70 nm. The emulsions do not show sharp decrease of I


concentration in blood as conventional emulsions with large particle size.
NC-239 (I, R = Bu) (28 mg) was dispersed in 4 g Panacetate 810 (triglyceride), homogenized with 800 mg Nikkol TO-10M [poly(oxyethylene) sorbitan fatty acid ester], 880 mg glycerin, and H2O to 40 mL (pH 7.4), charged into ampule, and sterilized to give an emulsion (average particle size 40 nm), which was i.v. administered to mice bearing lung cancer at 25 mg (as NC-239)/kg/day for 8 days to show T/C (treated group/control group) survival rate >18%, vs. 15%, for an emulsion with 250 nm average particle size.

IT 106224-68-4 106225-12-1 106225-21-2
RL: BICL (Biological study)
(antitumor emulsions containing, with improved bioavailability)


RN 106224-68-4 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, butyl ester (9CI) (CA INDEX NAME)



RN 106225-12-1 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, propyl ester (9CI) (CA INDEX NAME)




RN 106225-21-2 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, decyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 100 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN



ACCESSION NUMBER: 1989:136960 CAPLUS
DOCUMENT NUMBER: 110:136960
TITLE: Reactive dichloroquinoxaline group-containing dyes
INVENTOR(S): Jaeger, Horst; Stoehr, Frank Michael; Herd, Karl Josef; Henk, Hermann; Schwarz, Max; Koecher, Juergen
PATENT ASSIGNEE(S): Bayer A.G., Fed. Rep. Ger.
SOURCE: Ger. Offen., 62 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3707549	A1	19880922	DE 1987-3707549	19870310
EP 281898	A2	19880914	EP 1988-103052	19880301
EP 281898	A3	19890111		
EP 281898	B1	19910710		
R: CH, DE, FR, GB, LI				
PRIORITY APPLN. INFO.:			DE 1987-3707549	A 19870310
OTHER SOURCE(S):		CASREACT 110:136960; MARPAT 110:136960		
GI				




AB The title dyes (X02SD1)1-2GDN(R)Z, [D, D1 = direct bond, aromatic carbocyclic bridging groups, aromatic heterocyclic bridging group; G = chromophore residue; R = H, (un)substituted C1-4 alkyl; X = CH:CH2, CH2CH2Y; Y = alkali-cleavable substituent; Z = fiber-reactive residue], useful for dyeing or printing WO or carbonate group-containing materials, are prepared 1-Amino-8-hydroxy-3,6-naphthalenedisulfonic acid was condensed with 2,3-dichloroquinoxaline-6-carbonyl chloride, and the condensate coupled with diazotized 2-amino-6-sulfatoethylsulfonfyl-1-naphthalenesulfonic acid, producing I, which dyed cotton in a fast bluish-red shade.

IT 119385-60-3P
RL: PREP (Preparation)
(manufacture of, as reactive blue dye)
RN 119385-60-3 CAPLUS
CN 6-Quinoxalinecarboxamide, N,N'-[[[6,13-dichloro-4,11-bis[[[2-(sulfoxy)ethyl]sulfonyl]-3,10-triphenyldioxazinediyl]bis[[[2-amino-2,1-ethanediy]]bis[2,3-dichloro-N-[2-(sulfoxy)ethyl]- (9CI) (CA INDEX NAME)




IT 119385-47-6P
 RL: PREP (Preparation)
 (manufacture of, as reactive yellow dye)
 RN 119385-47-6 CAPLUS
 CN 3-Pyridinemethanesulfonic acid, 1-[2-[[[2,3-dichloro-6-quinolalyl]carbonyl]amino]ethyl]-1,2-dihydro-6-hydroxy-4-methyl-2-oxo-5-[[[1-sulfo-6-[[2-(sulfoxy)ethyl]sulfonyl]-2-naphthalenyl]azo]- (9CI) (CA INDEX NAME)



L13 ANSWER 101 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1989:23865 CAPLUS
 DOCUMENT NUMBER: 110:23865


TITLE: Preparation of 5-hydroxybenzo[a]phenazine-6-carboxylates as intermediates for antitumor agents
 INVENTOR(S): Uda, Yoshihiro; Kumazawa, Yukinari; Nakagami, Yoji; Asano, Takehiro; Soda, Kaoru; Sakakibara, Nisaku
 PATENT ASSIGNER(S): Taiho Pharmaceutical Co., Ltd., Japan
 SOURCE: Jpn. Kokai Tokkyo Koho, 8 pp.
 CODEN: JKXXAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 63083073	A2	19880413	JP 1986-229104	19860927
JP 06076393	B4	19940928		
PRIORITY APPL. INFO.:			JP 1986-229104	19860927
OTHER SOURCE(S):		MARPAT 110:23865		




AB Title compds. I (R1 = H, halo, Me, OH, alkoxy; R2 = CO2R4, CONSR5R6; R3 = alkyl; R4 = H, alkyl, cycloalkyl, PhCH2, Ph; R5, R6 = H, alkyl; RSR6N = heterocyclyl) are prepared as intermediates for benzo[a]phenazine-6-carboxamide antitumor agents. Treatment of Et 3-hydroxy-1,4-dihydro-1,4-dioxo-2-naphthoate with ClCO2Et in THF in the presence of Et3N gave Et 3-ethoxycarbonyloxy-1,4-dihydro-1,4-dioxo-2-naphthoate, followed by cyclocondensation with Me 4,5-diamino-2-methoxybenzoate in DMF gave I (R1 = MeO, R2 = MeOC, R3 = Et), which was refluxed with Me2N(CH2)2NH2 in C6H6 to afford the corresponding amide.

IT 106224-68-4P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of, as antitumor agent)
 RN 106224-68-4 CAPLUS
 CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, butyl ester (9CI) (CA INDEX NAME)




IT 106224-71-9P 106224-72-0P 106224-74-3P
 106224-75-3P 106224-76-4P 106224-78-6P


106224-81-1P 106224-87-7P 106224-90-2P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of, as intermediate for antitumor agents)
 RN 106224-71-9 CAPLUS
 CN Benzo[a]phenazine-6,9-dicarboxylic acid, 10-ethoxy-5-hydroxy-, diethyl ester (9CI) (CA INDEX NAME)




RN 106224-72-0 CAPLUS
 CN Benzo[a]phenazine-6,9-dicarboxylic acid, 10-ethoxy-5-hydroxy-, 9-butyl 6-ethyl ester (9CI) (CA INDEX NAME)




RN 106224-74-2 CAPLUS
 CN Benzo[a]phenazine-6,9-dicarboxylic acid, 5-hydroxy-10-propoxy-, diethyl ester (9CI) (CA INDEX NAME)




RN 106224-75-3 CAPLUS
 CN Benzo[a]phenazine-6,9-dicarboxylic acid, 5-hydroxy-10-propoxy-, 6-ethyl 9-propyl ester (9CI) (CA INDEX NAME)




RN 106224-76-4 CAPLUS
 CN Benzo[a]phenazine-6,9-dicarboxylic acid, 5-hydroxy-10-propoxy-, 9-butyl 6-ethyl ester (9CI) (CA INDEX NAME)



RN 106224-78-6 CAPLUS
 CN Benzo[a]phenazine-6,9-dicarboxylic acid, 10-butoxy-5-hydroxy-, 9-butyl 6-ethyl ester (9CI) (CA INDEX NAME)




RN 106224-81-1 CAPLUS
 CN Benzo[a]phenazine-6,9-dicarboxylic acid, 5-hydroxy-, diethyl ester (9CI) (CA INDEX NAME)




RN 106224-87-7 CAPLUS

CN Benzo[*al*]phenazine-6-carboxylic acid, 9-[(ethylamino)carbonyl]-5-hydroxy-10-methoxy-, ethyl ester (9CI) (CA INDEX NAME)



RN 106224-90-2 CAPLUS

CN Benzo[*al*]phenazine-6,9-dicarboxylic acid, 5-hydroxy-10-methoxy-, 9-(2,2-dimethylpropyl) 6-ethyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 102 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1988:590818 CAPLUS

DOCUMENT NUMBER: 109:190818

TITLE: Synthesis and antimicrobial activity of some new 2,3-dichloroquinaxaline-6-sulfonyl amino acid and dipeptide derivatives

AUTHOR(S): Kora, F. A.; Hussein, M. E.; El-Sayed, R. A.;

El-Maggar, A. M.

CORPORATE SOURCE: Fac. Sci., Al-Azhar Univ., Naar, Egypt

SOURCE: Journal of the Serbian Chemical Society (1987), 52(9),



529-35

DOCUMENT TYPE: CODEN: JSCSEN; ISSN: 0352-5139

LANGUAGE: Journal

OTHER SOURCE(S): English

GI CASREACT 109:190818



AB Title amino acids I (X = Gly, Ala, Val, Leu, DL-Leu, Ser, DL-Ser, Phe, Tyr; R = OH) and Me esters I (X = Gly, DL-Ala, Leu, Ser; R = OMe) were prepared by treating sulfonyl chloride II with the appropriate amino acids and amino acid Me esters. Hydrazides I (X = Gly, DL-Ala, Leu, Ser; R = NHNH2) were prepared by treating the corresponding Me esters with NH2NH2. Dipeptide Me esters I (X = Phe-DL-Ala, Phe-Leu, Tyr-Gly, Tyr-DL-Ala,

Tyr-Leu; R = OMe) were also prepared. The above compds. were active against a number of microorganisms.

IT 117195-85-4P 117195-86-5P 117195-87-6P

117195-88-7P 117195-89-8P 117195-90-1P

117195-91-2P 117195-97-8P 117195-98-9P

117195-99-0P 117196-00-6P 117196-01-7P

117196-02-8P 117196-03-9P 117196-04-0P


117196-05-1P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation and antimicrobial activity of)

RN 117195-85-4 CAPLUS


CN Glycine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)



RN 117195-86-5 CAPLUS

CN L-Alanine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 117195-87-6 CAPLUS

CN L-Valine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 117195-88-7 CAPLUS


CN L-Leucine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 117195-89-8 CAPLUS


CN Leucine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)



RN 117195-90-1 CAPLUS


CN L-Serine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117195-91-2 CAPLUS

CN Serine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)




RN 117195-97-8 CAPLUS

CN Glycine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]-, hydrazide (9CI) (CA INDEX NAME)



RN 117195-98-9 CAPLUS


CN Alanine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]-, hydrazide (9CI) (CA INDEX NAME)



RN 117195-99-0 CAPLUS

CN L-Leucine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]-, hydrazide (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 117196-00-6 CAPLUS

CN L-Serine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]-, hydrazide (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 117196-01-7 CAPLUS

CN Alanine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]-L-phenylalanyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117196-02-8 CAPLUS

CN L-Leucine, N-[(2,3-dichloro-6-quinoxaliny)lsulfonyl]-L-phenylalanyl-, methyl ester (9CI) (CA INDEX NAME)


methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117196-03-9 CAPLUS
CN Glycine, N-[(2,3-dichloro-6-quinoxaliny)sulfonyl]-L-tyrosyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117196-04-0 CAPLUS
CN Alanine, N-[(2,3-dichloro-6-quinoxaliny)sulfonyl]-L-tyrosyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117196-05-1 CAPLUS
CN L-Leucine, N-[(2,3-dichloro-6-quinoxaliny)sulfonyl]-L-tyrosyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 117195-93-4P 117195-94-5P 117195-95-6P
117195-96-7P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation, hydrazinolysis, and antimicrobial activity of)

RN 117195-93-4 CAPLUS
CN Glycine, N-[(2,3-dichloro-6-quinoxaliny)sulfonyl]-, methyl ester (9CI) (CA INDEX NAME)




RN 117195-94-5 CAPLUS
CN Alanine, N-[(2,3-dichloro-6-quinoxaliny)sulfonyl]-, methyl ester (9CI) (CA INDEX NAME)




RN 117195-95-6 CAPLUS
CN L-Leucine, N-[(2,3-dichloro-6-quinoxaliny)sulfonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117195-96-7 CAPLUS
CN L-Serine, N-[(2,3-dichloro-6-quinoxaliny)sulfonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 117195-92-3P 117222-08-9P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation, peptide coupling, and antimicrobial activity of)
RN 117195-92-3 CAPLUS
CN L-Tyrosine, N-[(2,3-dichloro-6-quinoxaliny)sulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 117222-08-9 CAPLUS
CN L-Phenylalanine, N-[(2,3-dichloro-6-quinoxaliny)sulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L13 ANSWER 103 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1988:38318 CAPLUS
DOCUMENT NUMBER: 108:38318
TITLE: Synthesis and biological activity of some new 2,3-dihydroxyquinoxaline-6-sulfonyl amino acids and dipeptide derivatives
AUTHOR(S): El-Naggar, A. M.; Kora, F. A.; El-Sayed, R. A.
CORPORATE SOURCE: Fac. Sci., Al-Azhar Univ., Cairo, Egypt
SOURCE: Journal of the Serbian Chemical Society (1986), 51(9-10), 441-7
CODEN: JSCSBN; ISSN: 0352-5139
DOCUMENT TYPE: Journal
LANGUAGE: English


GI



AB Title amino acid deriva. I (X = Ala, DL-Ala, Val, DL-Val, Leu, Phe, etc.) were prepared by sulfonylating the corresponding amino acid with sulfonyl chloride II. Some of the above amino acid deriva. were converted into their Me esters and hydrazides. I (X = Val, Leu, Phe, Tyr) were coupled with amino acid Me esters by the DCC method to give the corresponding dipeptide deriva. All of the above synthesized deriva. were active against a number of microorganisms, e.g., Bacillus cerues and Candida utilis.


IT 112169-97-8P 112169-98-9P 112169-99-0P
112170-00-0P 112170-01-1P 112170-02-2P
112170-03-3P 112170-04-4P 112170-05-5P
112170-06-6P 112170-07-7P 112170-08-8P
112170-18-0P 112170-19-1P 112170-26-0P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(preparation and antimicrobial activity of)
RN 112169-97-8 CAPLUS
CN L-Leucine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-L-tyrosyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112169-98-9 CAPLUS
CN L-Leucine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-L-phenylalanyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112169-99-0 CAPLUS
CN L-Leucine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-L-leucyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-00-0 CAPLUS
CN L-Leucine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-L-valyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-01-1 CAPLUS
CN Alanine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-L-valyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-02-2 CAPLUS
CN L-Phenylalanine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-, hydrazide (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-03-3 CAPLUS
CN Leucine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-, hydrazide (9CI) (CA INDEX NAME)




RN 112170-04-4 CAPLUS
CN L-Leucine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-, hydrazide (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-05-5 CAPLUS
CN Valine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-, hydrazide (9CI) (CA INDEX NAME)




RN 112170-06-6 CAPLUS
CN L-Valine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-, hydrazide (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-07-7 CAPLUS
CN Alanine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-, hydrazide (9CI) (CA INDEX NAME)




RN 112170-08-8 CAPLUS
CN L-Alanine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]-, hydrazide (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-18-0 CAPLUS
CN Serine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]- (9CI) (CA INDEX NAME)




RN 112170-19-1 CAPLUS
CN L-Serine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-26-0 CAPLUS
CN Glycine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]- (9CI) (CA INDEX NAME)



IT 112170-20-4P 112170-22-6P 112170-24-8P


112170-25-9P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and esterification and antimicrobial activity of)

RN 112170-20-4 CAPLUS
CN Leucine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]- (9CI) (CA INDEX NAME)




RN 112170-22-6 CAPLUS
CN Valine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)sulfonyl]- (9CI)

(CA INDEX NAME)




RN 112170-24-8 CAPLUS
CN Alanine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]- (9CI)
(CA INDEX NAME)




RN 112170-25-9 CAPLUS
CN L-Alanine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 112170-17-9P 112170-21-5P 112170-23-7P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and esterification and peptide coupling and antimicrobial activity of)
RN 112170-17-9 CAPLUS
CN L-Phenylalanine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-21-5 CAPLUS
CN L-Leucine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-23-7 CAPLUS
CN L-Valine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




IT 112170-09-9P 112170-10-3P 112170-11-3P
112170-12-4P 112170-13-5P 112170-14-6P
112170-15-7P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and hydrazinolysis and antimicrobial activity of)
RN 112170-09-9 CAPLUS
CN L-Phenylalanine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-10-2 CAPLUS
CN Leucine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]-, methyl ester (9CI) (CA INDEX NAME)




RN 112170-11-3 CAPLUS
CN L-Leucine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-12-4 CAPLUS
CN Valine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]-, methyl ester (9CI) (CA INDEX NAME)




RN 112170-13-5 CAPLUS
CN L-Valine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 112170-14-6 CAPLUS
CN Alanine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]-, methyl ester (9CI) (CA INDEX NAME)




RN 112170-15-7 CAPLUS
CN L-Alanine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 112170-16-6P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and peptide coupling and antimicrobial activity of)
RN 112170-16-8 CAPLUS
CN L-Tyrosine, N-[(1,2,3,4-tetrahydro-2,3-dioxo-6-quinoxaliny)lsulfonyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



L13 ANSWER 104 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1987:196459 CAPLUS
DOCUMENT NUMBER: 106:196459
TITLE: Preparation of quinoxalines as antidotes for acetanilide herbicides.
INVENTOR(S): Eicken, Karl; Spiegler, Wolfgang; Wuerzler, Bruno; Meyer, Norbert
PATENT ASSIGNEE(S): BASF A.-G., Fed. Rep. Ger.
SOURCE: Ger. Offen., 16 pp.
CODEN: GWXEXX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	----	-----	-----

DE 3533791 A1 19870326 DE 1985-3533791 19850921
 EP 216299 A1 19870401 EP 1986-112816 19860917
 R: CH, DE, FR, IT, LI
 JP 62072670 A2 19870403 JP 1986-219885 19860919
 SE 2003353 A6 19881101 SE 1986-2056 19860919
 PRIORITY APPL. INFO.: DE 1985-3533791 A 19850921
 OTHER SOURCE(S): CASREACT 106:196459
 GI




AB The title quinoxalines I [R1 = halo; R2 = H, alkyl, halo, CO2R4; R3 = H, CO2R4, R5R6C:NO; R4 = (un)substituted C1-6 alkyl, C3-6 alkenyl; R5, R6 = C1-6 alkyl, C3-6 alkenyl, cycloalkyl; R5R6 = (CH2)m; m = 1-4; n = 4-8] were prepared as antidotes for acetanilide herbicides II [A = C1-4 alkoxy, alkoxyalkyl, (un)substituted N-attached azolyl; Y = Br, Cl; R7 = H, C1-5 alkyl, alkoxy, R8, R9 = halo, R1] and their salts. 18.5 O
 6-chloro-2,3-quinoxalinediol was refluxed 4 h with 70.5 g PCl5 and 95 mL POCl3 to give 16.7 g I (R1 = R2 = Cl, R3 = 6-Cl, n = 1) (III). In tests with rice application of a mixture of III and II (A = 3,5-dimethylpyrazol-1-yl, Y = Cl, R7 = 2-Cl, R8 = 6-Cl, R9 = H) at 0.125 kg/ha and 0.025 kg/ha, resp., there was 98% kill of *Echinochloa crus-galli* with no damage to the rice plants.


IT 26773-13-7P 26773-25-1P 26773-32-0P
 26921-20-0P 108229-81-8P 108229-83-0P
 108229-84-1P 108229-85-2P 108229-86-3P
 108229-87-4P 108229-88-5P 108229-89-6P
 108229-90-9P 108229-92-1P 108229-94-3P
 108229-95-4P 108229-96-5P 108230-01-9P
 108230-02-0P 108230-03-1P 108258-55-5P
 108258-56-6P

RL: SPN (Synthetic preparation); PRSP (Preparation)
 (preparation of, as herbicide antidote for acetanilides)


RN 26773-13-7 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-hydroxyethyl)- (8CI, 9CI) (CA INDEX NAME)




RN 26773-25-1 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-ethyl- (8CI, 9CI) (CA INDEX NAME)




RN 26773-32-0 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-chloroethyl)- (8CI, 9CI) (CA INDEX NAME)




RN 26921-20-0 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-2-propenyl- (9CI) (CA INDEX NAME)




RN 108229-81-8 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, phenylmethyl ester (9CI) (CA INDEX NAME)




RN 108229-83-0 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, butyl ester (9CI) (CA INDEX NAME)




RN 108229-84-1 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, 2-propenyl ester (9CI) (CA INDEX NAME)




RN 108229-85-2 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, 2-ethoxyethyl ester (9CI) (CA INDEX NAME)




RN 108229-86-3 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, 2-phenylethyl ester (9CI) (CA INDEX NAME)




RN 108229-87-4 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, (4-methylphenyl)methyl ester (9CI) (CA INDEX NAME)




RN 108229-88-5 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, (2-chlorophenyl)methyl ester (9CI) (CA INDEX NAME)




RN 108229-89-6 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, [4-(1,1-dimethylethyl)phenyl]methyl ester (9CI) (CA INDEX NAME)




RN 108229-90-9 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, 2-chloroethyl ester (9CI) (CA INDEX NAME)



RN 108229-92-1 CAPLUS
 CN 6-Quinoxalinecarboxamide, N-butyl-2,3-dichloro- (9CI) (CA INDEX NAME)




RN 108229-94-3 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(phenylmethyl)- (9CI) (CA INDEX NAME)




RN 108229-95-4 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-phenylethyl)- (9CI) (CA INDEX NAME)


NAME)




RN 108229-96-5 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(1-phenylethyl)- (9CI) (CA INDEX NAME)




RN 108230-01-9 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, 1-phenylethyl ester (9CI) (CA INDEX NAME)




RN 108230-02-0 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, 2-furanylmethyl ester (9CI) (CA INDEX NAME)




RN 108230-03-1 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-furanylmethyl)- (9CI) (CA INDEX NAME)



RN 108258-55-5 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, (4-chlorophenyl)methyl ester (9CI) (CA INDEX NAME)




RN 108258-56-6 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2,3-dichloro-, (4-methoxyphenyl)methyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 105 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1987:50248 CAPLUS
DOCUMENT NUMBER: 106:50248
TITLE: Benzo[a]phenazine derivatives, with antitumor activity, and a process for their preparation
INVENTOR(S): Migita, Yoshihiro; Sguchi, Tadashi; Kumasawa, Yukinari; Nakagami, Joshi; Amano, Takehiro; Sota, Kaoru; Sakakibara, Jinsaku
PATENT ASSIGNER(S): Taiho Pharmaceutical Co., Ltd., Japan
SOURCE: Bur. Pat. Appl., 30 pp.
CODEN: SPXXDW
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 196910	A2	19861008	EP 1986-302395	19860401
EP 196910	A3	19870902		
EP 196910	B1	19910102		
R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE				
US 4686292	A	19870811	US 1986-838153	19860310
CA 1248106	A1	19880103	CA 1986-504378	19860318
JP 62000072	A2	19870106	JP 1986-64410	19860320
JP 05013149	B4	19920219		
ZA 8602183	A	19861126	ZA 1986-2183	19860324
ES 553500	A1	19870616	ES 1986-553500	19860326
AT 59642	E	19910115	AT 1986-302395	19860401
PRIORITY APPLN. INFO.:			JP 1985-65099	A 19850329
			EP 1986-302395	A 19860401
OTHER SOURCE(S):			CASREACT 106:50248; MARPAT 106:50248	


QI




AB Benzo[a]phenazine deriva. I (R1 = H, halo, Me, OH, alkoxy; R2 = CO2R5, CONR6R7; R3, R4 = H, alkyl; R5 = H, alkyl, cycloalkyl, Ph, PhCH2; R6, R7 = H, alkyl; NR6R7 = pyrrolidino, piperidino; n = 2, 3) are prepared as antitumor agents. A solution of ester II (R1 = OMe, R2 = CO2Me) in C6H6 was treated with H2NCH2CH2NMe2 and the mixture refluxed for 2 h to give I (R1 = OMe, R2 = CO2Me, R3 = R4 = Me, n = 2) (III). At 50 mg/kg/day i.p. for 5 days in mice transplanted with P388 leukemia cells, III increased survival time 5.88-fold, vs. 1.76-fold (maximum) for 5-FU at 25 mg/kg/day.

IT 106224-71-9P 106224-72-9P 106224-74-2P
106224-75-3P 106224-76-4P 106224-78-6P
106224-81-1P 106224-87-7P 106224-90-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


(preparation and amidation of)
RN 106224-71-9 CAPLUS
CN Benzo[a]phenazine-6,9-dicarboxylic acid, 10-ethoxy-5-hydroxy-, diethyl ester (9CI) (CA INDEX NAME)




RN 106224-72-0 CAPLUS
CN Benzo[a]phenazine-6,9-dicarboxylic acid, 10-ethoxy-5-hydroxy-, 9-butyl 6-ethyl ester (9CI) (CA INDEX NAME)




RN 106224-74-2 CAPLUS
CN Benzo[a]phenazine-6,9-dicarboxylic acid, 5-hydroxy-10-propoxy-, diethyl ester (9CI) (CA INDEX NAME)




RN 106224-75-3 CAPLUS
CN Benzo[a]phenazine-6,9-dicarboxylic acid, 6-hydroxy-10-propoxy-, 6-ethyl 9-propyl ester (9CI) (CA INDEX NAME)



RN 106224-76-4 CAPLUS
CN Benzo[a]phenazine-6,9-dicarboxylic acid, 5-hydroxy-10-propoxy-, 9-butyl 6-ethyl ester (9CI) (CA INDEX NAME)




RN 106224-78-6 CAPLUS
CN Benzo[a]phenazine-6,9-dicarboxylic acid, 10-butoxy-5-hydroxy-, 9-butyl 6-ethyl ester (9CI) (CA INDEX NAME)




RN 106224-81-1 CAPLUS
CN Benzo[a]phenazine-6,9-dicarboxylic acid, 5-hydroxy-, diethyl ester (9CI)


(CA INDEX NAME)




RN 106224-87-7 CAPLUS
CN Benzo[a]phenazine-6-carboxylic acid, 9-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, ethyl ester (9CI) (CA INDEX NAME)




RN 106224-90-2 CAPLUS
CN Benzo[a]phenazine-6,9-dicarboxylic acid, 5-hydroxy-10-methoxy-, 9-(2,2-dimethylpropyl) 6-ethyl ester (9CI) (CA INDEX NAME)




IT 106224-68-4P 106224-69-5P 106224-91-3P
106224-94-6P 106224-96-8P 106225-00-7P
106225-01-8P 106225-03-0P 106225-04-1P
106225-05-2P 106225-07-4P 106225-11-0P
106225-12-1P 106225-13-2P 106225-14-3P
106225-15-4P 106225-17-6P 106225-19-8P
106225-20-1P 106225-21-2P 106225-22-3P
106225-23-4P 106225-25-6P 106225-26-7P
RL: BAC (Biological activity or effector); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USRS (Uses)
(preparation of, as antitumor agent)
RN 106224-68-4 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, butyl ester (9CI) (CA INDEX NAME)




RN 106224-69-5 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-, 3-methylbutyl ester (9CI) (CA INDEX NAME)




RN 106224-91-3 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-, ethyl ester (9CI) (CA INDEX NAME)




RN 106224-94-6 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, 2,2-dimethylpropyl ester (9CI) (CA INDEX NAME)




RN 106224-96-8 CAPLUS
CN Benzo[a]phenazine-6,9-dicarboxamide, N6-[2-(dimethylamino)ethyl]-N9-ethyl-5-hydroxy-10-methoxy- (9CI) (CA INDEX NAME)




RN 106225-00-7 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-10-ethoxy-5-hydroxy-, ethyl ester (9CI) (CA INDEX NAME)




RN 106225-01-8 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-10-ethoxy-5-hydroxy-, butyl ester (9CI) (CA INDEX NAME)




RN 106225-03-0 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-propoxy-, ethyl ester (9CI) (CA INDEX NAME)




RN 106225-04-1 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-propoxy-, propyl ester (9CI) (CA INDEX NAME)




RN 106225-05-2 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-propoxy-, butyl ester (9CI) (CA INDEX NAME)




RN 106225-07-4 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 10-butoxy-6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-, butyl ester (9CI) (CA INDEX NAME)




RN 106225-11-0 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, ethyl ester (9CI) (CA INDEX NAME)




RN 106225-12-1 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, propyl ester (9CI) (CA INDEX NAME)




RN 106225-13-2 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, 1-methylethyl ester (9CI) (CA INDEX NAME)




RN 106225-14-3 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, 2-methylpropyl ester (9CI) (CA INDEX NAME)




RN 106225-15-4 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, pentyl ester (9CI) (CA INDEX NAME)



RN 106225-17-6 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, hexyl ester (9CI) (CA INDEX NAME)




RN 106225-19-8 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, heptyl ester (9CI) (CA INDEX NAME)




RN 106225-20-1 CAPLUS


CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, octyl ester (9CI) (CA INDEX NAME)




RN 106225-21-2 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, decyl ester (9CI) (CA INDEX NAME)




RN 106225-22-3 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, pentadecyl ester (9CI) (CA INDEX NAME)




RN 106225-23-4 CAPLUS
CN Benzo[a]phenazine-9-carboxylic acid, 6-[[[2-(dimethylamino)ethyl]amino]carbonyl]-5-hydroxy-10-methoxy-, phenylmethyl ester (9CI) (CA INDEX NAME)



RN 106225-25-6 CAPLUS
CN Benzo[a]phenazine-6,9-dicarboxamide, N6-[2-(dimethylamino)ethyl]-5-hydroxy-10-methoxy-N9-propyl- (9CI) (CA INDEX NAME)




RN 106225-26-7 CAPLUS
CN Benzo[a]phenazine-6,9-dicarboxamide, N9-butyl-N6-[2-(dimethylamino)ethyl]-5-hydroxy-10-methoxy- (9CI) (CA INDEX NAME)



L13 ANSWER 106 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1987:19959 CAPLUS
DOCUMENT NUMBER: 106:19959
TITLE: Lithium salts of fiber-reactive anionic dyes
INVENTOR(S): Meininger, Fritz; Schlaefer, Ludwig
PATENT ASSIGNER(S): Hoechst A.-G., Fed. Rep. Ger.
SOURCE: Ger. Offen., 25 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3443305	A1	19860528	DE 1984-3443305	19841128
EP 183142	A2	19860604	EP 1985-114511	19851115


EP 183142 A3 19890329
 R: CH, DE, FR, GB, IT, LI
 IN 164645 A 19890429
 US 4707545 A 19871117
 JP 6113074 A2 19860518
 BR 8505929 A 19860819
 CA 1253858 A1 19890509
 PRIORITY APPLN. INFO.:
 OTHER SOURCE(S): MARPAT 106:19959
 GI



AB Li salts of anionic dyes are prepared by neutralizing an aqueous solution or suspension of an acidic dye with a Ca salt forming a low solubility Ca dye salt, which is treated with Li2SO4 or LiHSO4 to form the corresponding Li salt. Thus, a Na salt of an acidic monazo dye was reacted with CaCl2 to form a Ca salt containing filter cake, which was treated with a solution of Li2SO4 forming I, which was storage-stable at 50° for several weeks. The solution was spray-dried to form an electrolyte-containing fine dye powder.

IT 104601-66-3
 RL: PROC (Process)
 (cation exchange of, with calcium chloride)


RN 104601-66-3 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxaliny)carbonyl]methylamino]methyl]-3-sulfonylphenyl]amino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)



IT 106046-41-7


RL: PROC (Process)
 (cation exchange of, with lithium oxalate)

RN 106046-41-7 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxaliny)carbonyl]methylamino]methyl]-3-sulfonylphenyl]amino]-9,10-dihydro-9,10-dioxo-, calcium salt (9CI) (CA INDEX NAME)



IT 106027-72-9P
 RL: IMP (Industrial manufacture); PREP (Preparation)
 (preparation of, as storage-stable spray-dryable reactive dye composition)

RN 106027-72-9 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxaliny)carbonyl]methylamino]methyl]-3-sulfonylphenyl]amino]-9,10-dihydro-9,10-dioxo-, dilithium salt (9CI) (CA INDEX NAME)



L13 ANSWER 107 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STM

ACCESSION NUMBER: 1986:610400 CAPLUS
 DOCUMENT NUMBER: 105:210400
 TITLE: Storage-stable dye solutions
 INVENTOR(S): Wolff, Joachim; Wolf, Karlheinz; Marschner, Werner
 PATENT ASSIGNEE(S): Bayer A.-G., Fed. Rep. Ger.
 SOURCE: Eur. Pat. Appl., 28 pp.
 CODEN: EPKXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 167952	A2	19860115	EP 1985-108117	19850629
EP 167952	A3	19890322		
EP 167952	B1	19910724		
R: CH, DE, FR, GB, LI				
DE 3504964	A1	19860123	DE 1985-3504964	19850213
US 4685933	A	19870811	US 1985-748267	19850624
JP 6103639	A2	19860221	JP 1985-149387	19850709
JP 05023306	B4	19930402		

PRIORITY APPLN. INFO.:
 DE 1984-3425813 A 19840713
 DE 1985-3504964 A 19850213

AB Concentrated aqueous solns. of organic dyes and water-solubilizing agents contain cyanamides N.tplbond.CNH2 or RC(NH)NH2 (I; R = NHC.tplbond.N, NHCONH2, NHCONHMe). These aqueous solns. may contain an anionic reactive dye 7-35, a solubility-increasing water-miscible organic compound and/or hydrotrope and/or dispersing agent 0-30, N.tplbond.CNH2 or I 0.1-15, an inorg. salt 0-10, and a buffer 0-5%. Thus, to 997.5 g aqueous suspension containing a sulfonated (phenylazo)naphthalene dye with a chlorodifluoropyrimidine group 11, inorg. salt mixture 1, and H3BO3 0.5%, 2.5 g dicyandiamide was added with stirring. A storage-stable aqueous solution was obtained showing no dye hydrolysis after 4 wk at 40°. This solution dyed rayon in a red-yellow tone.

IT 104993-68-2
 RL: USES (Uses)
 (concentrated aqueous solns. of, stabilizers against hydrolysis for)

RN 104993-68-2 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxaliny)carbonyl]methylamino]methyl]-3-sulfonylphenyl]amino]-9,10-dihydro-9,10-dioxo-, monolithium monosodium salt (9CI) (CA INDEX NAME)



D1-SO3H

Li

Na

L13 ANSWER 108 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STM
 ACCESSION NUMBER: 1986:610360 CAPLUS
 DOCUMENT NUMBER: 105:210360
 TITLE: Aqueous reactive dye solutions
 INVENTOR(S): Wolff, Joachim; Wolf, Karlheinz; Seipt, Guenter
 PATENT ASSIGNEE(S): Bayer A.-G., Fed. Rep. Ger.
 SOURCE: Ger. Offen., 11 pp.
 CODEN: GNMXXB
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3424145	A1	19860109	DE 1984-3424145	19840630

PRIORITY APPLN. INFO.:
 DE 1984-3424145 19840630
 AB Aqueous solns. of reactive dyes containing water-solubilizing groups are stabilized by addition of 0.1-25% R1(CR3R4)C.tplbond.CCR5R6R2 (I; R1, R2, R4, R6 = H, Cl-6 alkyl; R3, R5 = OH, (OCH2CHR7)OH; R7 = H, Me; m = 0, 1; n = 1-10). These solns. also contain a buffer 0-5, the reactive dye


PAGE 1-A

PAGE 2-A

10-35, an inorg. salt 0-10, and a solubility-increasing water-miscible organic compound and/or hydrotrope and/or dispersing agent 0-30%. Thus, an aqueous solution was formulated containing a sulfated anthraquinone dye with a dichloroquinoxaline reactive group 20, ϵ -caprolactam 10, MENCONHMe 10, and a mixture of inorg. salts (NaCl, LiCl, Na2SO4, and Li2SO4) 3%. This solution (200 g) was mixed with 50 g urea, and 550 g water, giving solution A; filtrn. 15 s after addition of 200 g 20% Na2CO3 showed precipitation. If to 800 g of solution A, 4 g I (m = 1, R1 = R2 = iso-Bu, R3 = R5 = Me, R4 = R6 = OH) in a mixture of ethylene glycol and polyethylene glycol alkylphenyl ether was added, followed by 200 g 20% Na2CO3 solution, no precipitation was observed.

IT 88103-23-5D, lithium and sodium salts
RL: USES (Uses)
(stabilizers for aqueous solns. of, alkynediols as)

RN 88103-23-5 CAPLUS
CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxalyl)carbonyl]methylamino]methyl]sulfonyl]amino]-9,10-dihydro-9,10-dioxo- (9CI) (CA INDEX NAME)




D1-SO3H

L13 ANSWER 109 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1984:630474 CAPLUS
DOCUMENT NUMBER: 101:230474
TITLE: Derivatives of 1,3-benzodioxole, 52. Preparation and reactions of 1,3-dioxolo[4,5-b]phenazines
Dallacker, Franz; Wagner, Alfred
Abt. Chem. Med., Tech. Hochschule Aachen, Aachen, D-5100, Fed. Rep. Ger.
Zeitschrift fuer Naturforschung, Teil B: Anorganische Chemie, Organische Chemie (1984), 39B(7), 936-49
CODEN: ZNBAD2; ISSN: 0340-5087

PAGE 1-A


PAGE 2-A

DOCUMENT TYPE: Journal
LANGUAGE: German
OTHER SOURCE(S): CASREACT 101:230474
OI




AB The 1,3-dioxolo[4,5-b]phenazines I (R, R7 = H, Me, CO2Me; R1R2 = H2, OCH2O; R3 = H, Me, CO2Me, OMe; R4 = H, CO2Me; R5 = H, CO2Et, Me, OMe; R6 = H, OMe; R5R6 = OCH2O) were formed by FeCl3 oxidation of the corresponding 2,2'-diaminodiphenylamines which were obtained by condensing the 6-chloro-8-cyclohexyl-11-R-5,8-dihydronaphtho[1,2-b]phenazin-5-ones
Schelz, D.; Rotsler, N.
Inst. Farbenchem., Univ. Basel, Basel, CH-4056, Switz.
Dyes and Pigments (1984), 5(1), 37-47
CODEN: DYPIDX; ISSN: 0143-7208
Journal
German

IT 93415-94-2P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)
RN 93415-94-2 CAPLUS
CN 1,3-Dioxolo[4,5-b]phenazine-7-carboxylic acid, ethyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 110 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1984:87233 CAPLUS
DOCUMENT NUMBER: 100:87233
TITLE: Ring closure of quinonimethane dyes and merocyanine analogs. Part 7. Synthesis and properties of 6-chloro-8-cyclohexyl-11-R-5,8-dihydronaphtho[1,2-b]phenazin-5-ones
Schelz, D.; Rotsler, N.
Inst. Farbenchem., Univ. Basel, Basel, CH-4056, Switz.
Dyes and Pigments (1984), 5(1), 37-47
CODEN: DYPIDX; ISSN: 0143-7208
Journal
German

AB The 1,3-dioxolo[4,5-b]phenazines I (R, R7 = H, Me, CO2Me; R1R2 = H2, OCH2O; R3 = H, Me, CO2Me, OMe; R4 = H, CO2Me; R5 = H, CO2Et, Me, OMe; R6 = H, OMe; R5R6 = OCH2O) were formed by FeCl3 oxidation of the corresponding 2,2'-diaminodiphenylamines which were obtained by condensing the 6-chloro-8-cyclohexyl-11-R-5,8-dihydronaphtho[1,2-b]phenazin-5-ones
Schelz, D.; Rotsler, N.
Inst. Farbenchem., Univ. Basel, Basel, CH-4056, Switz.
Dyes and Pigments (1984), 5(1), 37-47
CODEN: DYPIDX; ISSN: 0143-7208
Journal
German




AB Naphthophenazines I (R = OMe, Me, NHAc, H, F, Cl, Br, CO2Et, CF3) were obtained by reaction of 6-substituted 1-cyclohexyl-2,3-dimethylquinoxalium perchlorates with 2,3-dichloro-1,4-naphthoquinone [117-80-6]. The influence of R on general properties, lightfastness and on UV-visible and 1H-NMR spectra is discussed.

IT 87815-93-8
RL: RCT (Reactant); RACT (Reactant or reagent)
(reaction of, with dichloronaphthoquinone, ring formation in)

RN 87815-93-8 CAPLUS
CN Quinoxalium, 1-cyclohexyl-6-(ethoxycarbonyl)-2,3-dimethyl-, perchlorate (9CI) (CA INDEX NAME)

CM 1
CRN 87815-92-7
CMF C19 H25 N2 O2



CM 2
CRN 14797-73-0
CMF C1 O4




L13 ANSWER 111 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1984:35784 CAPLUS
DOCUMENT NUMBER: 100:35784
TITLE: Concentrated liquid compositions of cold-dyeing

INVENTOR(S): fiber-reactive dyes
Hoguet, Robert G.; Kalz, Dietmar; Thomas, Thomas J.;
Whetzel, Henry T.; Wolff, Joachim; Nonn, Konrad;
Wolf, Karlheinz
Bayer A.G., Fed. Rep. Ger.; Mobay Chemical Corp.
Eur. Pat. Appl., 34 pp.
CODEN: SPXKDW
Patent
German

PATENT ASSIGNEE(S):
SOURCE: Bayer A.G., Fed. Rep. Ger.; Mobay Chemical Corp.
Eur. Pat. Appl., 34 pp.
CODEN: SPXKDW
Patent
German

DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PRIORITY INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 92119	A2	19831026	EP 1983-103418	19830408
EP 92119	A3	19841107		
US 4435181	A	19840306	US 1982-370426	19820421
JP 58187460	A2	19831101	JP 1983-65768	19830415
BR 8302071	A	19831227	BR 1983-2071	19830420
CA 1205253	A1	19860603	CA 1983-443149	19831213
PRIORITY APPL. INFO.:			US 1982-370426	A 19820421
OTHER SOURCE(S):		MARPAT 100:35784		
OI				




AB Storage-stable, aqueous cold-dyeing reactive dye compns. are prepared which contain 10-50 weight dye(s) with a fiber-reactive haloheterocyclic group and particle size <100 μ , sufficient anionic dispersant or polymeric N-vinyl lactam dispersant to prevent agglomeration or settling out of dye particles, and sufficient electrolyte to inhibit hydrolysis of the reactive group during temperature cycles ranging from 20° to 50°.

A typical composition, stable for 3 wk during temperature cycles of 16 h at 20° and 8 h at 50°, contained dye I [78246-64-7] 31.5, lignosulfonate dispersant 3.0, NaCl 15.0, KH2PO4 0.2, and H2O 50.0%.

IT 78246-64-7
RL: USES (Uses)
(reactive dye, concentrated aqueous compns. containing, storage-stable)


RN 78246-64-7 CAPLUS
CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxalyl)carbonyl]methylamino]methyl]sulfonyl]amino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)

D1-SO₃H

● 2 Na

L13 ANSWER 112 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1984:8505 CAPLUS
 DOCUMENT NUMBER: 100:8505
 TITLE: Reactive dye preparations
 INVENTOR(S): Wolff, Joachim; Karlheinz, Wolf; Hoernle, Reinhold;
 Ditzler, Reiner; Falkenberg, Kurt
 PATENT ASSIGNEE(S): Bayer A.-G., Fed. Rep. Ger.
 SOURCE: Eur. Pat. Appl., 32 pp.
 CODEN: EPXKDW
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 87703	A1	19830907	EP 1983-101616	19830221
EP 87703	B1	19850410		
R: CH, DE, FR, GB, IT, LI				
DE 3207534	A1	19830908	DE 1982-3207534	19820303
DE 3215933	A1	19831103	DE 1982-3215933	19820429
PRIORITY APPL. INFO.:				
OTHER SOURCE(S): MARPAT 100:8505				
OI				




II

AB Dyes containing an NH₂ or Cl-4-alkylamino group and 1-5 sulfo groups, and having salt content 58%, are treated with a chloro- and/or fluoro-substituted heterocyclic compound in aqueous medium in the presence of an acid-binding agent and, optionally, a solubilizer, and the resultant liquid is optionally buffered or dried and milled. The compns. are used to prepare dyebaths and printing pastes. For example, diazotization of 0.3 mol 2,5,1-H₂N(H₂NCH₂)ClOHSO₃H, coupling with 0.3 mol 1,8,3,6-B₂NH(HO)ClO₄(SO₃H)₂, salting out with 15 weight% NH₄HCO₃, washing, and drying gave a dye (I) [70817-82-2] containing 3% salt. A soln of I in 700 mL H₂O containing 3.25 mol ε-caprolactam was treated with 0.3 mol 5-chloro-2,4,6-trifluoropyrimidine [697-83-6] in the presence of 0.32 mol Li₂CO₃ and finally 0.13 mol CaO, and adjusted to pH 7 with 0.6 weight% phosphate buffer to give a storage-stable solution containing 19 weight% II salt.

IT 88103-23-5P
 RL: PREP (Preparation)
 (manufacture of, as powder composition with improved water solubility)
 RN 88103-23-5 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxaliny)carbonyl]methylamino]methyl]sulfonyl]amino]-9,10-dihydro-9,10-dioxo- (9CI) (CA INDEX NAME)

D1-SO₃H


IT 87748-64-9P
 RL: PREP (Preparation)
 (storage-stable solution of, manufacture of)
 RN 87748-64-9 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxaliny)carbonyl]methylamino]methyl]sulfonyl]amino]-9,10-dihydro-9,10-dioxo-, dilithium salt (9CI) (CA INDEX NAME)

D1-SO₃H

● 2 Li

L13 ANSWER 113 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1984:8427 CAPLUS
 DOCUMENT NUMBER: 100:8427
 TITLE: Preparations of water-soluble organic dyes
 INVENTOR(S): Wolff, Joachim; Wolf, Karlheinz; Hoernle, Reinhold
 PATENT ASSIGNEE(S): Bayer A.-G., Fed. Rep. Ger.
 SOURCE: Ger. Offen., 23 pp.
 CODEN: GWXXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3207533	A1	19830908	DE 1982-3207533	19820303
EP 87705	A1	19830907	EP 1983-101619	19830221
EP 87705	B1	19850904		
R: CH, DE, FR, GB, LI				
JP 58162667	A2	19830927	JP 1983-31078	19830228
PRIORITY APPL. INFO.:				
OI				



I

AB Preps. containing water-soluble anionic or cationic dyes and di-Cl-4-alkyl sulfones are stable in storage. Thus, a mixture of reactive dye I [88112-50-9] 16.5, di-Me sulfone (II) [67-71-0] 10, phosphate buffer (for pH 7) 0.5, and H₂O 73 parts was stirred and filtered to give a solution. Storage of this solution for 4 wk at 50° resulted in only 4% hydrolysis of the dye, compared with 25% when II was replaced by tetramethylene sulfone.

IT 78246-64-7
 RL: USES (Uses)
 (aqueous solns. of, containing di-Me sulfone, storage-stable)
 RN 78246-64-7 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxaliny)carbonyl]methylamino]methyl]sulfonyl]amino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)



D1-SO₃H


PAGE 2-A

● 2 Na

L13 ANSWER 114 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1983:596640 CAPLUS
DOCUMENT NUMBER: 99:196640
TITLE: Concentrated liquid reactive dye preparations
INVENTOR(S): Wolff, Joachim; Wolf, Karlheinz; Ditzler, Reiner;
Hoernle, Reinhold
PATENT ASSIGNEE(S): Bayer A.-G., Fed. Rep. Ger.
SOURCE: Ger. Offen., 19 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3207534	A1	19830908	DE 1982-3207534	19820303
EP 87703	A1	19830907	EP 1983-101616	19820221
EP 87703	B1	19850410		
R: CH, DE, FR, GB, IT, LI				
JP 58162666	A2	19830927	JP 1983-31079	19830228
JP 02009629	B4	19900302		
BR 8301017	A	19831122	BR 1983-1017	19830302
PRIORITY APPL. INFO.:			DE 1982-3207534	A 19820303
			DE 1982-3215933	A 19820429


OI



AB Dyes of formula (M03S)nQNR (Q = dye residue; M = H, NH₄, Li, Na, K; n = 1-5; R = H, Cl-4 alkyl), with salt content 50 weight%, are treated in aqueous media (optionally containing a solubilizing agent) with a halo heterocyclic compound in the presence of an acid acceptor and then optionally buffered to give a concentrated solution of reactive dye (M03S)nQNR2.

(R1 = fiber-reactive heterocyclic group; M1 = NH₄, Li, Na, K; n and Q as defined above). For example, diazotization of 0.3 mol 4,3-H₂N(HO3S)C₆H₃NHAc [96-78-6], coupling with 0.3 mol 1-(4-sulfophenyl)-5-pyrazolone-3-carboxylic acid [118-47-8], deacetylation, dissoln. of the presake by LiOH in 450 mL H₂O, treatment with 0.33 mol 2,3-dichloro-6-quinoxalinecarboxyl chloride [1919-43-3]

OTHER SOURCE(S): CASREACT 99:196628
OI




AB The title compds. (I; R = cyclohexyl, Me; R1 = H₂N, MeO, Me, AcNH, H, halogen, EtO₂C, F₃C, MeSO₂, O₂N), useful as precursors for naphthophenazinone dyes were synthesized starting with 2-nitrohalobenzenes. The preferred method of condensing the diamine intermediate (II) with 2,3-butanedione [431-03-8] in mixts. of HOAc and HClO₄ was not successful whenever R1 was a strongly electron-withdrawing substituent. But in those cases R1 stabilized the corresponding enamines III, which could be obtained very easily.

IT 87815-93-8P
RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (preparation and NMR spectrum of)

RN 87815-93-8 CAPLUS

CN Quinoxalium, 1-cyclohexyl-6-(ethoxycarbonyl)-2,3-dimethyl-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 87815-92-7
CMF C19 H25 N2 O2

CM 2

CRN 14797-73-0
CMF C1 O4


while adding Li₂CO₃ to neutralize HCl, filtration, and buffering to pH 7 with 0.6% phosphate gave a storage-stable liquid crystalline preparation containing 22 weight% I [87730-51-6].

IT 87748-64-9
RL: USSES (Uses)
(dye, manufacture of concentrated storage-stable solution of)

RN 87748-64-9 CAPLUS

CN 2-Anthracenesulfonic acid, 1-amino-4-[[4-[[[(2,3-dichloro-6-quinoxalyl)carbonyl]methylamino]methyl]sulfonyl]amino]-9,10-dihydro-9,10-dioxo-, dilithium salt (9CI) (CA INDEX NAME)

PAGE 1-A

D1-SO₃H

● 2 Li

PAGE 2-A

L13 ANSWER 115 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1983:596628 CAPLUS
DOCUMENT NUMBER: 99:196628
TITLE: Synthesis of 1-aryl- and 1-alkyl-2,3-dimethylquinoxalium perchlorates. Part 3. Synthesis of 1,2,3-trimethyl-6-X- and 1-cyclohexyl-2,3-dimethyl-6-X-quinoxalium perchlorates
Schelz, D.; Rotzler, N.
Inst. Farbenchem., Univ. Basel, Basel, CH-4056, Switz.
Dyes and Pigments (1983), 4(4), 305-20
CODEN: DYPIDK; ISSN: 0143-7208
DOCUMENT TYPE: Journal
LANGUAGE: German



L13 ANSWER 116 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1983:181108 CAPLUS
DOCUMENT NUMBER: 98:181108
TITLE: Crossconjugated cyanines and merocyanines, obtained from salts of 1-substituted 2,3-dimethylquinoxalines. Part 2. Oxidative transformation of color bases
Schelz, Dieter
Inst. Farbenchem., Univ. Basel, Basel, 4056, Switz.
Helvetica Chimica Acta (1982), 65(5), 1607-16
CODEN: HCACAV; ISSN: 0018-019X
DOCUMENT TYPE: Journal
LANGUAGE: German
OI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Products (I; R = Ph, 4-ClC₆H₄, Me; R1 = NO₂, CF₃, Br, H, OMe, etc.) of the readily oxidized II (X = N; R and R1 as defined) and II (X = NH+; R and R1 as defined) are sensitive to solvolysis, especially when R and R1 are electron-withdrawing substituents. In some cases, I could be identified as the oxidation products of III. The oxidation of III (R = Me, R1 = 5,6-benzo.

A = ClO₄ [52736-76-2] by alkaline K₂Fe(CN)₆ leading to IV [84268-37-1] was compared with the voltammetric oxidation of V [68797-83-1] and related to the capto, dative-stabilized radicals proposed by H.G. Viehe et al. (1979). IR-NMR spectra of I were discussed with regard to E-Z isomers.

IT 84267-32-3P
RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (preparation and NMR spectrum of)


RN 84267-32-3 CAPLUS

CN Quinoxalium, 2,2'-(1,3-ethenediyl)bis[6-(ethoxycarbonyl)-3-methyl-1-phenyl-, (E)-, diperchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 84267-31-2
CMF C38 H34 N4 O4

Double bond geometry as shown.



CM 2

CRN 14797-73-0
CMF C1 O4



L13 ANSWER 117 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER:
DOCUMENT NUMBER:

1983:55539 CAPLUS
98:55539
TITLE:
Crossconjugated cyanines and merocyanines, obtained from salts of 1-substituted 2,3-dimethylquinoxalines. Part 2. Oxidative transformation of color bases
AUTHOR(S):
CORPORATE SOURCE:
SOURCE:
DOCUMENT TYPE:
LANGUAGE:
OTHER SOURCE(S):
GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


AB Products (I; R = Ph, 4-ClC6H4, Me; R1 = NO2, CF3, Br, H, OMe, etc.) of the readily oxidized II (X = N; R and R1 as defined) and II (X = NH; R and R1 as defined) are sensitive to solvolysis, especially when R and R1 are electron-withdrawing substituents. In some cases, I could be identified as the oxidation products of III. The oxidation of III (R = Me, R1 = 5,6-benzo, A = ClO4) [52736-76-2] by alkaline K3Fe(CN)6 leading to IV [84268-37-1] was compared with the voltammetric oxidation of V [68797-93-1] and related to the captodative-stabilized radicals proposed by H. G. Viche et al. (1979). 1H-NMR spectra of I were discussed with regard to E-Z isomers.
IT 84267-32-3P
RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (preparation and NMR spectrum of)
RN 84267-32-3 CAPLUS
CN Quinoxalium, 2,2'-(1,2-ethenediyl)bis(6-(ethoxycarbonyl)-3-methyl-1-

phenyl-, (S)-, diperchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 84267-31-2
CMF C38 H34 N4 O4

Double bond geometry as shown.



CM 2


CRN 14797-73-0
CMF C1 O4



L13 ANSWER 118 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER:
DOCUMENT NUMBER:


1982:194867 CAPLUS
96:194867
TITLE:
Microbial mutagenicity and toxicity of newly synthesized heterocyclic N-oxides
AUTHOR(S):
CORPORATE SOURCE:
SOURCE:
DOCUMENT TYPE:
LANGUAGE:
GI



AB Newly synthesized heterocyclic N-oxides were tested for their mutagenicity

using the Ames test. DX1 (I) [81485-18-9] was potentially mutagenic in Salmonella typhimurium TA 100 and 98 with and without the S-9 mixture WO 25 [81485-17-8] And WO 20 [81485-16-7], being structurally related to I, did not show any genetic change in the strains used. The antibiotic activity of these chems. was also tested using gram-neg. and gram-pos. bacteria. I had more killing effect in gram-pos. bacteria than WO 25 and WO 20.


IT 81485-16-7
RL: ADV (Adverse effect, including toxicity); BIOL (Biological study) (mutagenicity and toxicity of)
RN 81485-16-7 CAPLUS
CN D-Phenylalanine, N-[[[3-[[[(1S,2R)-2-hydroxy-1-methyl-2-phenylethyl]methylamino]carbonyl]-2-methyl-1,4-dioxido-6-quinoxaliny]carbonyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 119 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER:
DOCUMENT NUMBER:

1982:87009 CAPLUS
96:87009
TITLE:
Cross-conjugated cyanines and merocyanines, obtained from salts of 1-substituted 2,3-dimethylquinoxalines. Part 1. Isolation of the dye bases from spontaneous transformation or oxidation of the reactants with copper(II) acetate or silver oxide
AUTHOR(S):
CORPORATE SOURCE:
SOURCE:
DOCUMENT TYPE:
LANGUAGE:
OTHER SOURCE(S):
GI




AB Quaternary salts I (R = Me, Ph, p-ClC6H4; R1 = H, electron acceptor or donor; R2 = Me, Ph, p-ClC6H4), in some cases in the presence of the corresponding II, undergo spontaneous conversion to III (all groups as defined for I) when dissolved in DMSO or DMF. Yields are 24-47%. Higher

yields (up to 66%) are obtained by oxidation of I, II, or I-II mixts. with Cu(OAc)2 or Ag2O. Viable and 1H-NMR spectra data for the dyes are given, and their structural relationship to S. Huenig's (1980) two-step redox systems is discussed.

IT 68765-65-1
RL: RCT (Reactant); RACT (Reactant or reagent) (oxidative dimerization of)
RN 68765-65-1 CAPLUS
CN Quinoxalium, 6-(ethoxycarbonyl)-2,3-dimethyl-1-phenyl-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 68765-64-0
CMF C19 H19 N2 O2



CM 2



CRN 14797-73-0
CMF C1 O4




L13 ANSWER 120 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER:
DOCUMENT NUMBER:


1981:497719 CAPLUS
95:97719
TITLE:
The synthesis of "stretched-out" analogs of lumazine, 6,7-dimethylumazine and 2-amino-5,6,7,8-tetrahydro-6,7-dimethyl-4-pteridinone
AUTHOR(S):
CORPORATE SOURCE:
SOURCE:
DOCUMENT TYPE:
LANGUAGE:
GI



AB Treating 2,4,5-(H2N)3C6H2CO2Et with glyoxal and MeCOOMe gave I (R = H, Me), which were treated with urea to give pyrazinoquinazolinone II. III was similarly prepared
 IT 78795-09-2P 78795-10-5P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation and cyclocondensation of, with urea)
 RN 78795-09-2 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 7-amino-, ethyl ester (9CI) (CA INDEX NAME)



RN 78795-10-5 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 7-amino-2,3-dimethyl-, ethyl ester (9CI) (CA INDEX NAME)





L13 ANSWER 121 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1981:463663 CAPLUS
 DOCUMENT NUMBER: 95:63663
 TITLE: 2,3-Dichloroquinoxaline-6-carboxamide derivatives
 Gleinig, Harald; Lehre, Juergen; Jovicic, Dorde;
 INVENTOR(S): Schubert, Klaus; Oomm, Walter; Goessling, Claus
 PATENT ASSIGNEE(S): Bayer A.-G., Fed. Rep. Ger.
 SOURCE: Ger. Offen., 12 pp.
 CODEN: GWXXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2942364	A1	19810423	DE 1979-2942364	19791019
DE 2942364	C2	19861120	DE 1979-2942364	A 19791019


PRIORITY APPLN. INFO.:
 GI



AB Title compds. (I; Z = organic radical; R = H, Cl-4 alkyl; n = 1-4) are prepared in high yield by treating Z(NHR)n in aqueous medium at 5-20° and pH 3.5-5 with molten 2,3-dichloroquinoxaline-6-carboxyl chloride [1919-43-3] at 110-180° which is introduced beneath the surface of the aqueous phase via a spray nozzle. The method is especially useful for preparing I in which


Z is a dye residue, e.g., azo, anthraquinone, or phthalocyanine. The preparation of II [78246-64-7] and several other fiber-reactive dyes is described.
 IT 78181-07-4P 78246-64-7P
 RL: IMP (Industrial manufacture); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)
 (dye, manufacture of)
 RN 78181-07-4 CAPLUS
 CN Cuprate(3-), [3-[[[4-[[[5-[[[2,3-dichloro-6-quinoxaliny]carbonyl]methylamino]methyl]-2-sulphophenyl]azo]-2-hydroxy-5-methylphenyl]azo]-4-hydroxy-2,7-naphthalenedisulfonato(5-)]-], trisodium (9CI) (CA INDEX NAME)

PAGE 1-A




● 3 Na⁺

PAGE 1-B



RN 78246-64-7 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[[[4-[[[2,3-dichloro-6-quinoxaliny]carbonyl]methylamino]methyl]sulphophenyl]amino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)

PAGE 1-A



D1- SO₃H

● 2 Na

PAGE 2-A


L13 ANSWER 122 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1981:425134 CAPLUS

DOCUMENT NUMBER: 95:25134
 TITLE: Quinoxalinyloxyphenoxymethane carboxylic acid derivatives and their use as herbicides
 INVENTOR(S): Serban, Alexander; Watson, Keith Geoffrey; Farquharson
 PATENT ASSIGNEE(S): ICI Australia Ltd., Australia
 SOURCE: Eur. Pat. Appl., 63 pp.
 CODEN: EPXKDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 23785	A2	19810211	EP 1980-302411	19800717
EP 23785	A3	19810429		
EP 23785	B1	19850403		
R: AT, BE, CH, DE, FR, GB, IT, NL, SE				
AU 540234	B2	19841108	AU 1980-59547	19790717
AU 8059547	A1	19810806		
CA 8003928	A	19810624	ZA 1980-3928	19800630
IL 60506	A1	19861231	IL 1980-60506	19800706
CA 1314549	A1	19930316	CA 1980-356027	19800711
HU 26554	O	19830928	HU 1980-1762	19800715
HU 186299	B	19850729		
DK 8003068	A	19810118	DK 1980-3068	19800716
DK 160426	B	19910311		
DK 160426	C	19910819		
BR 8004413	A	19810127	BR 1980-4413	19800716
ES 493431	A1	19810701	ES 1980-493431	19800716
CS 239908	B2	19860116	CS 1980-5044	19800716
SU 1261564	A1	19860930	SU 1980-2951003	19800716
JP 56039077	A2	19810414	JP 1980-96960	19800717
JP 06013489	B4	19940223		
AT 12495	E	19850415	AT 1980-302411	19800717
US 4655819	A	19870407	US 1981-334384	19811224
US 4803273	A	19890207	US 1986-939694	19861209
DK 8901684	A	19890407	DK 1989-1684	19890407
DK 168380	B1	19940321		
DK 8901685	A	19890407	DK 1989-1685	19890407
DK 162521	B	19911111		
DK 162521	C	19920330		

PRIORITY APPLN. INFO.:
 AU 1979-9617 A 19790717
 AU 1980-3093 A 19800411
 US 1980-164933 A2 19800701
 EP 1980-302411 A 19800717
 AU 1981-7201 A 19810112
 US 1981-334384 A3 19811224

OTHER SOURCE(S): CASREACT 95:25134
 GI




AB The title compds. I (X = optionally substituted OC6H4O, OC6H4S, SC6H4S; R = H, optionally substituted alkyl, acyl; R1 = H, optionally substituted alkyl; RR1 = alkylene; R2 = cyano, carbamoyl, optionally esterified CO2H, substituted Me; R3 = H, halogen, cyano, thiocyanato, optionally substituted NH2, aliphatic, OH, SH, CO2H, or CONH2; m, n = 0, 1; p = 0-2) were prepared. Thus, 2,6-dichloroquinoxaline was treated with 4-HOC6H4OCHMeCO2Me to give 70% I. At 1 kg/ha preemergence II gave 100% control of ryegrass and Japanese millet.

IT 78104-80-0P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 78104-80-0 CAPLUS


CN 6-Quinoxalinecarboxylic acid, 3-(4-(2-ethoxy-1-methyl-2-oxoethoxy)phenoxy)-, ethyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 123 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1981:141191 CAPLUS
DOCUMENT NUMBER: 94:141191
TITLE: Diazo copper complex dyes
INVENTOR(S): Jaeger, Horst
PATENT ASSIGNEE(S): Bayer A.-G., Fed. Rep. Ger.
SOURCE: Ger. Offen., 11 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2925210	A1	19810122	DE 1979-2925210	19790622
EP 23955	A1	19810218	EP 1980-103234	19800611
DE 23955	B1	19810921		
R: CH, DE, FR, GB				
JP 56005858	A2	19810121	JP 1980-82214	19800619
PRIORITY APPLN. INFO.:			DE 1979-2925210	A 19790622

GI




AB Diazo copper complex dyes (I: R = H, Cl-4 alkyl; R1 = H, Me, Et, OMe, OEt; Z = 1- or 2-hydroxynaphthalenesulfonic acid derivative residue) are manufactured by coupling diazotized 3,4-H2N(HO3S)C6H3CH2NRR2 (R defined as above, R2 = acyl component) with a coupling component 4,2-R1(H2N)C6H3OR3 (R1 defined as above, R3 = Me, Et), diazotizing the aminoazo intermediate, coupling with the hydroxynaphthalenesulfonic acid derivative, dealkylatively copperizing and removing R2, and condensing with 2,3-dichloro-6-quinoxalinecarboxyl chloride [1919-43-3]. Thus, I (R = R1 = Me, Z = Z1, azo bond in 2-position, OH in 1-position) [77000-78-3] was prepared by this method.

IT 77000-78-3P
RL: IMP (Industrial manufacture); PREP (Preparation)
(preparation of)

RN 77000-78-3 CAPLUS


CN Cuprate (3-), 3-[[4-[[5-[[[(2,3-dichloro-6-quinoxalinyloxy)methyl]methylamino]methyl]-2-sulfonyl]azo]-2-hydroxy-5-methylphenyl]azo]-4-hydroxy-2,7-naphthalenedisulfonate(5-)-, trihydrogen (9CI) (CA INDEX NAME)

PAGE 1-A



● 3 H*

PAGE 1-B




L13 ANSWER 124 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1981:92561 CAPLUS
DOCUMENT NUMBER: 94:92561
TITLE: Studies on sulfenamidides. V. Anodic oxidation of 4'-substituted 2-nitrobenzenesulfenamidides at a reticulated vitreous carbon electrode
AUTHOR(S): Sayo, Hiroteru; Mori, Koichi; Michida, Takashi
CORPORATE SOURCE: Fac. Pharm. Sci., Kobe-Gakuin Univ., Kobe, 673, Japan
SOURCE: Chemical & Pharmaceutical Bulletin (1980), 28(12), 3707-10
CODEN: CPBTAL; ISSN: 0009-2363
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Constant current electrolysis of 4'-R-2-nitrobenzenesulfenamidides (I; R = OMe, Me, Cl, CO2Et) was carried out in MeCN containing 0.1M ethyltriethylammonium trifluoromethanesulfonate, 1% trifluoroacetic acid, and 1% trifluoroacetic anhydride at a reticulated vitreous C (RVC) electrode. The quantity of electricity to be fed into the electrolytic cell was determined from the anodic potential vs. time curves. The yields of 2,7-di-R-phenazines (R = OMe, Me, Cl, and CO2Et) were 56, 24, 42, and 33%, resp. The RVC anode was useful for electrolysis of I because the considerable yields of phenazines were obtained within several minutes without using an expensive potentiostat.

IT 72848-45-4P
RL: PREP (Preparation)
(preparation of, by electrochem. oxidation of ethoxycarbonylnitrobenzenesulfenamidide on glassy carbon in acetonitrile)

RN 72848-45-4 CAPLUS

CN 2,7-Phenazinedicarboxylic acid, diethyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 125 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1980:93992 CAPLUS
DOCUMENT NUMBER: 92:93992
TITLE: Studies on sulfenamidides. IV. Oxidation of 4'-, 3'-, and 2'-substituted 2-nitrobenzenesulfenamidides with


lead dioxide
AUTHOR(S): Sayo, Hiroteru; Mori, Koichi; Michida, Takashi
CORPORATE SOURCE: Fac. Pharm. Sci., Kobe-Gakuin Univ., Kobe, 673, Japan
SOURCE: Chemical & Pharmaceutical Bulletin (1979), 27(10), 2115-20
CODEN: CPBTAL; ISSN: 0009-2363
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The oxidation of 2-O2NC6H4SNHC6H4R [R = 4-Br (I), 4-CO2Et (II), 4-COMe (III), 4-OEt (IV), 4-NO2 (V), 4-SO2NH2 (VI), 3-Me (VII), 2-OMe (VIII), and 2-Me (IX)] by PbO2 was carried out in MeCN containing 1% CF3CO2H and 1% (CF3CO)2O. The oxidation of I-V and VII gave 2,7-disubstituted phenazines, whereas that of VI, VIII and IX did not. ACNHC6H4NO2-2 was obtained in all cases and (2-O2NC6H4R)2 was obtained from II-IV, VI, and VIII. The oxidation of IX gave a small amount of 2'-methyl-N-[[2-nitrophenyl]thio]-p-benzoquinoneimine, while that of VIII gave a mixture of 2'-methoxy-N-[[2-nitrophenyl]thio]-o- and -p-benzoquinoneimines.

IT 72848-45-4P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 72848-45-4 CAPLUS

CN 2,7-Phenazinedicarboxylic acid, diethyl ester (9CI) (CA INDEX NAME)





L13 ANSWER 126 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1979:152610 CAPLUS
DOCUMENT NUMBER: 90:152610
TITLE: N2-Arylsulfonyl-L-argininamides
INVENTOR(S): Okamoto, Shosuke; Kikumoto, Ryoji; Tamao, Yoshikuni; Okubo, Kazuo; Tazuka, Toru; Tomomura, Shinji; Hijikata, Akiko
PATENT ASSIGNEE(S): Mitsubishi Chemical Industries Co., Ltd., Japan
SOURCE: Ger. Offen., 147 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 15
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2801478	A1	19780720	DE 1978-2801478	19780113
DE 2801478	C2	19910131		
US 4068773	A	19780103	US 1977-760745	19770119
US 4073913	A	19780214	US 1977-760668	19770119
US 4093712	A	19780606	US 1977-760672	19770119
US 4097472	A	19780627	US 1977-760676	19770119
US 4101653	A	19780718	US 1977-760929	19770119
US 4097591	A	19780627	US 1977-776195	19770310
JP 54003037	A2	19790111	JP 1977-66508	19770606
JP 60010028	B4	19850314		
US 4125604	A	19781114	US 1977-804334	19770607
US 4131673	A	19781226	US 1977-804368	19770607
US 4140681	A	19790220	US 1977-804331	19770607

PRIORITY APPLN. INFO.:

US 1977-760568	A	19770119
US 1977-760672	A	19770119
US 1977-760676	A	19770119
US 1977-760745	A	19770119
US 1977-760929	A	19770119
US 1977-771695	A	19770310
JP 1977-66508	A	19770606
US 1977-804331	A	19770607
US 1977-804368	A	19770607
JP 1974-128774	A	19741108
JP 1974-128775	A	19741108
JP 1974-136695	A	19741129
JP 1975-136697	A	19750119
JP 1975-236358	A	19750225
JP 1975-23635	A	19750226
JP 1975-26768	A	19750305
JP 1975-293157	A	19750311

OTHER SOURCE(S) : MARPAT 90:152610
GI



CN1C=NC2=C(N1)C(=C(C=C2)C(=C(C=C3C=C(C=C3)C(=O)O)C=C3)C=C3)C=C3
CN1C=NC2=C(N1)C(=C(C=C2)C(=C(C=C3C=C(C=C3)C(=O)O)C=C3)C=C3)C=C3

as dye precursors.


IT 68765-65-1P
RU: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
(preparation and NMR of)

RN 68765-65-1 CAPLUS

CN Quinoxalium, 6-(ethoxycarbonyl)-2,3-dimethyl-1-phenyl-, perchlorate
(9CI) (CA INDEX NAME)


CC1=C(C)N(C(=O)OCC)c2ccccc2N1
$$\begin{array}{c} \text{O} \\ \parallel \\ \text{O}=\text{Cl}-\text{O} \\ \parallel \\ \text{O} \end{array}$$

L13 ANSWER 128 of 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1978:120955 CAPLUS
 DOCUMENT NUMBER: 88:120955
 TITLE: Some reactions of 2,3-diaminoindole derivatives.
 Synthesis of indolo[2,3-b]quinoxalines
 Kuriilo, G. N.; Roetova, N. I.; Cherkasova, A. A.;
 Grinev, A. N.
 CORPORATE SOURCE: Vses. Nauchno-Issled. Khim.-Farm. Inst. im.
 Ordzhonikidze, Moscow, USSR
 SOURCE: Khimiya Geterotsiklicheskiikh Soedinenii (1977), (12),
 1645-7
 CODEN: KUSSAQ; ISSN: 0453-6234
 DOCUMENT TYPE: Journal
 LANGUAGE: Russian
 OTHER SOURCE(S): CASREACT 88:120955
 OT



AB Treatment of indoles I (R = H, o- and p-Me, p-Cl, p-CO₂Et; R₁ = H, Me) with Cu acetate in MeOH gave 40-64% II (R₂ = MeO); II (R = H, p-Cl; R₁ = H; R₂ = piperidino) were obtained in 45-51% yield in the presence of piperidine. Heating II (R = H, o-Me, p-Cl, p-CO₂Et; R₁ = H; R₂ = OMe) at 200-10° for 5 min gave 15-13% III (R₃ = H, 4-Me, 2-Cl, 2-CO₂Et);


III (R3 = 2-Cl, 2-CO2Et) monooxides were obtained in 61-87% yield by oxidation of III by H2O2.
IT 65880-42-4P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)
RN 65880-42-4 CAPLUS
CN 6H-Indolo[2,3-b]quinoxaline-2-carboxylic acid, 6-methyl-, ethyl ester (9CI) (CA INDEX NAME)




L13 ANSWER 129 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1977:553420 CAPLUS
DOCUMENT NUMBER: 87:153420
TITLE: 2,4-Diaryl[1,3,4H]thiadiazines fused to quinoxalines
INVENTOR(S): Elliott, Arthur John
PATENT ASSIGNEE(S): du Pont de Nemours, S. I., and Co., USA
SOURCE: U.S., 10 pp.
CODEN: USXXAM
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 4025510	A	19770524	US 1975-636792	19751201
US 4025510	A	19770524	US 1975-636792	19751201


PRIORITY APPLN. INFO.:
GI For diagram(s), see printed CA Issue.
AB 2,4-Diaryl-1,3,4-thiadiazines (I, R = Ph, substituted Ph, 4-quinolinyl; R1 = Ph, substituted Ph; A = quinoxaline, pyrazine, pyrimidine, pyridine, pyridazine residue) are prepared and used to dye polyester fibers fast yellow shades. Thus, a mixture of N-thiobenzoyl-N'-phenylhydrazine [13437-75-7], 2,3-dichloroquinoxaline [2213-63-0], and Et3N were refluxed in MeCN to give I [63811-31-4]. The other 34 I were similarly prepared [63811-18-7, 63811-22-3].
IT RL: TEM (Technical or engineered material use); USES (Uses)
(dye, for polyester fibers, preparation of)
RN 63811-18-7 CAPLUS
CN 1H-[1,3,4]Thiadiazino[5,6-b]quinoxaline-7-carboxamide, N-(3-methoxypropyl)-1,3-diphenyl- (9CI) (CA INDEX NAME)




RN 63811-22-3 CAPLUS
CN 1H-[1,3,4]Thiadiazino[5,6-b]quinoxaline-7-carboxamide, N,N-diethyl-1,3-diphenyl- (9CI) (CA INDEX NAME)



IT 26887-34-3 63810-79-7
RL: RCT (Reactant); RACT (Reactant or reagent)
(reaction of, with (thiobenzoyl)phenylhydrazine)
RN 26887-34-3 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N,N-diethyl- (8CI, 9CI) (CA INDEX NAME)



RN 63810-79-7 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(3-methoxypropyl)- (9CI) (CA INDEX NAME)




L13 ANSWER 130 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1977:469732 CAPLUS
DOCUMENT NUMBER: 87:69732
TITLE: Fiber-reactive azo dyes
INVENTOR(S): Jaeger, Horst
PATENT ASSIGNEE(S): Bayer A.-G., Fed. Rep. Ger.
SOURCE: Ger. Offen., 49 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2549570	A1	19770512	DE 1975-2549570	19751105
DE 2549570	C2	19810519		

JP 52058730	A2	19770514	JP 1976-130580	19761101
GB 1540604	A	19790214	GB 1976-45502	19761102
GB 1540605	A	19790214	GB 1977-22502	19761102
FR 2330738	A1	19770603	FR 1976-33489	19761105
FR 2330738	B1	19800808		
CH 624426	A	19810731	CH 1977-12221	19771006

PRIORITY APPLN. INFO.:
GI DE 1975-2549570 A 19751105
CH 1976-13821 A 19761102
GB 1976-45502 A 19761102




AB Fiber-reactive azo dyes (I, R = 5-chloro-2,6-difluoro-4-pyrimidinyl, 2,3-dichloro-6-quinoxaliny; R1 = naphthalenesulfonic acid, pyrazole, acetocetanilide, azo chromophore residue; Z = direct bond, CO) were prepared and used to dye and print cotton and wool fast yellow to blue shades. Thus, 1-amino-2-sulfo-3-(aminomethyl)-4-methoxybenzene [63353-60-6] was prepared, diazotized, coupled with 1,3,6-HOClOH5(SO3H)2 [578-85-8], the resulting azo compound treated with 2,4,6-trifluoro-5-chloropyrimidine [697-83-6], and the reaction mixture salted to give I (R = 5-chloro-2,6-difluoro-4-pyrimidinyl, R1 = 1,3,6,2-HO(SO3H)2ClOH4; Z = direct bond) [63353-68-4], dyeing cotton a fast yellowish red shade.
IT 63395-43-7
RL: TEM (Technical or engineered material use); USES (Uses)
(dye, for cellulosic fibers, preparation of)
RN 63395-43-7 CAPLUS
CN Cuprate(3-), [3-[[[4-[[[3-[[[2,3-dichloro-6-quinoxaliny]carbonyl]amino]methyl]-4-methoxy-2-sulfonyl]azo]-2-hydroxy-5-methylphenyl]azo]-4-hydroxy-2,7-naphthalenedisulfonato(5-)]-], trihydrogen (9CI) (CA INDEX NAME)



PAGE 1-A

• 3 H •


IT 63353-62-8
RL: TEM (Technical or engineered material use); USES (Uses)
(dye, for cotton, preparation of)
RN 63353-62-8 CAPLUS
CN Benzenesulfonic acid, 6-[[[1-(2-chloro-5-sulfonylphenyl)-4,5-dihydro-3-methyl-5-oxo-1H-pyrazol-4-yl]azo]-2-[[[2,3-dichloro-6-quinoxaliny]carbonyl]amino]methyl]-3-methoxy- (9CI) (CA INDEX NAME)



L13 ANSWER 131 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1977:89889 CAPLUS
DOCUMENT NUMBER: 86:89889
TITLE: Azepon[4,5-b]quinoxalines
INVENTOR(S): Hurnaus, Rudolf; Griss, Gerhart; Grell, Wolfgang; Sauter, Robert; Reichl, Richard; Leitold, Matyas
PATENT ASSIGNEE(S): Thomas, Dr. Karl, G.m.b.H., Fed. Rep. Ger.
SOURCE: Ger. Offen., 46 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2519258	A1	19761111	DE 1975-2519258	19750430
PRIORITY APPLN. INFO.:			DE 1975-2519258	A 19750430

GI




AB Tetrahydro-1H-azepino[4,5-b]quinoxalines (I; R = e.g., H, Me, Ph, PhCH₂, Ac, Bz, CO₂H, CH₂CH₂CO₂H; R₁ = R₂ = H, OH, AcO, EtOCO; R₃n = e.g., H, 8-Cl, 7-NO₂, 8-Me, 8-CO₂H, 8,9-Me₂, 8-MeO), useful as appetite depressants and bactericides (no date), are prepared by various known methods, mostly involving reaction between an o-phenylenediamine and an azepinedione. The azepinedione can be obtained by cyclization of an iminodipropionic acid derivative. Thus, reaction of PhCH₂N(CH₂CH₂CO₂Me) with Na and Me₃SiCl in Me₂COH₄ gives 1-benzyl-2,3,6,7-tetrahydro-4,5-bis(trimethylsiloxy)-1H-azepine which is oxidized with Br to the azepine-4,5-dione which then reacts with 1,2-(H₂N)C₆H₄ in AcOH to give after 4 hr at 100° 83% 1.HCl (R = PhCH₂, R₁ = R₂n = R₃ = H).

IT 61793-52-0P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 61793-52-0 CAPLUS

CN 1H-Azepino[4,5-b]quinoxaline-8-carboxylic acid, 2,3,4,5-tetrahydro-3-(phenylmethyl)-, ethyl ester (9CI) (CA INDEX NAME)




L13 ANSWER 132 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1976:412370 CAPLUS
DOCUMENT NUMBER: 85:12370
TITLE: Stable polymer images by photopolymerization in a matrix
INVENTOR(S): Baumann, Nikolaus
PATENT ASSIGNEE(S): Ciba-Geigy A.-G., Switz.
SOURCE: Ger. Offen., 109 pp.
CODEN: GWXXBX

DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2526674	A1	19760102	DE 1975-2526674	19750609
CH 594704	A	19780131	CH 1974-7956	19740611
CH 604208	A	19780831	CH 1974-7957	19740611
FR 2274951	A1	19760109	FR 1975-17754	19750606
FR 2274951	B1	19831028		
CA 1077760	A1	19800520	CA 1975-228849	19750609
BE 830049	A1	19751210	BE 1975-157171	19750610
JP 51030286	A2	19760315	JP 1975-71385	19750611

US 4043819 A 19770823 US 1976-743011 19761118
PRIORITY APPLN. INFO.: CH 1974-7956 A 19740611
CH 1974-7957 A 19740611
CH 1975-4843 A 19750416
US 1975-584444 A1 19750606

GI




AB Photopolymerizable compns. capable of forming stable polymer relief images are composed of an ethylenically unsatd. photopolymerizable monomer, a chemical hardenable, nonlight-sensitive, swellable macromol. compound as binder, a hardening agent, and a quinoxaline derivative (I; R = H, Me; R₁ = Me, Ph, NaO₃SC₆H₄, p-MeOC₆H₄; R₂ = H, SO₃H), which together with the monomer or the binder forms a redox pair, as photoinitiator. Thus, a gelatin-subbed cellulose triacetate support was overcoated with a solution containing calcium diacrylate 36.07, acrylamide 3.89, gelatin 7.68, glycerin 2.14, the ether of polyethylene glycol with N-(methylol)perfluoroalkylsulfonamide 0.107, β-(3,5-dimethyl-1-pyrazolyl)acrolein (hardener) 0.081, and I (R, R₂ = H; R₁ = NaO₃SC₆H₄) 1.03 g/m², dried at 30°, contact exposed under a photocg. step wedge (12 steps) for approx. 30 sec to a 400-W high-pressure Hg lamp at 40 cm, rinsed with water, colored with a cationic dye, rinsed with water, and dried to show 12 steps. The maximum color d. was 3.2 and the absolute sensitivity at an optical d. of 1 was 8.2 + 10⁻³ J/cm².

IT 32388-05-9P 32388-06-0P 32388-08-2P
32388-09-3P 37966-43-1P 52996-75-5P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)


RN 32388-05-9 CAPLUS

CN 6-Quinoxalinecarboxylic acid, 2,3-diphenyl-, ethyl ester (8CI, 9CI) (CA INDEX NAME)



RN 32388-06-0 CAPLUS


CN 6-Quinoxalinecarboxylic acid, 2,3-dimethyl-, ethyl ester (6CI, 8CI, 9CI) (CA INDEX NAME)



RN 32388-08-2 CAPLUS


CN 6-Quinoxalinecarboxamide, N,N-diethyl-2,3-dimethyl- (8CI, 9CI) (CA INDEX NAME)

NAME)




RN 32388-09-3 CAPLUS

CN 6-Quinoxalinecarboxamide, N,N-diethyl-2,3-diphenyl- (8CI, 9CI) (CA INDEX NAME)




RN 37966-43-1 CAPLUS

CN 6-Quinoxalinecarboxylic acid, 2,3-bis[(acetyloxy)methyl]-, ethyl ester (9CI) (CA INDEX NAME)



RN 52996-75-5 CAPLUS

CN 6-Quinoxalinecarboxylic acid, 2,3-bis(bromomethyl)-, ethyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 133 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1975:423165 CAPLUS
DOCUMENT NUMBER: 83:22365
TITLE: Effect of 1-phenazinecarboxylic acid derivatives on experimental tumors
AUTHOR(S): Sidorik, O. A.; Shevchenko, I. N.
CORPORATE SOURCE: Inst. Probl. Onkol., Kiev, USSR
SOURCE: Pirolochevskii Aktivnye Veshchestva (1966-1992) (1974), 6, 92-4
CODEN: FAVUAI; ISSN: 0533-1153


DOCUMENT TYPE: Journal
LANGUAGE: Russian
GI For diagram(s), see printed CA Issue.

AB When injected s.c. at 200 mg/kg/day into rats for 12-15 days or i.p. at 50 mg/kg/day into mice for 8-12 days, 1-phenazinecarboxylic acid Na salt (I) [1144-02-1] or glycine N-1-phenazinecarboxyl acid Na salt (II) [55327-47-4] significantly inhibited the growth of Ehrlich carcinoma and erythromyeloidia. The inhibitory effects of the prepns. on lymphoma NK/Ly or sarcoma 45 were less pronounced and both compds. were inactive against Guerin carcinoma.

IT 55327-47-4
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(neoplasia inhibitor)

RN 55327-47-4 CAPLUS

CN Glycine, N-(2-phenazinylcarbonyl)-, monosodium salt (9CI) (CA INDEX NAME)



● Na


L13 ANSWER 134 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1975:156377 CAPLUS
DOCUMENT NUMBER: 82:156377
TITLE: Piperazinyl quinoxalines
INVENTOR(S): Engelhard, Edward L.; Lumma, William C., Jr.; Saari, Waifred S.
PATENT ASSIGNEE(S): Merck and Co., Inc.
SOURCE: Ger. Offen., 36 pp.
CODEN: GWXXBX

DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2433397	A1	19750206	DE 1974-2433397	19740711
FI 7401939	A	19750114	FI 1974-1939	19740625
DK 7403426	A	19750303	DK 1974-3426	19740626
NO 7402351	A	19750114	NO 1974-2351	19740627
SE 7408486	A	19750114	SE 1974-8486	19740627
SE 417316	B	19810309		
SE 417316	C	19810625		
NL 7408705	A	19750115	NL 1974-8705	19740627
AU 7470731	A1	19760108	AU 1974-70731	19740702
GB 1440722	A	19760623	GB 1974-30176	19740708
ES 428107	A1	19761116	ES 1974-428107	19740709
FR 2236499	A1	19750207	FR 1974-24114	19740711
DD 112127	C	19750320	DD 1974-179871	19740711
BE 817608	A1	19750113	BE 1974-146519	19740712
ZA 7404466	A	19760225	ZA 1974-4466	19740712
CH 605919	A	19780103	CH 1974-9648	19740712

JP 50037791 A2 19750408 JP 1974-79774 19740713
 PRIORITY APPLN. INFO.: US 1973-379022 A 19730713
 US 1974-465381 A 19740429

GI For diagram(s), see printed CA Issue.
 AB Piperazinyloquinolines I (R = H, Me, CO₂Me, COCH₂Ac, CH₂CH₂OH, Ac, CO₂H, CH₂CH₂Ph, CO₂CH₂CO₂Et, allyl, CH₂CH₂CHCl; R₁ = H, Cl, Me, CO₂H, Ph, CO₂Et, SPh, Ac, NHCH₂CH₂OH, NH₂, OEt; R₂ = halo, NO₂, OMe, CF₃ etc. in 5-6 positions) and some related compds. (50 compds.) were prepared for use as antidepressants, appetite depressants, and analgesics. Thus reaction of 2,3-dichloro-6-cyanoquinoline with N-formylpiperazine and reduction over Pd-C gave I (R = R₁ = H, R₂ = 6-CN).
 IT 55686-57-2
 RN 55686-57-2 CAPLUS
 CN 6-Quinoxalinecarboxamide, 3-chloro-N-(2-hydroxyethyl)-2-[4-(2-hydroxyethyl)-1-piperazinyl]-, monohydrochloride (9CI) (CA INDEX NAME)



IT 26773-13-7
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (reaction of, with piperazine deriva.)
 RN 26773-13-7 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-hydroxyethyl)- (8CI, 9CI) (CA INDEX NAME)




L13 ANSWER 135 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1974:570266 CAPLUS
 DOCUMENT NUMBER: 81:170266
 TITLE: Photopolymerization of ethylenically unsaturated compounds
 INVENTOR(S): Baumann, Nikolaus; Schlunke, Hans P.
 PATENT ASSIGNEE(S): Ciba-Geigy A.-G.
 SOURCE: Ger. Offen., 77 pp.
 CODEN: GWXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE


DE 2360350	A1	19740606	DE 1973-2360350	19731204
CH 573448	A	19760315	CH 1972-17658	19721205
NL 1715878	A	19740607	NL 1973-15878	19731120
FR 2217355	A1	19740906	FR 1973-42221	19731127
US 4001017	A	19770104	US 1973-420176	19731129
CA 980350	A1	19760316		
IT 997924	A	19751230	CA 1973-187233	19731203
GB 1436589	A	19760519	IT 1973-54054	19731203
GB 1436590	A	19760519	GB 1973-55925	19731203
BE 808179	A1	19740604	GB 1975-10419	19731203
JP 49086383	A2	19740819	BE 1973-138476	19731204
JP 59024147	B4	19840607	JP 1973-135455	19731205
JP 49087780	A2	19740822	JP 1973-135454	19731205
JP 59028205	B4	19840711		
FR 2221453	A1	19741011	FR 1974-13181	19740416
FR 2221453	B1	19780908		

PRIORITY APPLN. INFO.: CH 1972-17658 A 19721205
 CH 1973-14714 A 19731017


AB Quinoxaline deriva. and salts (sim.150) were described which were useful with electron donors, such as Na p-toluenesulfinate (I) (824-79-3) and triphenylphosphine [603-35-0], as photoredox catalysts for the preparation of acrylamide-barium diacrylate copolymer [37281-67-7] (e.g., on photog. film supports) or a similar polymer upon exposure to radiation (200-450 nm). Thus, a solution of 1.4 M aqueous Ba diacrylate 180, 1.6 M aqueous acrylamide 60, 5% aqueous gelatin 30, and 0.25% aqueous FC 170 (wetting agent) 30 ml was mixed with 2 ml EtOH containing 10 mg 6,7-ethylenedioxy-2,3-bis(hydroxymethyl)quinoxaline (II) [52996-38-0] and 2 ml 0.016 M aqueous I, coated on gelatin-coated cellulose triacetate film, dried, irradiated with a Hg lamp through a photog. step-wedge for 30 sec, and developed with a dye to give discernible shades corresponding to the steps.
 IT 32388-05-9 32388-06-0 32388-08-2
 32388-09-3 37966-43-1 52996-75-5
 RL: CAT (Catalyst use); USES (Uses)
 (catalysts, contg electron donors, for photopolymer)
 RN 32388-05-9 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-diphenyl-, ethyl ester (8CI, 9CI) (CA INDEX NAME)




RN 32388-06-0 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-dimethyl-, ethyl ester (6CI, 8CI, 9CI) (CA INDEX NAME)




RN 32388-08-2 CAPLUS
 CN 6-Quinoxalinecarboxamide, N,N-diethyl-2,3-dimethyl- (8CI, 9CI) (CA INDEX NAME)




RN 32388-09-3 CAPLUS
 CN 6-Quinoxalinecarboxamide, N,N-diethyl-2,3-diphenyl- (8CI, 9CI) (CA INDEX NAME)



RN 37966-43-1 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-bis[(acetyloxy)methyl]-, ethyl ester (9CI) (CA INDEX NAME)



RN 52996-75-5 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-bis(bromomethyl)-, ethyl ester (9CI) (CA INDEX NAME)




L13 ANSWER 136 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1974:146079 CAPLUS
 DOCUMENT NUMBER: 80:146079
 TITLE: Nuclear magnetic resonance studies of heterocyclic bridged biphenyls
 AUTHOR(S): Hall, D. Muriel; Hwang, Huaun-Yong; Bhanthumnavin, Biravana
 CORPORATE SOURCE: Dep. Chem., Bedford Coll., London, UK

SOURCE: Journal of the Chemical Society, Perkin Transactions 2: Physical Organic Chemistry (1972-1999) (1973), (15), 2131-4
 CODEN: JCPKBN; ISSN: 0300-9580

DOCUMENT TYPE: Journal
 LANGUAGE: English

GI For diagram(s), see printed CA Issue.
 AB Condensation reactions between 1,2-diamines and biphenyl-2,2'-dicarboxaldehyde or 9,10-phenanthraquinone, and between 2,2'-diaminobiphenyl and 1,2-diketones, gave polycyclic products with 5-, 6-, 7-, and 8-membered heterocyclic rings, the NMR spectra of which are discussed. The 15H-dibenzo[*a,c*]benzimidazo[1,2-*a*]azepines I (R = H, CO₂Et) are fluxional. The dibenzo[*a,c*]phenazines II showed large downfield shifts (δ approx.9.5) for some aromatic protons.
 IT 51448-37-4P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 RN 51448-37-4 CAPLUS
 CN Dibenzo[*a,c*]phenazine-11-carboxylic acid, ethyl ester (9CI) (CA INDEX NAME)



L13 ANSWER 137 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1974:109835 CAPLUS
 DOCUMENT NUMBER: 80:109835
 TITLE: Azo reactive dyes
 INVENTOR(S): Jaeger, Horst
 PATENT ASSIGNEE(S): Bayer A.-G.
 SOURCE: Ger. Offen., 47 pp.
 CODEN: GWXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2232541	A1	19740117	DE 1972-2232541	19720703
DE 2232541	B2	19771027		
IT 990615	A	19750710	IT 1973-26026	19730628
BE 801661	A1	19740102	BE 1973-132901	19730629
JP 49052828	A2	19740522	JP 1973-73009	19730629
JP 55043025	B4	19801104		
CH 739539	A	19750530	CH 1973-9539	19730629
CH 572546	B	19760213		
CA 994330	A1	19760803	CA 1973-175278	19730629
CH 582739	A	19761215	CH 1975-15476	19730629
NL 7309200	A	19740107	NL 1973-9200	19730702
DD 107302	C	19740720	DD 1973-171990	19730702
ES 416498	A1	19760301	ES 1973-416498	19730702
GB 1413122	A	19760407	GB 1973-31420	19730702
GB 1431323	B	19760407	GB 1975-19759	19730702
AT 320100	B	19750127	AT 1973-5845	19730703
FR 2236905	A1	19750207	FR 1973-24415	19730703

US 4126609 A 19781121 US 1973-376184 19730703
AT 7400179 A 19760515 AT 1974-179 19740110
AT 334113 B 19760110
JP 52063488 A2 19770525 JP 1976-11493 19760206
JP 57029592 B4 19820623
US 4049704 A 19770920 US 1976-656251 19760209
DE 1972-2232541 A 19720703
AT 1973-5845 A 19730703
US 1973-376184 A3 19730703


PRIORITY APPLM. INFO.:

AB Azo and diazo fiber-reactive dyes (I, R = 1-(sulfonylphenyl)-4-pyrazolyl derivative, aminohydroxysulfonaphthyl derivative, [(sulfonylphenyl)azo]sulfonaphthyl derivative; R1 = 2,6-difluoro-5-chloro-4-pyrimidinyl, 2,6-dichloroquinoline-6-carbonyl) were prepared and were used to dye cotton fast greenish yellow to navy blue shades. Thus, 2,1-AcNHClOH6SO3H in H2SO4 was treated with N-(hydroxymethyl)phthalimide at 15-20 deg. for 24 hr and the product treated with aqueous NaOH at 170 deg. to give 2-amino-5-(aminomethyl)-1-naphthalenesulfonic acid [52084-84-1] which was diazotized and coupled with 1,6,3,5-NO(BaNH)ClOH4(SO3H)2 to give an azo intermediate (II); treatment of II with 2,4,6-trifluoro-5-chloropyrimidine gave azo reactive dye (III) [51366-30-4]. The other I were similarly prepared

IT 52084-87-49
RL: IMF (Industrial manufacture); PREP (Preparation)

(Preparation of)

RN 52084-87-4 CAPLUS
CN 1-Naphthalenesulfonic acid, 2-[[1-(2-chloro-5-sulfonylphenyl)-4,5-dihydro-3-methyl-5-oxo-1H-pyrazol-4-yl]azo]-5-[[1-(2,3-dichloro-6-quinoxaliny)carbonyl]amino]methyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 138 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1973:418669 CAPLUS
DOCUMENT NUMBER: 79:18669


TITLE: Reaction of sodium borohydride with heteroaromatic nitro compounds
AUTHOR(S): Rao, Koppaka V.; Jackman, Dennis
CORPORATE SOURCE: J. Hillis Miller Health Cent., Univ. Florida, Gainesville, FL, USA
SOURCE: Journal of Heterocyclic Chemistry (1973), 10(2), 213-15
CODEN: JHTCAD; ISSN: 0022-152X

DOCUMENT TYPE: Journal
LANGUAGE: English
GI For diagram(s), see printed CA Issue.
AB Quinoxalines (I, R = 5-, 6-NO2, 6-CN, 6-CO2Et, 6-CF3) and 5-, 6-, 7-, 8-nitroquinoline (II) were reduced selectively by NaBH4 in HOAc at 5° to give 1,2,3,4-tetrahydro-derivs. of I and 1,2-dihydro-derivs.

of II resp. 5-Nitroisoquinoline was reduced to the 1,2,3,4-tetrahydro derivative in HOAc at 5° but yielded the 1,2-dihydro derivative in aqueous MeOH.

IT 6924-72-7
RL: RCT (Reactant); RACT (Reactant or reagent)
(reduction of, by sodium borohydride in acetic acid)

RN 6924-72-7 CAPLUS
CN 6-Quinoxalinecarboxylic acid, ethyl ester (6CI, 7CI, 8CI, 9CI) (CA INDEX NAME)



L13 ANSWER 139 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 1973:73615 CAPLUS

DOCUMENT NUMBER: 78:72615

TITLE: Fiber-reactive azo dyes

INVENTOR(S): Jaeger, Horst; Dehmelt, Georg

PATENT ASSIGNEE(S): Farbenfabriken Bayer A.-G.

SOURCE: Ger. Offen., 40 pp.
CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE

DE 2113298 A 19720921 DE 1971-2113298 19710319

DE 2113298 B2 19770512

DE 2113298 C3 19771229

NL 7203510 A 19720921 NL 1972-3510 19720316

BE 740048 B 19720918 BE 1972-115221 19720317

FR 2130422 A5 19721103 FR 1972-9516 19720317

FR 2130422 B1 19751024

IT 953527 A 19730810 IT 1972-22048 19720317

AT 309619 B 19730827 AT 1972-2310 19720317

DD 102159 C 19731212 DD 1972-161627 19720317

AT 316479 B 19740710 AT 1972-7973 19720317

GB 1378244 A 19741227 GB 1972-12626 19720317

CA 998388 A1 19761012 CA 1972-137324 19720317

CH 606341 A 19781031 CH 1974-8948 19720317

JP 54027020 B4 19790907 JP 1972-26718 19720317

ES 400909 A 19750116 ES 1972-400909 19720318

US 4118382 A 19781003 US 1977-600573 19770525

PRIORITY APPLM. INFO.:

DE 1971-2113298 A 19710319


US 1972-235856 A1 19720317

AB 4'-Amino-3',5'-disulfoacetacetanilide (I) [37615-83-1] was used as the coupling component to prepare 6 fiber-reactive azo dyes (II, Q = phenyl, naphthotriazolyl, or benzothiazolyl) containing a fiber-reactive group which dyed cotton light- and wetfast yellow to greenish yellow shades. Thus, 1,4,2,6-(H2N)2C6H2(SO3H)2 was condensed with diketene in aqueous NaOH to give I. 2,5-H2N(O2N)C6H3SO3H was diazotized and coupled with I, the intermediate nitro azo compound reduced with Na2S, and the amino derivative condensed with 2,3-dichloroquinoline-6-carbonyl chloride to give azo dye III [37615-47-7], greenish yellow on cotton. In another typical example, 2-(methylsulfonyl)-6-methoxy-7-aminobenzothiazole was diazotized and coupled with I to give azo dye IV [37615-48-8], yellow on cotton.


IT 40859-07-2P
RL: IMF (Industrial manufacture); PREP (Preparation)

(Preparation of)

RN 40859-07-2 CAPLUS
CN 2H-Naphtho[1,2-d]triazole-5,9-disulfonic acid, 7-[[1-[[[4-amino-3,5-disulfonylphenyl]amino]carbonyl]-2-oxopropyl]azo]-2-[[1-[[[2,3-dichloro-6-quinoxaliny)carbonyl]methylamino]methyl]phenyl]- (9CI) (CA INDEX NAME)



PAGE 1-A



PAGE 1-B

L13 ANSWER 140 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1972:552222 CAPLUS
DOCUMENT NUMBER: 77:152222

TITLE: Substituted quinoxalines for the inhibition of gastric acidity
INVENTOR(S): Bolhofer, William A.; Baldwin, John J.
CORPORATE SOURCE: Merck and Co., Inc.
SOURCE: U.S., 5 pp.
CODEN: USXXAM

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE

US 3691166 A 19720912 US 1969-886791 19691219

PRIORITY APPLM. INFO.:

GI For diagram(s), see printed CA Issue.


AB Seven 2,3-dichloro[(2-hydroxyethyl)carbamoyl]quinoxalines (I, when R = CONH(CH2)2OH, R1 = H, Cl, R2 = H, Cl, Me, R3 = H, Cl, Me; when R1 = CONH(CH2)2OH, R = R2 = H, R3 = Cl, Me), gastric acid inhibitors, were prepared by refluxing the 2,3- and 3,4-diaminobenzoic acids with di-Et oxalate, converting the resulting 2,3-dihydroxy-5-(or 6)-quinoxalinecarboxylic acids with PC15-POCl3 to the 2,3-dichloroquinoxalinecarboxylic acid chlorides and treating these with HO(CH2)2NH2 at room temperature

IT 37901-98-7P 37902-03-7P
RL: BPH (Synthetic preparation); PREP (Preparation)

(Preparation of)


RN 37901-98-7 CAPLUS

CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-hydroxyethyl)-8-methyl- (9CI) (CA INDEX NAME)



RN 37902-03-7 CAPLUS

CN 6-Quinoxalinecarboxamide, 2,3,8-trichloro-N-(2-hydroxyethyl)- (9CI) (CA INDEX NAME)



L13 ANSWER 141 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 1972:513283 CAPLUS

DOCUMENT NUMBER: 77:113283

TITLE: Proton resonance spectra of heterocycles. VIII.

Substituent effects on coupling constants in bicyclic heteroaromatic compounds and the prediction of chemical shifts from coupling constants

Katritzky, A. R.; Takeuchi, Y.

Sch. Chem. Sci., Univ. East Anglia, Norwich, UK

SOURCE: Journal of the Chemical Society, Perkin Transactions 2: Physical Organic Chemistry (1972-1999) (1972), (11), 1682-5

CODEN: JCPKBI; ISSN: 0300-9580

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Substituents affect coupling conste. in bicyclic heteroaromatic compds.

e.g., substituted 2,1,3-selenadiazole, benzofuroxan, quinoxaline, quinoline; they increase 3J and 4J, slightly decrease 4J and 5J, and have little effect on 3J. The regularities discussed allowed prediction of J in substituted systems. Ortho-substituent effects on chemical shifts relate to the corresponding 3J values in the unsubstituted compound, reflecting the dependence of both on partial bond fixation.


IT 6924-72-7

RL: PRP (Properties)

(substituent coupling conste. in, substituent chemical shift in relation to)

RN 6924-72-7 CAPLUS


CN 6-Quinoxalinecarboxylic acid, ethyl ester (6CI, 7CI, 8CI, 9CI) (CA INDEX NAME)



L13 ANSWER 142 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1972:503277 CAPLUS
 DOCUMENT NUMBER: 77:103277
 TITLE: Fiber reactive dyes
 INVENTOR(S): Dehmelt, Georg; Jaeger, Horst
 PATENT ASSIGNER(S): Farbenfabriken Bayer A.-G.
 SOURCE: Ger. Offen. 86 pp.
 CODEN: GWXIBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2055967	A	19720518	DE 1970-2055967	19701113
IT 951598	A	19730710	IT 1971-30972	19711111
CH 549632	A	19740531	CH 1973-4115	19711111
BE 775265	A1	19720301	BE 1971-110436	19711112
NL 7115598	A	19720516	NL 1971-15598	19711112
FR 2114670	A5	19720630	FR 1971-40675	19711112
GB 1369856	A	19741009	GB 1971-52667	19711112
JP 56003389	B4	19810124	JP 1971-89976	19711112

PRIORITY APPLN. INFO.:
 AB Fourteen H₂O-soluble fiber-reactive azo, phthalocyanine, nitro, and anthraquinone dyes containing SO₂NHSO₂(CH₂)_nNMeX groups (X = 2,3-dichloroquinoxalin-6-carbonyl or 5-chloro-2,6-difluoropyrimidin-4-yl, n = 3 or 4) were prepared and used to dye cellulose and wool wetfast shades. For example, p-H₂NC₆H₄SO₂NHSO₂CH₂CH₂CH₂NHMe .fer. 1-(2-chloro-4-sulfonylphenyl)-3-methyl-5-pyrazolone was condensed with 2,3-dichloroquinoxaline-6-carbonyl chloride to give fiber-reactive dye I [35934-00-0].
 IT 35934-00-0P 38097-34-6P 38097-35-7P
 38153-46-7P
 RL: IMF (Industrial manufacture); PRBP (Preparation)
 (preparation of)
 RN 35934-00-0 CAPLUS
 CN Benzenesulfonic acid, 3-chloro-4-[4-[[[4-[[[2,3-dichloro-6-quinoxaliny]carbonyl]methylamino]propyl]sulfonyl]amino]sulfonyl]phenyl]azo-4,5-dihydro-3-methyl-5-oxo-1H-pyrazol-1-yl]- (9CI) (CA INDEX NAME)



PAGE 1-A


D1-SO₃⁻




• H⁺

RN 38097-35-7 CAPLUS
 CN Cuprate(1-), [C-(aminosulfonyl)-C,C-bis[[[4-[[[2,3-dichloro-6-quinoxaliny]carbonyl]methylamino]butyl]sulfonyl]amino]sulfonyl]-29H,31H-phthalocyanine-C-sulfonate(3-)-N29,N30,N31,N32]-, hydrogen (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 38097-34-6 CAPLUS
 CN Cuprate(1-), [C-(aminosulfonyl)-C,C-bis[[[4-[[[2,3-dichloro-6-quinoxaliny]carbonyl]methylamino]butyl]sulfonyl]amino]sulfonyl]-29H,31H-phthalocyanine-2-sulfonate(3-)-N29,N30,N31,N32]-, hydrogen (9CI) (CA INDEX NAME)

PAGE 1-A




D1-SO₃⁻




• H⁺

RN 38153-46-7 CAPLUS
 CN 2-Naphthalenesulfonic acid, 7-(acetylamino)-3-[[[4-[[[2,3-dichloro-6-quinoxaliny]carbonyl]methylamino]propyl]sulfonyl]amino]sulfonyl]phenyl]azo-4-hydroxy- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




L13 ANSWER 143 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1972:461948 CAPLUS
 DOCUMENT NUMBER: 77:61948
 TITLE: Synthesis and study of phenazine derivatives. XVII. Synthesis and properties of some phenazine derivatives and their N-mono- and N,N-dioxides
 AUTHOR(S): Batulina, R. Kh.; Konyukhov, V. N.; Pushkareva, Z. V.; Yarysheva, I. A.
 CORPORATE SOURCE: Ural. Politekhn. Inst. im. Kirova, Sverdlovsk, USSR
 SOURCE: Khimiya Geterotsiklicheskih Soedinenii (1972), (4), 563-7
 CODEN: KGSSAQ; ISSN: 0132-6244
 DOCUMENT TYPE: Journal
 LANGUAGE: Russian
 OI For diagram(s), see printed CA issue.
 AB Five 2-phenazincarboxamide 10-oxides (I, R = H, Et, R1 = H, Et, Ph,


p-MeOC6H4, RR1 = (CH2CH2)20, n = 0) and 4 2-phenazinecarboxamide
5,10-dioxides I (n = 1) were obtained in 46-86% yield. Polarog. of 36 I
and previously obtained phenazinecarboxamides was reported.
IT 30806-87-2 30806-88-3 30905-67-0
30905-73-8 30905-74-9 37648-78-5
37648-80-9 37648-82-1
RL: RCT (Reactant); RACT (Reactant or reagent)
(polarog. of)
RN 30806-87-2 CAPLUS
CN Glycine, N-(2-phenazinylcarbonyl)-, ethyl ester (8CI, 9CI) (CA INDEX
NAME)




RN 30806-88-3 CAPLUS
CN Glycine, N-(2-phenazinylcarbonyl)-, ethyl ester, dioxide (9CI) (CA INDEX
NAME)




RN 30905-67-0 CAPLUS
CN L-Valine, N-(2-phenazinylcarbonyl)-, ethyl ester (9CI) (CA INDEX NAME)
Absolute stereochemistry.




RN 30905-73-8 CAPLUS
CN L-Valine, N-(2-phenazinylcarbonyl)-, ethyl ester, dioxide (9CI) (CA INDEX
NAME)
Absolute stereochemistry.




RN 30905-74-9 CAPLUS
CN L-Valine, N-(2-phenazinylcarbonyl)-, dioxide (9CI) (CA INDEX NAME)
Absolute stereochemistry.




RN 37648-78-5 CAPLUS
CN 2-Phenazinecarboxylic acid, ethyl ester, 5,10-dioxide (9CI) (CA INDEX
NAME)




RN 37648-80-9 CAPLUS
CN 2-Phenazinecarboxylic acid, ethyl ester (9CI) (CA INDEX NAME)




RN 37648-82-1 CAPLUS
CN 2-Phenazinecarboxamide, N,N-diethyl- (9CI) (CA INDEX NAME)




IT 30806-89-4
RL: RCT (Reactant); RACT (Reactant or reagent)
(polarog. of)
RN 30806-89-4 CAPLUS
CN Glycine, N-(2-phenazinylcarbonyl)-, dioxide (9CI) (CA INDEX NAME)




IT 37648-67-2P 37648-71-8P 37648-75-2P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and polarog. of)
RN 37648-67-2 CAPLUS
CN 2-Phenazinecarboxamide, N,N-diethyl-, 10-oxide (9CI) (CA INDEX NAME)



RN 37648-71-8 CAPLUS
CN 2-Phenazinecarboxamide, N,N-diethyl-, 5,10-dioxide (9CI) (CA INDEX NAME)



RN 37648-75-2 CAPLUS
CN 2-Phenazinecarboxylic acid, ethyl ester, 10-oxide (9CI) (CA INDEX NAME)




L13 ANSWER 144 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1972:454870 CAPLUS
DOCUMENT NUMBER: 77:54870
TITLE: Quinoxaline catalysts for the silver-dye bleach
process
INVENTOR(S): Schlunke, Hans P.; Egli, Christian
PATENT ASSIGNEE(S): Ciba-Geigy A.-G.
SOURCE: Ger. Offen., 83 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2144297	A	19720109	DE 1971-2144297	19710903
DE 2144297	C2	19821202		
CH 553428	A	19740830	CH 1970-13253	19700904
FR 2106207	A5	19720428	FR 1971-31379	19710830
US 3796576	A	19740312	US 1971-176749	19710831
GB 1360046	A	19740717	GB 1971-41053	19710902
BE 772142	A1	19720303	BE 1971-107759	19710903
JP 54003620	B4	19790224	JP 1971-68809	19710904
US 3875158	A	19750401	US 1973-344815	19730326
			CH 1970-13253	A 19700904
			US 1971-176749	A2 19710831

AB Quinoxalines, having a favorable redox potential and adequate solubility in the dye bleach bath, carry in their 2- and 3-positions a -CH2- group linked to Br, C, O, S, or N. They are synthesized from 1,2-dinitrobenzene or o-nitroaniline compds. by hydrogenation to the 1,2-diamines, followed by condensation with 1,2-dicarbonyl compds. in a N atmospheric. The halogen in the 2,3-bis(bromomethyl)quinoxalines can readily be exchanged by reaction with Lewis bases. Thus, 2,3-bis(bromomethyl)-6,7-di(methoxy)quinoxaline is obtained by reduction of 1,2-dimethoxy-4,5-dinitrobenzene and condensation with BrCH2COCCH2Br. Reaction with (1) K2CO3 in aqueous EtOH replaces the Br by OH, (2) KOAc by OAc, and (3) NaOMe by OMe. The quinoxalines are added to the bleach bath (1-100 mg/l.) or incorporated in a layer of the photog. material.

IT 37966-43-1
RL: USES (Uses)
(photog. silver-dye bleach bath containing, for color processing)
RN 37966-43-1 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2,3-bis(acetyloxy)methyl-, ethyl ester (9CI) (CA INDEX NAME)




L13 ANSWER 145 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1972:434584 CAPLUS
 DOCUMENT NUMBER: 77:34584
 TITLE: Anticancerous 2-substituted phenazine 5,10-dioxides
 PATENT ASSIGNEE(S): Societe des usines chimiques de Rhone-Poulenc
 SOURCE: Fr. CAM, 2 pp. Addn. to Fr. M 4745 (CA 69:67421f).
 CODEN: FMXXBK
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 193		19680410	FR 1965-60633	19650618

GI For diagram(s), see printed CA issue.
 AB N,N'-Carbonyldiimidazole in DMF was treated with 2-carboxyphenazine 5,10-dioxide to give 2-imidazolocarboxylphenazine 5,10-dioxide. 1-Benzylpiperazine was added, and the solution concentrated to give 2-(4-benzyl-1-piperazinyl)carbonylphenazine 5,10-dioxide (I) (R = 4-benzyl-1-piperazinyl). Similarly prepared were I (R = [2-(dimethylamino)ethyl]amino), [3-(dimethylamino)-2-methylpropyl]amino, [3-(dimethylamino)propyl]amino, 32-(1-pyrrolidinyl)ethyl]amino, [2-(4-methyl-1-piperazinyl)ethyl]amino and 2-(4-ethyl-1-piperazinyl)].
 IT 13458-25-8P 13458-27-OP 13458-29-2P
 13458-30-5P 14559-63-8P
 RL: BPN (Synthetic preparation); PREP (Preparation) (preparation of)


RN 13458-25-8 CAPLUS
 CN 2-Phenazinecarboxamide, N-[2-(dimethylamino)ethyl]-, 5,10-dioxide (8CI, 9CI) (CA INDEX NAME)




RN 13458-27-0 CAPLUS
 CN 2-Phenazinecarboxamide, N-[3-(dimethylamino)-2-methylpropyl]-, 5,10-dioxide (8CI, 9CI) (CA INDEX NAME)




RN 13458-29-2 CAPLUS
 CN 2-Phenazinecarboxamide, N-[2-(1-pyrrolidinyl)ethyl]-, 5,10-dioxide (8CI, 9CI) (CA INDEX NAME)



RN 13458-30-5 CAPLUS
 CN 2-Phenazinecarboxamide, N-[3-(dimethylamino)propyl]-, 5,10-dioxide (8CI, 9CI) (CA INDEX NAME)




RN 14559-63-8 CAPLUS
 CN 2-Phenazinecarboxamide, N-[2-(4-methyl-1-piperazinyl)ethyl]-, 5,10-dioxide (8CI, 9CI) (CA INDEX NAME)



L13 ANSWER 146 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1972:419006 CAPLUS
 DOCUMENT NUMBER: 77:19006
 TITLE: Polarography and ir spectra of 2,2'-biphenaziny and

AUTHOR(S): its derivatives
 CORPORATE SOURCE: Gordienko, L. L.; Rozum, Yu. S.; Prokopenko, V. P.
 SOURCE: Kiev. Tekhnol. Inst. Pishch. Prom., Kiev, USSR
 Elektrokimiya (1971), 7(12), 1830-3
 CODEN: ELKXKX; ISSN: 0424-8570
 DOCUMENT TYPE: Journal
 LANGUAGE: Russian
 AB Polarog. data for 9 2,2'-biphenazines (I, R = MeO, Me, Cl; R1 = Me; R2 = CO2H, CO2CSH11, CO2Me) were related to the Hammett σ const. for the substituents.
 IT 37552-97-9
 RL: PREP (Properties)
 (polarography and ir spectrum of)
 RN 37552-97-9 CAPLUS
 CN [2,2'-Biphenazine]-7,7'-dicarboxylic acid, dipentyl ester (9CI) (CA INDEX NAME)



PAGE 1-A

PAGE 1-B


— Me

L13 ANSWER 147 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1971:465269 CAPLUS
 DOCUMENT NUMBER: 75:65269
 TITLE: Metal-containing monoazo fiber-reactive dyes
 INVENTOR(S): Jager, Horet; Schundehutte, Karl H.; Machatzke, Heinz
 PATENT ASSIGNEE(S): Farbenfabriken Bayer A.-G.
 SOURCE: U.S., 5 pp.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 3585181	A	19710615	US 1969-849600	19690807


GI For diagram(s), see printed CA issue.
 AB Hydroxy azonaphthalene Cu complex fiber reactive dyes (I, A = sulfo-1,2-naphthylenes, R = 2,3-dichloro-6-quinoxalylcarbonyl, 4,6-dichloro-8-triazin-2-yl, R1 = H, SO3H, R2 = H, SO3H, R3 = H, SO3H), useful for dyeing natural and regenerated cellulose, were prepared. Thus, 2-[7-[2-[(2,3-dichloro-6-quinoxalylcarbonyl)methylamino]acetamidomethyl]-2,8-dihydroxy-3,6-disulfo-1-naphthyl]azo]-4-sulfo-1-naphthol 1:1 copper complex (I, A = 4-sulfo-1,2-naphthylene, R = 2,3-dichloro-6-quinoxalylcarbonyl, R1 = R3 = SO3H, R2 = H) was prepared by condensing I

(A = 4-sulfo-1,2-naphthylene, R = H, R1 = R3 = SO3H, R2 = H) with 2,3-dichloroquinoline-6-carbonyl chloride at pH 7-8, and used to dye cotton fast blue shades.
 IT 16207-38-8P 16207-39-9P 16265-96-6P
 16265-97-7P 33111-13-8P
 RL: IMP (Industrial manufacture); PREP (Preparation) (preparation of)
 RN 16207-38-8 CAPLUS
 CN Copper, [trihydrogen 3-[[[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamidomethyl]-4,6-dihydroxy-5-[[1-hydroxy-4-sulfo-2-naphthyl]azo]-2,7-naphthalenedisulfonato(2-)]]- (8CI) (CA INDEX NAME)



PAGE 1-A


PAGE 2-A




● 3 H*

RN 16207-39-9 CAPLUS
 CN Copper, [trihydrogen 3-[[[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamidomethyl]-2,8-dihydroxy-4-sulfo-1-naphthyl]azo]-4-hydroxy-1,5-naphthalenedisulfonato(2-)]]- (8CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A


● 3 H⁺

RN 16265-96-6 CAPLUS
 CN Copper, [[tetrahydrogen 3-[[[2-[[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamidomethyl]-2,8-dihydroxy-6-sulfo-1-naphthyl]azo]-4-hydroxy-1,5-naphthalenedisulfonato(2-)]-(8CI) (CA INDEX NAME)]

PAGE 1-A




PAGE 2-A


● 3 H⁺

RN 16265-97-7 CAPLUS
 CN Copper, [[tetrahydrogen 6-[[[2-[[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamidomethyl]-3,3',5-trihydroxy-4,4'-azodi-2,7-naphthalenedisulfonato(2-)]-(8CI) (CA INDEX NAME)]

PAGE 1-A




PAGE 2-A


● 4 H⁺

RN 13111-15-8 CAPLUS
 CN Copper, [[dihydrogen 3-[[[2-[[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamidomethyl]-2-hydroxy-8-methanesulfonamido-1-naphthyl]azo]-4-hydroxy-1,5-naphthalenedisulfonato(2-)]-(8CI) (CA INDEX NAME)]


PAGE 1-A



PAGE 2-A

● 2 H⁺

L13 ANSWER 148 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1971:406107 CAPLUS
 DOCUMENT NUMBER: 75:4307
 TITLE: Anomalous nucleosides and related compounds. XVI. Phenazinyloxyphosphates
 AUTHOR(S): Chernetskii, V. P.; Vlasika, N. Ya.
 CORPORATE SOURCE: Inst. Mikrobiol. Virusol. im. Zabolotnogo, Kiev, USSR
 SOURCE: Khimiya Geterotsiklicheskikh Soedinenii (1970), (7), 985-90
 CODEN: KGSSAQ; ISSN: 0132-6244
 DOCUMENT TYPE: Journal
 LANGUAGE: Russian
 GI For diagram(s), see printed CA Issue.
 AB Glycine Me ester hydrochloride was treated with PCl3 in CSH5N 30 min, 1-phenazinecarboxylic acid added, and the mixture heated 3 hr at 100° to give 50% I (R = OMe, R1 = H). Similarly obtained were 6 other I analogs and II.
 IT 32271-88-8P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 RN 32271-88-8 CAPLUS
 CN Glycine, N-(2-phenazinyloxyphosphoryl)-, methyl ester (8CI) (CA INDEX NAME)




L13 ANSWER 149 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1971:133012 CAPLUS
 DOCUMENT NUMBER: 74:133012
 TITLE: Quinoxaline derivative catalysts for photographic
 silver dye bleach baths
 INVENTOR(S): Schlunke, Hans P.; Ronco, Karl
 PATENT ASSIGNER(S): CIBA Ltd.
 SOURCE: Ger. Offen., 72 pp.
 CODEN: GWXXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2010280	A	19700924	DE 1970-2010280	19700305
DE 2010280	C3	19791115		
DE 2010280	B2	19790322		
CH 508226	A	19710531	CH 1969-508226	19690313
US 3656953	A	19720418	US 1970-16207	19700303
FR 2034876	A5	19701218	FR 1970-8444	19700310
BE 747252	A	19700914	BE 1970-747252	19700312
NL 7003551	A	19700915	NL 1970-3551	19700312
NL 167523	B	19810716		
NL 167523	C	19811216		
GB 1299402	A	19721213	GB 1970-1299402	19700312
SU 363336	D	19730525	SU 1970-1416280	19700312
JP 49010054	B4	19740308	JP 1970-21168	19700313
			CH 1969-3820	A 19690313


PRIORITY APPL. INFO.:
 AB They are obtained by the condensation of properly substituted diamines with 1,2-dicarbonyl comds., α -halo ketones, or α -oximino ketones, followed by oxidation with Na m-nitrobenzenesulfonate. 43 synthesized examples are listed with their m.ps. For use 1-100 mg is added to 1 l. of bleach bath, or they are incorporated in the dye or another layer of the photog. material.

IT 32388-05-99 32388-06-0P 32388-08-2P
 32388-09-3P
 RL: PREP (Preparation)
 (manufacture of, and used in photographic silver-dye bleach process)
 RN 32388-05-9 CAPLUS
 CN 6-Quinoxalinecarboxylic acid, 2,3-diphenyl-, ethyl ester (8CI, 9CI) (CA INDEX NAME)




RN 32388-06-0 CAPLUS


CN 6-Quinoxalinecarboxylic acid, 2,3-dimethyl-, ethyl ester (8CI, 8CI, 9CI) (CA INDEX NAME)



RN 32388-08-2 CAPLUS
 CN 6-Quinoxalinecarboxamide, N,N-diethyl-2,3-dimethyl- (8CI, 9CI) (CA INDEX NAME)



RN 32388-09-3 CAPLUS
 CN 6-Quinoxalinecarboxamide, N,N-diethyl-2,3-diphenyl- (8CI, 9CI) (CA INDEX NAME)




L13 ANSWER 150 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1971:87918 CAPLUS
 DOCUMENT NUMBER: 74:87918
 TITLE: N-Oxides of N-phenazinyl derivatives of some α -amino acids
 AUTHOR(S): Batulina, R. Kh.; Pushkareva, Z. V.; Konyukhov, V. N.; Bobarykina, K. Yu.; Platonova, G. N.
 CORPORATE SOURCE: Ural. Politekh. Inst. im. Kirova, Sverdlovsk, USSR
 SOURCE: Khimiko-Farmatsveticheski Zhurnal (1970), 4(11), 18-22
 CODEN: KHFFAN; ISSN: 0033-1134
 DOCUMENT TYPE: Journal
 LANGUAGE: Russian
 GI For diagram(s), see printed CA Issue.


AB 1-Phenazinecarboxylic acid was obtained from condensation of o-H₂NCH₂COOH and PhNO₂. 2-Phenazinecarboxylic acid (I) was prepared by oxidative condensation of p-toluidine with PhNO₂. I as the acid chloride reacted with amino acid Et esters to give II. III and IV were obtained from II (R = iso-Pr) by oxidation with 30 H₂O₂ in HOAc.

IT 30806-07-2P 30806-08-3P 30806-09-4P
 30905-67-0P 30905-73-8P 30905-74-9P
 RL: SPW (Synthetic preparation); PREP (Preparation)
 (preparation of)


RN 30806-07-2 CAPLUS
 CN Glycine, N-(2-phenazinylcarbonyl)-, ethyl ester (8CI, 9CI) (CA INDEX NAME)



RN 30806-08-3 CAPLUS
 CN Glycine, N-(2-phenazinylcarbonyl)-, ethyl ester, dioxide (9CI) (CA INDEX NAME)




RN 30806-09-4 CAPLUS
 CN Glycine, N-(2-phenazinylcarbonyl)-, dioxide (9CI) (CA INDEX NAME)




RN 30905-67-0 CAPLUS
 CN L-Valine, N-(2-phenazinylcarbonyl)-, ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 30905-73-8 CAPLUS
 CN L-Valine, N-(2-phenazinylcarbonyl)-, ethyl ester, dioxide (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 30905-74-9 CAPLUS
 CN L-Valine, N-(2-phenazinylcarbonyl)-, dioxide (9CI) (CA INDEX NAME)

Absolute stereochemistry.





L13 ANSWER 151 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1970:521552 CAPLUS
 DOCUMENT NUMBER: 73:121552
 TITLE: Fiber-reactive phthalocyanine dyes
 PATENT ASSIGNER(S): Farbenfabriken Bayer A.-G.
 SOURCE: Fr. Demande, 12 pp.
 CODEN: FRXXBL
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2011515		19700306	FR 1969-19221	19690610
GB 1262583				
			GB	
			DE	19680610



PRIORITY APPL. INFO.:
 AB The title comds. (I, M = Cu or Ni, x = 3 or 4, Pc = phthalocyanine), useful for dyeing cotton fast greenish blue shades, were prepared by treating MPc (x-SO₂Cl)₃₋₄ with AcNMeCH₂CH₂NHMe, optionally adding NH₄Cl, deacetylating, and acylating with 2,3-dichloroquinoxaline-6-carbonyl chloride in the presence of Na₂CO₃. Thus, CuPc was chlorosulfonated and the resultant CuPc(3-SO₂Cl)₄ converted to turquoise blue I (M = Cu, x = 3, n = 2.5, n = 0, p = 1.5). Similarly 5 other I were prepared

IT 28901-03-9P 29116-77-6P
 RL: IMP (Industrial manufacture); PREP (Preparation)
 (preparation of)

RN 28901-03-9 CAPLUS
 CN Copper, [dihydrogen {[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)ethyl]methylsulfamoyl}phthalocyaninedisulfonato(2-)]-, disodium salt (8CI) (CA INDEX NAME)

● 2 Na⁺

RN 29116-77-6 CAPLUS
 CN Copper, (trihydrogen [[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)ethyl]methylsulfamoyl]phthalocyaninetrisulfonato(2-)]-, trisodium salt (8CI) (CA INDEX NAME)


● 3 Na⁺

L13 ANSWER 152 OF 161 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1970:100749 CAPLUS
 DOCUMENT NUMBER: 72:100749
 TITLE: Quinoxalines useful in treatment of peptic ulcers
 INVENTOR(S): Bolhofer, William A.; Baldwin, John J.
 PATENT ASSIGNER(S): Merck and Co., Inc.
 SOURCE: S. African, 25 pp.
 CODEN: SFXKAB
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
ZA 6707613		19690619		
DE 1695532			DE	
FR 1588778			FR	
FR 7331			FR	
GB 1180249			GB	
US 3510487		19700505	US	19661228
US 3655894		19720411	US	19690828
PRIORITY APPLN. INFO.:			US	19661228

OTHER SOURCE(S): MARPAT 72:100749


GI For diagram(s), see printed CA issue.
 AB The title compds. (I or II) were prepared by treating an amine with 6-(chlorocarbonyl)-2,3-dichloroquinoline (III) or III deriva. The compds. inhibit acid secretion. Thus, 0.048 mole CH₂NH₂CH₂OH in 2.3 ml. dioxane added to 0.023 mole III in 53 ml dioxane gave 6-[(2-hydroxyethyl)carbamoyl]-2,3-dichloroquinoxaline, m. 181-3° (MeCN). Similarly were prepared 6-(diethylcarbamoyl)-2,3-dichloroquinoxaline, m. 73-5°, 6-[(2-methoxyethyl)-carbamoyl]-2,3-dichloroquinoxaline, m. 141-2°, and the following I (X = H) (R, R₁, and m.p. given): CH₂CH₂OH, H, 161.5-3.5°; Me, H, 213-15°; Me, Me, 138-9°; Et, H, 191-6°; Et, Et, 73-5°; Pr, H, 158-60°; CH₂CH₂OH, H, 181-3°; CH₂CH₂OH, CH₂CH₂OH, 123-5°; CH₂CH₂OH, Me, 128-30°; CHMeCH₂OH, H, 157-9°; CH₂CHMeOH, H, 180-2°; CH₂C(Me)₂OH, H, 185-7°; CH₂CH₂CH₂OH, H, 136-8°; CH₂CH₂OMe, H, 141-2°; CH₂CH₂Cl, H, 165-6°; CH₂CH=CH₂, H, 166.0-7.5°; HOCH₂CH₂Ph, H, 217-18°; tetrahydrofurfuryl, H, 149-50°; cyclopropyl, H, 208-10°; and 11 others. 2,4,5-Cl₃(NH₂)₂C₆H₂CO₂H (5.23 g) and 210 ml (CO₂Et)₂ gave 7.22 g 2,3-dihydroxy-7-chloroquinoxaline-6-carboxylic acid (IV), m. >300°. IV (5 g) refluxed 45 min with 12.95 g POCl₃ gave 280 mg 6-(chlorocarbonyl)-2,3,7-trichloroquinoxaline (V), m. 93-5° (hexane). V (0.07 g) in 5 ml dioxane treated at 27° with 0.029 g NH₂CH₂CH₂OH in 2 ml dioxane gave 0.02 g 6-[(2-hydroxyethyl)carbamoyl]-2,3,7-trichloroquinoxaline (VI), m. 204-5° (BuCl-MeCN). 6-[(2-Dimethylaminoethyl)carbamoyl]-2,3-dichloroquinoxaline (VII) HCl salt (570 mg) in 50 ml EtOAc treated with 25 ml saturated NaHCO₃ gave 440 mg VII, m. 158-60° (BuCl). Similarly, 1 g 6-(2-morpholinoethyl)carbamoyl)-2,3-dichloroquinoxaline (VIII) HCl salt in 100 ml Et₂O gave 450 mg VIII, m. 163-6° (iso-PrOH); and 0.61 g NH₂CH₂CH₂OH, 30 ml CHCl₃, and 1.30 g 2,3-dichloro-5-(chlorocarbonyl)-2,3-dichloroquinoxaline gave II (X = R = H, R₁ = HOCH₂CH₂), m. 154.5-6.5°. NH₂CH₂CH₂NMe₂ (0.004 mole) in 5 ml dioxane added at room temperature to 0.004 mole III in 10 ml dioxane gave VII.HCl, m. 224-6°. Similarly were prepared the following I.HCl (X = H, R, R₁, and m.p. given): (CH₂)₃CH₂NMe₂, H, 178.0-9.5°; CH₂CH₂NMe₂, H, 139.0-1.5°; CH₂CH₂CH₂NMe₂, H, 228.5-30.5°; and 2 others. MeOC₆H₄CO₂H (24.2 g) in 500 ml EtOH hydrogenated over 5% Pd/C at 40 psi until 0.6 mole H had been absorbed gave 2,4,5-MeO(NH₂)₂C₆H₂CO₂H, which gave 6-(2-hydroxyethylcarbamoyl)-7-methoxy-2,3-dichloroquinoxaline.
 IT 26773-13-7P 26773-14-8P 26773-17-1P
 26773-18-2P 26773-19-3P 26773-21-7P
 26773-22-8P 26773-25-1P 26773-26-2P
 26773-27-3P 26773-28-4P 26773-29-5P
 26773-30-8P 26773-31-9P 26773-32-0P
 26840-63-1P 26840-68-4P 26840-73-3P
 26840-74-4P 26840-75-5P 26840-76-6P
 26840-77-7P 26840-78-8P 26887-34-3P
 26887-35-4P 26921-20-0P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 RN 26773-13-7 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-hydroxyethyl)- (8CI, 9CI) (CA INDEX NAME)




RN 26773-14-8 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-methoxyethyl)- (8CI) (CA INDEX NAME)




RN 26773-17-1 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3,7-trichloro-N-(2-hydroxyethyl)- (8CI) (CA INDEX NAME)




RN 26773-18-2 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-[2-(dimethylamino)ethyl]- (8CI) (CA INDEX NAME)



RN 26773-19-3 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-morpholinoethyl)- (8CI) (CA INDEX NAME)




RN 26773-21-7 CAPLUS
 CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-[2-(dimethylamino)ethyl]-, hydrochloride (8CI) (CA INDEX NAME)




● x HCl


RN 26773-22-8 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-hydroxyethyl)-, acetate (ester) (8CI) (CA INDEX NAME)




RN 26773-35-1 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-ethyl- (8CI, 9CI) (CA INDEX NAME)




RN 26773-26-2 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-bis(2-hydroxyethyl)- (8CI) (CA INDEX NAME)




RN 26773-27-3 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-hydroxyethyl)-N-methyl- (8CI) (CA INDEX NAME)




RN 26773-38-4 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-hydroxy-1-methylethyl)- (8CI) (CA INDEX NAME)




RN 26773-29-5 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-hydroxypropyl)- (8CI) (CA INDEX NAME)




RN 26773-30-8 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-hydroxy-2-methylpropyl)- (8CI) (CA INDEX NAME)




RN 26773-31-9 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(3-hydroxypropyl)- (8CI) (CA INDEX NAME)




RN 26773-32-0 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-chloroethyl)- (8CI, 9CI) (CA INDEX NAME)




RN 26840-63-1 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(tetrahydrofurfuryl)- (8CI) (CA INDEX NAME)




RN 26840-68-6 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2,3-dihydroxypropyl)- (8CI) (CA INDEX NAME)




RN 26840-73-3 CAPLUS
CN 6-Quinoxalinecarboxamide, N-(2-acetamidoethyl)-2,3-dichloro- (8CI) (CA INDEX NAME)



RN 26840-74-4 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(β-hydroxyphenethyl)- (8CI) (CA INDEX NAME)




RN 26840-75-5 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(4-(diethylamino)butyl)-, monohydrochloride (8CI) (CA INDEX NAME)




● HCl

RN 26840-76-6 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-[2-(diethylamino)ethyl]-, monohydrochloride (8CI) (CA INDEX NAME)



● HCl


RN 26840-77-7 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-[3-(dimethylamino)propyl]-, monohydrochloride (8CI) (CA INDEX NAME)



● HCl

RN 26840-78-8 CAPLUS


CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-(2-morpholinoethyl)-, monohydrochloride (8CI) (CA INDEX NAME)



● HCl


RN 26887-34-3 CAPLUS

CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N,N-diethyl- (8CI, 9CI) (CA INDEX NAME)




RN 26887-35-4 CAPLUS

CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-propyl- (8CI) (CA INDEX NAME)



RN 26921-20-0 CAPLUS

CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-2-propenyl- (9CI) (CA INDEX NAME)



L13 ANSWER 153 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 1970:70530 CAPLUS

DOCUMENT NUMBER: 72:70530

TITLE: Preparation and laboratory evaluation of cellulose-based ion permselective membranes

Sueszer, A.; Bandel, E.; Flitman, M.

CORPORATE SOURCE: Negev Inst. Arid Zone Res., Beer-Sheva, Israel

SOURCE: Desalination (1969), 7(1), 47-50

CODEN: DSIJAH; ISSN: 0011-9164

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Ion permselective membranes were prepared by treating cellophane membranes with trichloropyrimidine reactive dyes, i.e. Drimarene Black Z-BL or Reaction Yellow RL, or dichloroquinoxaline reactive dyes, i.e. N-(3-dimethylaminopropyl)-2,3-dichloro-6-quinoxalinecarboxamide. Promising results were obtained when electro-dialytic desalination stacks prepared with the membranes were used in the desalination of water.


IT 24604-56-6

RL: OCCU (Occurrence)

(reaction products, with cellophane)

RN 24604-56-6 CAPLUS

CN 6-Quinoxalinecarboxamide, 2,3-dichloro-N-[3-(dimethylamino)propyl]- (8CI) (CA INDEX NAME)



L13 ANSWER 154 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 1969:514152 CAPLUS

DOCUMENT NUMBER: 71:114152

TITLE: Reactive dyes

PATENT ASSIGNEE(S): Imperial Chemical Industries Ltd.

SOURCE: Fr., 6 pp.

CODEN: FRXXAK

DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 1531271		19680628	FR 1967-114662	19670718
DE 1619594				
GB 1130395				
GB 1180395				
US 3489502		19700113		19670712
PRIORITY APPLN. INFO.:				19660718
				19670619

GI For diagram(s), see printed CA Issue.

AB Comps. of the general formula I, where Y is an alkylene or arylene radical and Z is a heterocyclic group, dye cellulose textiles reddish blue. Thus, a solution of 5.16 parts I (Y = CH₂CH₂, Z = H) (II), and 0.6 part NaOH in 145 parts H₂O and 20 parts Me₂CO was added during 15 min. with stirring to a suspension of 3.7 parts cyanuric chloride (III) in 25 parts Me₂CO and 75 parts H₂O and ice at 0-3° (pH held at 8.5-9 with 2N NaOH), and stirred for 1.25 hrs. to give I (Y = CH₂CH₂, Z = 4,6-dichloro-s-triazin-2-yl (Q)) (IV). A solution of 6.88 parts di-Na salt of I (Y = 2,5-C₆H₃SO₃H(R), Z = H) in 130 parts H₂O was added during 20 min. with stirring to a suspension of 1.94 parts III in 75 parts H₂O and ice and 12 parts Me₂CO at 0-5° (pH held at 5-5.5 with 2N NaOH), stirred for 45 min. at 0-5° and pH 5 ± 0.2, and sieved to eliminate excess III to give the di-Na salt of I (Y = R, Z = O) (V). The solns. of IV and V were mixed, the pH adjusted to 6.4, a solution of 12 parts 3-Et₃NC₆H₄SO₃-Na (VI) and 0.75 part NaHSO₄ in 40 parts H₂O added with

stirring followed by 10 parts NaCl, the precipitated dye filtered, washed with

a solution of 24 parts NaCl, 0.37 part NaHSO₄, and 6 parts VI in 240 parts H₂O, and dried at room temperature. Similarly, other dyes were prepared (reactants given): II, I (Y = CH₂CHMe, Z = H) (VII), III; II, VII, 5-cyano-2,4,6-trichloropyrimidine: II, VII, 2,3-dichloro-6-quinoxalinecarboxyl chloride; I (Y = CH₂CH₂, Z = CH₂CH₂OR), III, mixed with IV.


IT 25238-46-4P 25238-47-5P

RL: IMF (Industrial manufacture); PREP (Preparation)


(preparation of)

RN 25238-46-4 CAPLUS

CN 2-Anthracenesulfonic acid, 1-amino-4-[m-[[2-(2,3-dichloro-6-quinoxalinecarboxamido)ethyl]sulfamoyl]anilino]-9,10-dihydro-9,10-dioxo- (8CI) (CA INDEX NAME)




PAGE 1-A




RN 25238-47-5 CAPLUS

CN 2-Anthracenesulfonic acid, 1-amino-4-[m-[[2-(2,3-dichloro-6-quinoxalinecarboxamido)ethyl]sulfamoyl]anilino]-9,10-dihydro-9,10-dioxo- (8CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A



L13 ANSWER 155 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN

ACCESSION NUMBER: 1969:492648 CAPLUS

DOCUMENT NUMBER: 71:92648

TITLE: Anthraquinone fiber-reactive dyes

INVENTOR(S): Harms, Wolfgang; Gehrke, Gunter; Hohmann, Walter;

Bien, Hans S.

PATENT ASSIGNEE(S): Farbenfabriken Bayer A.-G.

SOURCE: Brit., 25 pp.

CODEN: BRXXAA

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
GB 1147297		19690402	GB 1967-54744	19671201
DE 1644612				
FR 1551267				
PRIORITY APPLN. INFO.:				19661207

GI For diagram(s), see printed CA Issue.

AB Comps. of the general formula I where X is a group containing at least one reactive halogen, are water soluble yellow-green to green reactive dyes for


cotton. I are prepared by reacting the appropriate 5-aminoanthraquinone with a reactant containing at least 2 reactive halogen atoms. Thus 13.2 parts I (X = H, Y = OH) in 600 parts H₂O at 50 is treated with 8 parts 2,3-dichloro-6-quinoxalincarboxamide (QCOCl) and 0.5 part PhCl and the pH maintained at 5.5-6.5 by dropwise addition of N NaOH. After 1 hr. PhCl is removed by aeration, the mixture clarified and salted to precipitate I

(R = H, X = QCO) a yellowish green dye. Similarly other yellowish green I (Y = OH) are prepared (R and X given): H, QSO₂, 2-methoxysulfonyl-4-methyl-5-chloro-6-pyrimidinyl (Z); H, 4-ZNHC6H4CO; H, 3-ZNHC6H4SO₂; H, 2,4-difluoro-5-chloro-6-pyrimidinyl; H, ClCH₂CO; H, MeNHCH₂CO; H, QSO₂NMeCH₂CO; H, QCONMeCH₂CO; H, BrCH₂CHBrCO; H, 1,4-dichloro-6-phthalazinecarbonyl; H, 4,6-dichloro-s-triazin-2-yl(Q'); H, 2,4,5-trichloro-6-pyrimidinyl; H, 4-chloro-6-methoxy-s-triazin-2-yl. Similarly other green I were prepared (R, X, and Y given): H, QCO, NH₂; H, QSO₂, NH₂; H, QCO, Cl; H, QSO₂, Cl; H, Q', Cl; H, QCO, Br; H, QCO, NHCH₃(SO₃Na)Me; H, QSO₂, NHCH₃(SO₃Na)Me. Also prepared were green II (X = CO) and blue-green II (X = SO₂).

IT 23945-98-49 23946-02-39
RL: IMP (Industrial manufacture); PREP (Preparation)

RN 23945-98-4 CAPLUS
CN m-Toluenesulfonic acid, 6,6'-[[5-[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]-8-hydroxy-1,4-anthraquinonylene]diimino]di-, disodium salt (8CI) (CA INDEX NAME)


PAGE 1-A



● 2 Na

RN 23946-02-3 CAPLUS
CN m-Toluenesulfonic acid, 6,6'-[[5-[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]-8-hydroxy-1,4-anthraquinonylene]diimino]di-, disodium salt (8CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A



● 2 Na

L13 ANSWER 156 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1969:482641 CAPLUS
DOCUMENT NUMBER: 71:82641

TITLE: Anthraquinone fiber-reactive dyes
INVENTOR(S): Bien, Hans-S.; Harms, Wolfgang; Schmitz, Reinold;
Leister, Heinrich
PATENT ASSIGNER(S): Farbenfabriken Bayer A.-G.
SOURCE: Brit., 23 pp.
CODEN: BRXXAA

DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
GB 1147110		19690402	GB 1967-53799	19671127
DE 1644611			DE	
FR 1546177			FR	
US 3828040		19740806	US 1970-64007	19700717
			DE	19661130

PRIORITY APPLN. INFO.:

Q1 For diagram(s), see printed CA issue.

AB 1, where X is a group containing a reactive halogen, and II, where R is sulfonyl or sulfato, are water-soluble yellowish green to bluish green dyes for cotton. I are prepared by reacting the appropriate 5-amino-1,4-bisaminoanthraquinone with a reactant containing 22 reactive halogen atoms. A mixture of 15 parts 5-aminoquinizarin, 100 parts p-MeC₆H₄NH₂, 8 parts H₂SO₄, and 8.3 parts concentrated HCl is treated at 75° with 2 parts Zn dust during 1 hr., heated at 90-95° 2.5 hrs., treated at 95° with 15 parts powdered KOH, the melt aerated, cooled to 65°, and treated with 100 parts MeOH to give I (X = R = H) (III), green needles. III (20 parts) is added during 3 hrs. to a mixture of 48 parts 20% oleum and 52 parts concentrated H₂SO₄. The mixture is treated with 34 parts 20% oleum, stirred until disulfonation is complete, and treated with 300 parts ice to precipitate I (X = H, R = SO₃H) (IV). A solution of 8


parts Na salt of IV in 150 parts H₂O is heated at 80-85° with 7 parts 2,4,5,6-tetrachloropyrimidine (V) for 12 hrs. while the pH is kept at 5.5-7.0 by dropwise addition of N aqueous NaOH, excess V boiled off, and the green dye I (X = 2,4,5-trichloro-6-pyrimidinyl, R = SO₃Na) (Va) is salted out. The following I (R = SO₃Na) are similarly prepared from IV (X and shade on cotton given): 4-chloro-6-methoxy-s-triazin-2-yl, green; 2,3-dichloro-6-quinoxa-lylcarbonyl, green; 2,3-dichloro-6-quinoxalinesulfonyl, bluish green; 4,6-dichloro-s-triazin-2-yl, green; 1,4-dichloro-6-phthalazincarbonyl, green; (2,3-dichloro-N-methyl-6-quinoxaline-sulfonamido)acetyl, green; ClCH₂CO, - (nylon, green); 2-(methylsulfonyl)-6-benzothiazolylsulfonyl, green; BrCH₂CHBrCO, - (wool, green). The following intermediates (I, R = SO₃Na) were similarly prepared (X given): p-O₂NCH₄CO (VI); m-O₂NCH₄CO₂ (VII). A solution of VI (from 55 parts Na salt of IV) in 850 parts H₂O is treated at 70-75° for 30 min. with 17.2 parts Me₂S.H₂O and 11 parts NaHCO₃ in 180 parts H₂O and salted to precipitate I (X = p-H₂NCH₄CO, R = SO₃Na). Similarly, VII is converted to I (X = m-H₂NCH₄CO₂, R = SO₃Na). In a similar manner to that described above, the following I (R = SO₃Na) are prepared (X and shade on cotton given): p-(2,3-dichloro-6-quinoxalinesulfonamido)benzoyl, green; p-(2,3-dichloro-6-quinoxalinesulfonamido)phenylsulfonyl, bluish green; p-(5-chloro-6-methyl-2-(methylsulfonyl)-4-pyrimidinylamino)-benzoyl, green; p-(5-chloro-6-methyl-2-(methylsulfonyl)-4-pyrimidinylamino)phenylsulfonyl, bluish green; m-(2,3-dichloro-6-quinoxalinesulfonamido)phenylsulfonyl, bluish green; p-(5-chloro-2,6-difluoro-4-pyrimidinylamino)benzoyl, fast green. Similarly, as was Va, were prepared the following I (R and shade on cotton given): C₆H₄SO₃K, blue; OSO₃Na (VIII), blue. Also prepared was the 2,3-dichloro-6-

quinoxalinesulfonamido analog of VIII, greenish blue on cotton.


IT 24031-65-0P
RL: IMP (Industrial manufacture); PREP (Preparation)

RN 24031-65-0 CAPLUS
CN m-Toluenesulfonic acid, 6,6'-[[5-[2-(2,3-dichloro-N-methyl-6-quinoxalinesulfonamido)acetamido]-1,4-anthraquinonylene]diimino]di- (8CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A




L13 ANSWER 157 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1969:38891 CAPLUS
DOCUMENT NUMBER: 70:38891
TITLE: Fiber reactive azo dyes
INVENTOR(S): Jaeger, Horst; Schuendehutte, Karl H.
PATENT ASSIGNER(S): Farbenfabriken Bayer A.-G.
SOURCE: Brit., 10 pp.
CODEN: BRXXAA
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE

GB 1130228 19681009 GB 1967-40454 19670905
DE 1644170 DE
FR 1538115 FR DE 19660910


PRIORITY APPLN. INFO.:
GI For diagram(s), see printed CA issue.
AB The moist paste of the monoozo compound obtained by diazotizing 31.3 parts 4,3-H₂N(HO₃S)C₆H₃CH₂NHMe and coupling with 46.1 parts 1,5,4,7-(HO₃S)2C₁₀H₄(NH₂)₂ is slurried in 500 part H₂O, treated with 200 parts 20% NH₄OH and 80 parts CuSO₄.5H₂O, the mixture heated at 90-5° until the red color has disappeared, treated with Na₂S to precipitate Cu₂S and the solution of the resultant triazole (I, QNH₂) diazotized and coupled (acid) with 34.7 parts 1,3,7-HO(HO₃S)C₁₀H₅NH₂. The moist paste of the azotriazole derivative is stirred into 1000 parts H₂O, treated with 12 parts NaNO₂, poured into a solution of 12.5 parts concentrated H₂SO₄ in 200 parts ice-water, and stirred overnight, and salted to give 6,2,8,1-HO₃S-(HO)2C₁₀H₄N: HQ. The moist cake is dissolved in 500 parts H₂O at 40°, treated with 40 parts 2,3-dichloroquinoxaline-6-carboxylic acid chloride, maintained at pH 5.7 for 10 hrs. by addition of Na₂CO₃, added to 1000 parts H₂O, clarified with charcoal, and salted. The moist cake is suspended in 4000 parts ice-water and added to a solution of 40 parts CuSO₄.5H₂O in 400 parts H₂O, kept at pH 7 by addition of aqueous NaOH, and the dark red solution mixed dropwise with 250 parts 3% H₂O₂ (turns blue) and salted to give II, a clear blue, fiber reactive dye.
IT 22873-61-6P
RL: IMP (Industrial manufacture); PRP (Preparation) (preparation of)
RN 22873-61-6 CAPLUS
CN Copper, [tetrahydrogen 2-[α-(2,3-dichloro-N-methyl-6-quinoxalinedicarboxamido)-6-sulfo-m-tolyl]-7-[(2,8-dihydroxy-6-sulfo-1-naphthyl)azo]-6-hydroxy-2H-naphtho[1,2-d]triazole-5,9-disulfonate(2-)]-(8CI) (CA INDEX NAME)

PAGE 1-A




IT 19625-39-9P
RL: FORM (Formation, nonpreparative); PRP (Preparation) (formation of)
RN 19625-39-9 CAPLUS
CN Copper, [μ-(dihydrogen 5,8-dihydroxy-6,7-quinoxalinedicarboxylato(2-))]bis(dihydrogen 5,8-dihydroxy-6,7-quinoxalinedicarboxylato)di-, hexaethyl ester (8CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A



● 4 H⁺


L13 ANSWER 158 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1968:413368 CAPLUS
DOCUMENT NUMBER: 69-13368
TITLE: Chelating reagents containing N-heterocycles. V. Dihydroxyquinoxaline studies. Solubility, ionization constant, and chelating behavior
AUTHOR(S): Oguchi, Shoshichi
CORPORATE SOURCE: Tokyo Gakugei Univ., Tokyo, Japan
SOURCE: Bulletin of the Chemical Society of Japan (1968), 41(4), 980-7
CODEN: BCSJAB; ISSN: 0009-2673
DOCUMENT TYPE: Journal
LANGUAGE: English
AB The solubility, ionization constant, and chelating behavior of dihydroxyquinoxaline, mainly 5,8-dihydroxyquinoxaline (I) derivs., were measured. The introduction of a hydroxyl group into quinoxaline greatly lowers the solubility. The order of the decreasing solubility of 2,3-disubstituted I in H₂O (at 20°) is: O(CH₂)₂-OEt < H < 1H < Me < Ph < EtO < Cl < SH < Et. For I the introduction of electron-repelling groups into 2,3-positions raises both the pK_{OH} and pK_{NH} values, while the introduction of electron-attracting groups into the same positions lowers both pK_{NH} and pK_{OH}. I and its derivs. form colored precipitate with metal ions, but some derivs. which have pK_{NH} values lower than zero fail to show any precipitation or coloration. 6,7-Dihydroxyquinoxaline forms precipitate, and the precipitation colors distinctively with Cr(III) or Fe(II). The composition and stability constant (K) of Cu(II) chelates of I and its derivs. in a dioxane-H₂O (10:90 by volume) solution were studied spectrometrically. The Cu chelate of I had a metal:ligand ratio of 1:1 and a log K value of 6.28 at pH 4.0. 27 references.
IT 2427-91-0
RL: PRP (Properties) (chelation properties and ionization and solubility of)
RN 2427-91-0 CAPLUS
CN 6,7-Quinoxalinedicarboxylic acid, 5,8-dihydroxy-, diethyl ester (7CI, 8CI) (CA INDEX NAME)




PAGE 3-A



IT 2427-91-ODP, 6,7-Quinoxalinedicarboxylic acid, 5,8-dihydroxy-, diethyl ester, copper complex
RL: SPN (Synthetic preparation); PRP (Preparation) (preparation of)
RN 2427-91-0 CAPLUS
CN 6,7-Quinoxalinedicarboxylic acid, 5,8-dihydroxy-, diethyl ester (7CI, 8CI) (CA INDEX NAME)



L13 ANSWER 159 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1968:21412 CAPLUS
DOCUMENT NUMBER: 68-21412
TITLE: Proton resonance spectra of heterocycles. IV. Quinoxaline and monosubstituted quinoxalines
AUTHOR(S): Brignell, Peter J.; Katritzky, Alan R.; Reavill, Roger E.; Cheeseman, Gordon W. H.; Sarsfield, A. A.
CORPORATE SOURCE: Univ. East Anglia, Norwich, UK
SOURCE: Journal of the Chemical Society (Section) B: Physical Organic (1967), (11), 1241-3
CODEN: JCSPEC; ISSN: 0045-6470
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Chemical shifts and coupling consts. are reported for fourteen monosubstituted quinoxalines. These parameters are correlated with the effect of the substituents on the electron distribution. 16 references.
IT 6924-72-7
RL: PRP (Properties) (nuclear magnetic resonance of)
RN 6924-72-7 CAPLUS
CN 6-Quinoxalinedicarboxylic acid, ethyl ester (6CI, 7CI, 8CI, 9CI) (CA INDEX NAME)



L13 ANSWER 160 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1967:500993 CAPLUS
DOCUMENT NUMBER: 67:100993
TITLE: Metallized azo dyes containing 2,3-dichloroquinoline-6-carboxylamino groups
INVENTOR(S): Jaeger, Horst; Gerlach, Klaus
PATENT ASSIGNEE(S): Farbenfabriken Bayer A.-G.
SOURCE: Fr., 9 pp.
CODEN: FRXXAK
DOCUMENT TYPE: Patent
LANGUAGE: French
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 1470128		19670217	FR 1966-51204	19660225
			DE	19650226

PRIORITY APPLN. INFO.:

OI For diagram(s), see printed CA Issue.

AB 1, II, and III are blue dyes for cotton. Thus, 27.6 parts 2,3,5-HO(HO3S)(MeNHCH2CONH)C6H2NH2 is diazotized and coupled with 34.1 parts 8,3,6,1-H2N(NaO3S)2C10H4OH, to give an azo dye which is dissolved in 1000 parts water at 45°, treated with a solution of 24.9 parts CuSO4 at pH 5-6 (Na2CO3), and treated at 45° with 26.1 part 2,3-dichloroquinoline-6-carboxylic acid chloride (OCl) at pH 5-6 (Na2CO3) to give I, dark powder, which gives a violet-blue aqueous solution and reddish blue shades on cotton. Also prepared are (color on cotton given): II, greenish blue; III, 1:2 Cr complex, black; III, 1:3 Co complex, black.

IT 16246-84-7P 16520-32-4P


RL: IMF (Industrial manufacture); PREP (Preparation)

(preparation of)

16246-84-7 CAPLUS


CN Copper, [trihydrogen 4-amino-6-[[5-[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]-2-hydroxy-3-sulfonyl]azo]-5-hydroxy-1,3-naphthalenedisulfonato(2-)]- (8CI) (CA INDEX NAME)

PAGE 1-A



● 3 H⁺


PAGE 1-B



RN 16520-32-4 CAPLUS


CN Copper, [trihydrogen 4-amino-6-[[6-[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]-2-hydroxy-4-sulfo-1-naphthyl]azo]-5-hydroxy-1,3-naphthalenedisulfonato(2-)]- (8CI) (CA INDEX NAME)

PAGE 1-A



● 3 H⁺

PAGE 1-B



L13 ANSWER 161 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1967:482944 CAPLUS

DOCUMENT NUMBER: 67:82944

TITLE: Fiber-reactive dyes

INVENTOR(S): Jaeger, Horst; Schuendehutte, Karl H.; Machatzke, Heinz

PATENT ASSIGNER(S): Farbenfabriken Bayer A.-G.

SOURCE: Fr., 7 pp.

CODEN: FRXXAK

DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 1474432		19670324	FR 1966-56034	19660401
DE 1544516			DE	
GB 1106023			GB	
			DD	19650403

PRIORITY APPLN. INFO.:

OI For diagram(s), see printed CA Issue.

AB Blue monazo dyes containing the reactive group O are prepared by treatment of a

copperized azo dye with ClCH2CONHCH2OH (I) followed by MeNH2 and finally with 2,3-dichloroquinoline-6-carboxylic acid chloride (II). For example, a solution of 63.2 parts Cu complex of 2,8,3,6,1-(HO)2(HO3S)2C10H3N.NC1OHS(OH)SO3H-2,1,4 in 450 parts ice-cold 96% H2SO4 is stirred in an ice bath, treated with 18.5 parts finely ground I, stirred for 12 hrs. at 10-15°, and poured into 1500 parts of ice. The product is precipitated by addition of 75 parts NaCl, filtered, redissolved in 1000

parts water, adjusted to pH 7, and reprecipitated with NaCl. The wet paste is stirred with 100 parts 25% MeNH2 and 100 parts water for 24-48 hrs. at ambient temperature, adjusted to pH 5 with concentrated HCl, and filtered. A solution of

0.1 mole of this product in 400 parts water at 30-40° at pH 7 is stirred and treated with a suspension of 28 parts II together with a solution of Na2CO3 to keep pH 7-8, precipitated with KCl, filtered, and dried to give

III (W = Y = SO3H, X = Z = H, R = OH), dyeing blue shades. Similarly, III (Z = SO3H) are prepared (R, W, X, and Y given): OH, H, SO3H, H; MeSO2NH, H, H, H; OH, H, H, SO3H. IV is prepared similarly.

IT 16207-38-8P 16207-39-9P 16265-96-6P

16265-97-7P 31111-15-8P. Copper, [dihydrogen 3-[[[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]methyl]-

2-hydroxy-8-methanesulfonamido-1-naphthyl]azo]-4-hydroxy-1,5-naphthalenedisulfonato(2-)]- (8CI) (CA INDEX NAME)


RL: IMF (Industrial manufacture); PREP (Preparation)

(preparation of)

16207-38-8 CAPLUS


CN Copper, [trihydrogen 3-[[[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]methyl]-4,6-dihydroxy-5-[[1-hydroxy-4-sulfo-2-naphthyl]azo]-2,7-naphthalenedisulfonato(2-)]- (8CI) (CA INDEX NAME)

PAGE 1-A





● 3 H⁺

PAGE 2-A





RN 16207-39-9 CAPLUS



CN Copper, [trihydrogen 3-[[[7-[[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]methyl]-2,8-dihydroxy-4-sulfo-1-naphthyl]azo]-4-hydroxy-1,5-naphthalenedisulfonato(2-)]- (8CI) (CA INDEX NAME)

●3 H⁺



RN 16265-96-6 CAPLUS
 CN Copper, [[trihydrogen 3-[[[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]methyl]-2,6-dihydroxy-6-sulfo-1-naphthyl]azo]-4-hydroxy-1,5-naphthalenedisulfonato(2-)]- (8CI) (CA INDEX NAME)

●3 H⁺

RN 16265-97-7 CAPLUS
 CN Copper, [[tetrahydrogen 6-[[[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]methyl]-3,3',5-trihydroxy-4,4'-azodi-2,7-naphthalenedisulfonato(2-)]- (8CI) (CA INDEX NAME)

●4 H⁺

RN 33111-15-8 CAPLUS
 CN Copper, [[dihydrogen 3-[[[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]methyl]-2-hydroxy-6-methanesulfonamido-1-naphthyl]azo]-4-hydroxy-1,5-naphthalenedisulfonato(2-)]- (8CI) (CA INDEX NAME)

●2 H⁺

L13 ANSWER 162 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1967:455160 CAPLUS
 DOCUMENT NUMBER: 67:55160
 TITLE: Stabilized fiber-reactive dyes
 INVENTOR(S): Kissa, Erik
 PATENT ASSIGNER(S): du Pont de Nemours, E. I., and Co.
 SOURCE: U.S., 9 pp.
 CODEM: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 3313797		19670411	US 1963-252056	19630117

GI For diagram(s), see printed CA Issue.
 AB 2,3-Dichloroquinoxaline dyes for cotton were prepared and stabilized against hydrolysis by formulation with Na2SO4 or K2SO4. Thus, 36 parts 2,3-dichloro-6-quinoxalinecarboxamido chloride (QC1) was added to a solution of 42 parts 4,8,2-(HO3S)2C10H5NH2 + 3-MeC6H4NH2 in 700 parts H2O at 40° and pH 7.5-8, the mixture stirred overnight and salted with 10 parts Na2SO4 to give a light yellow powder containing .apprx.82% I, 11% H2O, and 7% Na2SO4. Similarly, other amino dyes were acylated with QC1 and the resultant amides salted (or blended) with Na2SO4 or K2SO4 (amino dye used and shade of product on cotton given): 4-(2-sulfo-5-aminophenylazo) derivative of 1-(2,5-dichloro-4-sulfo-phenyl)-3-methyl-5-pyrazolone (II), greenish yellow [also prepared by coupling 2,4-H2N(QNH)C6H3SO3H with II]; 1:1 Cu

complex of 2,5,7,6-H₂N(HO) (HO₃S)C₁₀H₄N:NC₆H₃(OH)SO₃H-2,5, rubine; III, blue; CuPc(SO₃Na)₂.5(SO₂NH₂)₂.1.4SO₂NH₂CH₃(NH₂)SO₃Na-3,4 (Pc = phthalocyanine), turquoise; 3,4-AcNH[2,4-HO₃S(4-HO₃SC₆H₄N:N)C₆H₃NH₂]. A similar dye was prepared by acylating 1,8,3,6,7-H₂N(HO) (NaO₃S)C₁₀H₃N:NC₆H₄SO₃Na-2 with 2-chloro-6-quinoxalinecarboxylic acid and salting with Na₂SO₄.

IT 16014-03-2P
RL: IMP (Industrial manufacture); PRSP (Preparation)
(preparation of)

RN 16014-03-2 CAPLUS
CN 2-Anthracenesulfonic acid, 1-amino-4-[4-(2,3-dichloro-N-ethyl-6-quinoxalinecarboxamido)-3-sulfoenilino]-9,10-dihydro-9,10-dioxo-, disodium salt (8CI) (CA INDEX NAME)



L13 ANSWER 163 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1967:86597 CAPLUS
DOCUMENT NUMBER: 66:86597
TITLE: Reactive azo dyes
INVENTOR(S): Siegel, Edgar; Sassen, Klaus
PATENT ASSIGNER(S): Farbenfabriken Bayer A.-G.
SOURCE: Ger., 6 pp.
CODEN: GWXXAW
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 1233519		19670202	DE	19610307

AB Azo dyes containing a 6-substituted-2,3-dichloroquinoxaline group (Q) and useful for dyeing cellulose, wool, polyamide, and polyester fibers were prepared. Thus, 34.7 parts 2,4,8-H₂NClO₅(SO₃Na)₂ was diazotized and coupled with 10.7 parts 3-MeC₆H₄NH₂, the aminoazo dye salted, filtered, washed, dissolved at pH 7 in 700 parts H₂O by adding NaOH, stirred at pH 7-7.5 with a solution of 25 parts QNCO in C₆H₆, salted, and filtered to give a fast reddish yellow dye for cotton. Similarly, the following dyes were prepared (reactants and shade on fiber given): QNH₂ + 1-(2,5-dichloro-4-sulfoenyl)-3-methyl-5-pyrazolone, reddish-yellow on cellulose; QNH₂ + PhN(CH₂CH₂OH)₂, orange on polyester and polyamide fibers; QNH₂ + PhOH, reddish yellow on polyamide fibers; 2-HO₃SC₆H₄NH₂ + 1,3,6,9-HO(HO₃S)C₁₀H₄NHCOCH₂CH₂NMeCO₂, bluish red on cellulose, wool, and

polyamide fibers; 1,2,3,6,8,7-HO(QH:N) (HO₃S)C₁₀H₂N:NC₆H₃(SO₃H)NH₂-2,4, QNCO, greenish blue on cellulose.

IT 14573-57-0P
RL: IMP (Industrial manufacture); PRSP (Preparation)
(preparation of)

RN 14573-57-0 CAPLUS
CN 2,7-Naphthalenedisulfonic acid, 5-[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]-4-hydroxy-3-[(o-sulfonyl)azo]- (7CI, 8CI) (CA INDEX NAME)



L13 ANSWER 164 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1967:37957 CAPLUS
DOCUMENT NUMBER: 66:37957
TITLE: Phenazine derivatives
PATENT ASSIGNEE(S): Rhone-Poulenc S. A.
SOURCE: Meth. Appl., 10 pp.
CODEN: NAXXAN
DOCUMENT TYPE: Patent
LANGUAGE: Dutch
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
NL 6603503		19660926	NL 1966-3503	19660317
FR 1462194			FR	
FR 89671			FR	
GB 1068985			GB	
US 1455926		19690715	US	19660318
			FR	19650325
			FR	19660204

PRIORITY APPL. INFO.:


GI For diagram(s), see printed CA Issue.
AB The title comds. (I, X = NH₂ or substituted-amino or N-heterocyclic moiety) are prepared by reaction of phenazine-2-carboxylic acid 5,10-dioxide (II) with N,N'-carbonyldiimidazole (III) to give I (X = 2-imidazolyl), which then is treated with an NH₂ derivative. Thus, to 17.2 g. III (88%) in 500 cc. dry HCO₂Me₂ (IV) 12 g. II is added, the mixture kept 24 hrs. at normal temperature, 21 g. 1-methylpiperazine added, and after 4 hrs. the mixture cooled to 5° to obtain 11.4 g. I (X = 4-methyl-1-piperazinyl), m. 209-10°. Similarly the following I were obtained [X and (m.p.) given]: X = 2-dimethylaminoethylamino (180°); 4-ethyl-1-piperazinyl (160-2°); 3-dimethylamino-2-propylamino (164-6°); 4-benzyl-1-piperazinyl (162°); [2-(1-pyrrolidinyl)ethylamino] (184-6°); 3-(dimethylamino)-propylamino (161-3°); [2-(4-methyl-1-piperazinyl)ethylamino] (170-2°). The comds. and their salts or quaternary N derivs. are anticancer agents.

IT 13458-25-8P 13458-27-0P 13458-29-2P
13458-30-5P 14559-63-8P
RL: SPW (Synthetic preparation); PRSP (Preparation)


(preparation of)
RN 13458-25-8 CAPLUS
CN 2-Phenazinecarboxamide, N-[2-(dimethylamino)ethyl]-, 5,10-dioxide (8CI, 9CI) (CA INDEX NAME)




RN 13458-27-0 CAPLUS
CN 2-Phenazinecarboxamide, N-[3-(dimethylamino)-2-methylpropyl]-, 5,10-dioxide (8CI, 9CI) (CA INDEX NAME)




RN 13458-29-2 CAPLUS
CN 2-Phenazinecarboxamide, N-[2-(1-pyrrolidinyl)ethyl]-, 5,10-dioxide (8CI, 9CI) (CA INDEX NAME)



RN 13458-30-5 CAPLUS
CN 2-Phenazinecarboxamide, N-[3-(dimethylamino)propyl]-, 5,10-dioxide (8CI, 9CI) (CA INDEX NAME)



RN 14559-63-8 CAPLUS
CN 2-Phenazinecarboxamide, N-[2-(4-methyl-1-piperazinyl)ethyl]-, 5,10-dioxide (8CI, 9CI) (CA INDEX NAME)




L13 ANSWER 165 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1966:51469 CAPLUS
DOCUMENT NUMBER: 64:51469
ORIGINAL REFERENCE NO.: 64:9569d-f
TITLE: Structure vs. reactivity in quinoxalinecarboxylic acids and esters
AUTHOR(S): Gurn, Wilson F., Jr.; Joulie, Madeleine M.
CORPORATE SOURCE: Univ. of Pennsylvania, Philadelphia
SOURCE: Journal of Organic Chemistry (1965), 30(11), 3982-5
CODEN: JOCEAH; ISSN: 0022-3263
DOCUMENT TYPE: Journal
LANGUAGE: English

AB In an attempt to establish a correlation between the calculated electron densities in an unperturbed quinoxaline nucleus and the reactivities of its deriva., the pKa values of 2-, 2,3-, 6-, 5-, and 2,3-dimethyl-5-quinoxalinecarboxylic acids (I, II, III, IV, V) were measured. The carbonyl frequencies of the corresponding Me and Et esters (VI-XII) were determined by ir spectroscopy and tabulated together with those of Et and Me pyrazinecarboxylate. Good correlation seemed to exist between pKa values of the acids I-V and the electron d. calculated by Longuet-Higgins and Coulson (CA 41, 4978b) but only poor correlation with those reported by Basur and Bhattacharya (CA 52, 864i and by Pullman (CA 41, 197b). The split carbonyl bands observed for the esters VI-XII should be ascribed to conformational isomerism rather than to Fermi resonance. Relative pKa values of I-V were predictable from electron densities of the unsubstituted quinoxaline ring even though the carboxylate anions formed during the determination must perturb the ring densities. The effect is apparently small in relation to the perturbed electron d. caused by the ring H atoms in the unsubstituted quinoxaline nucleus.

IT 6924-72-7, 6-Quinoxalinecarboxylic acid, ethyl ester
(spectrum of, reactivity and)

RN 6924-72-7 CAPLUS
CN 6-Quinoxalinecarboxylic acid, ethyl ester (6CI, 7CI, 8CI, 9CI) (CA INDEX NAME)



L13 ANSWER 166 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1965:489504 CAPLUS
DOCUMENT NUMBER: 63:89504

PATENT INFORMATION:

sulfoethyl)-6-quinoxalinecarboxamido[ethyl]phenyl]sulfamoyl]phthalocyanine

Cobalt(II) (3-). [C-]

disulfonato(5-)-N29,N30,N31,N32]-, trihydrogen (9CI) (CA INDEX NAME)

$$2 \left[\text{Dl-SO}_3^- \right]$$
 $\bullet_3 \text{ H}^+$

disulfonato(5-)-N29,N30,N31,N32]-, trihydrogen (9CI) (CA INDEX NAME)




$$2 \left[\text{Dl-SO}_3^- \right]$$
 $\bullet_3 \text{ H}^+$

{CA INDEX NAME}




$$2 \left[\text{D1-SO}_3^- \right]$$

● 4+




disulfonato(5-)-N29,N30,N31,N32]-, trihydrogen (9CI) (CA INDEX NAME)



RN 31215-23-3 CAPLUS
CN Cuprate(3-), [C-[[[4-[[2-[[[2,3-dichloro-6-quinoxaliny]carbonyl]methylamino]butyl]-5-sulfonyl]amino]sulfonyl]-2,9H,31H-phthalocyanine-C,C-disulfonato(5-)-N29,N30,N31,N32]-, trihydrogen (9CI) (CA INDEX NAME)



RN 31216-61-2 CAPLUS
CN Cobaltate(3-), [C-[[[4-[[2-[[[2,3-dichloro-6-quinoxaliny]carbonyl]methylamino]butyl]-5-sulfonyl]amino]sulfonyl]-2,9H,31H-phthalocyanine-C,C-disulfonato(5-)-N29,N30,N31,N32]-, trihydrogen (9CI) (CA INDEX NAME)



L13 ANSWER 167 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1965.455160 CAPLUS
DOCUMENT NUMBER: 63:55160
ORIGINAL REFERENCE NO.: 63:10100f-h,10101a-b
TITLE: Reactive dyes
INVENTOR(S): Rothman, Leonard A.
PATENT ASSIGNEE(S): E. I. du Pont de Nemours & Co.
SOURCE: 11 pp.
DOCUMENT TYPE: Patent
LANGUAGE: Unavailable
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 1384789		19650108	FR 1963-955250	19631127
GB 1000527			GB	
US 3232931		19660201	US 1962-240747	19621128



PRIORITY APPLN. INFO.:
OI For diagram(s), see printed CA Issue.
AB Comps. of the general formula I, where Pc is phthalocyanine, R1 is H,

alkyl, or aryl, A is a p-C6H4 or sulfonylphenylene, R2 is H or St, R3 is a heterocyclic residue, and a + b + c = 3-4, dye cotton turquoise. Thus, 23 parts CuPc was added to 160 parts ClSO3H at 5° keeping the temperature below 25°, the mixture stirred 15 min., heated during 1.5 hrs. to 135 ± 5°, stirred 3.5 hrs. at that temperature, then the green solution cooled to room temperature, poured into a mixture of ice and H2O at <5°, the precipitate filtered and washed with 1% HCl at 5°, to give CuPc(3-SO3H)x(3-SO2Cl)y(x + y = 3-4) (II). 2,4-(H2N)2C6H3SO3H (III) (22.5 parts) was added to II in 1200 parts ice-H2O, the pH adjusted to 5 with 10 N NaOH, and then to 9.9 with aqueous NH3, the mixture heated to 25-30° and stirred until the pH stayed at 8.8-9.2 without further addition of NH3 (about 15 hrs.), acidified with HCl, the precipitate filtered, washed with aqueous HCl to remove excess III, the filter cake mixed with 1500 parts H2O at 35-40°, the suspension adjusted to pH 7.2 with NaOH, 21 parts 2,3-dichloro-6-quinoxalinecarbonyl chloride (IV) added, the mixture stirred 6 hrs. at 35-40°, keeping the pH at 7-7.5 with NaOH, the turquoise solution filtered, NaCl carefully added to 10%, the precipitate filtered, washed with 20% aqueous NaCl, and vacuum-dried at 60° to give I [R = R1 = R2 = H, A = 4-sulfo-m-phenylene, R3 = 2,3-dichloro-6-quinoxaliny], a = 1.5, b = 1.3, c = 1.1). Similarly, other I were prepared (R1, A, R2, R3, a, b, and c given): H, 3-sulfo-p-phenylene, H, 3-chloro-6-quinoxazoliny, 0.9, 1.2, 1.6; H, 3-sulfo-p-phenylene, St, 2,3-dichloro-6-quinoxazoliny, 0.3, 2.3, 1.1; C6H4SO3Na-3, p-C6H4, H, 2,3-dichloro-6-quinoxazoliny, -, -, -; St, 4-sulfo-p-phenylene, H, 2,3-dichloro-6-quinoxazoliny, -, -, -; H, 4-sulfo-p-phenylene, H, 1,4-dichloro-6-phthalaziny, -, -, -. Also, a turquoise dye was obtained by using CuPc(4-SO3H)4, ClSO3H, III, and IV. Preparation of intermediates: trimellitic anhydride was treated with H2-NH2. H2O in AcOH to give 1,4-dihydroxy-6-phthalazinecarboxylic acid, which with PCl5 and POCl3 gave the dichloro acid chloride, m. 130-2°. 2,3-Dihydroxy-6-quinoxalinecarboxylic acid with COCl2 in HCONMe2 and p-C6H4Me2 gave IV, m. 111-13° similarly were prepared 2-and 3-chloro-6-quinoxalinecarbonyl chloride, m. 127-9° and 120-2°, resp.

IT 31157-46-7, Copper, [hydrogen [[4-(2,3-dichloro-N-ethyl-6-quinoxalinecarboxamido)-3-sulfonyl]sulfonyl]diulfamoylphthalocyaninesulfonato(2-)]-, disodium salt (mixture-containing)

RN 31157-46-7 CAPLUS
CN Cuprate(2-), [C-C-bis(aminosulfonyl)-C-[[[4-[[[2,3-dichloro-6-quinoxaliny]carbonyl]ethylamino]-3-sulfonyl]amino]sulfonyl]-2,9H,31H-phthalocyanine-C-sulfonato(4-)-N29,N30,N31,N32]-, disodium (9CI) (CA INDEX NAME)



D1-SO₃⁻●2 Na⁺

L13 ANSWER 168 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1965:439568 CAPLUS
 DOCUMENT NUMBER: 63:39568
 ORIGINAL REFERENCE NO.: 63:7145b-f
 TITLE: Chromium- and cobalt-containing azo dyes
 PATENT ASSIGNEE(S): J. R. Geigy A.-G.
 SOURCE: 23 pp.
 DOCUMENT TYPE: Patent
 LANGUAGE: Unavailable
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
NL 6411791		19650412	NL 1964-11791	19641009
PRIORITY APPLN. INFO.:			CH	19631011


AB 2,5-HO(O2N)C6H3N:NCPh-NNHC6H4SO2NH2-m (I) 22, from equimolar amt. diazotized 2,4-H2N(O2N)C6H3OH (II) and m-H2NSO2C6H4NH2: CHPh, and 2,5-HO(EtO2S)C6H3N:NCPh-NNHC6H4SO2NH2-m (III), from diazotized 2,4-H2N(EtO2S)C6H3OH (IV) and o-AcCH2CONH-C6H4Cl (V), in HCONH2 300 treated at 80-5° with Co(OAc)2 15 parts and salted with NaCl gave a dark powder which dyes wool yellowish olive-green. I 13.2 and III 29.7 parts gave similarly a stronger yellowish dye; I 30.8 and III 12.7 parts gave a more grayish dye. Similar Co complex dyes were obtained from I 22 with IV + m-AcCH2CONH-C6H4Cl 21.2, with IV + AcCH2CONHPh 19.5, and with II + V 18.5 parts. 2,5-HO(O2N)-C6H3N:NCPh-NNHC6H4SO2NH2CH2CH2OH-m 24.2 and 1,2-[2,4-HO(O2N)C6H3N:N]C10H6OH 15.5 in HCONH2 300 treated at 80-5° with Co(OAc)2 15 parts gave a dark powder which dyes wool navy-blue. Cr complex (1:1) (VI) 22.2 of 2,4,1-HO-(HO3S)C10H6NH2 → 2-ClOH7OH in HCONH2 333 treated with o-HO3CC6H4N:NCPh-NNHC6H4SO2NH2 17.2 and heated with Na2CO3 15 parts at 100-5° gave a dark powder; it dyes gray shades. The same dye was also obtained by treating VI 22.3 parts in EtOH 800 vols. and H2O 150 parts with VII 17.2 parts and 10N NaOH 15 vols. and heating at 60-5°. Similar dyes were prepared from VI 22.3 with 2,5-H2N(PH-MeSO2)C6H3CO2H → PhCH: NNHPh (VIII) 25, with 2,5-H2N(PH-MeSO2)C6H3CO2H → VIII 25.7, or with 4,3-HO(H2N)C6H3SO2-Ph → VIII. p-isomer (IX) of I 22 and

2,4-HO(PhSO2)C6H3N:CHC6H4OH-o 17.7 in HCONH2 300 with Co(OAc)2 15 parts at 80-5° gave a dark powder, olive-green on wool. Similar dyes were obtained from IX 22 with 2,4-HO(PhN:N)C6H3CH:NC6H3-(OH)NO2-2,5 18.1 or with 2,4-HO(PhSO2)C6H3N:CHC6H3(OH)-Cl-2,5 19.4 parts. 3,2,5-Cl(HO)C6H2N:NCPh-NNHC6H4SO2NH2-m 33.7 and IV + 3-methyl-1-phenyl-5-pyrazolone (X) 19.3 with Co(OAc)2 15 parts gave a dark powder, brown-olive on wool. Similar Co complex dyes were prepared from mixts. of 5,2-Cl(HO)C6H3N:NCPh-NNHC6H4SO2NH2-m (XI) 21.5 and 1,2-[2,5-HO(O2N)C6H3N:N]C10H6NH2 15.4 (olive-gray), and of 2,5,4-HO(EtSO2)C6H3N:NCPh-NNHC6H4SO2NH2-m p 27.3 and 5,2,4-Me(HO)C6H2N:NCPh-NNHC6H4SO2NH2 → 2-ClOH7OH 19.1 parts (olive-blue), of XI 21.5, III 10.6, and IV + X 9.65 (olive), and of I 11, XI 10.8, and III 21.2 parts (olive-green). I 22 and 2,5-HO(MeSO2)C6H3NH2 → 2,7-HOClO5NHAc 20 with Cr-(OAc)216 and Na2CO3 15 parts in HCONH2 333 vols. gave an olive-gray dye.


IT 4445-61-8, 2,7-Naphthalenedisulfonic acid, 4-amino-6-[[α-(2,3-dichloro-N-methyl-6-quinoxalinedicarboxamido)-4-sulfo-m-tolyl]azo]-5-hydroxy-3-[(p-nitrophenyl)azo]- (preparation of)

RN 4445-61-8 CAPLUS
 CN 2,7-Naphthalenedisulfonic acid, 4-amino-6-[[α-(2,3-dichloro-N-methyl-6-quinoxalinedicarboxamido)-4-sulfo-m-tolyl]azo]-5-hydroxy-3-[(p-nitrophenyl)azo]- (7CI, 8CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



L13 ANSWER 169 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1965:439567 CAPLUS
 DOCUMENT NUMBER: 63:39567
 ORIGINAL REFERENCE NO.: 63:7144g-h, 7145a-b
 TITLE: Monoazo dyes
 PATENT ASSIGNEE(S): Farbenfabriken Bayer A.-G.
 SOURCE: 21 pp.
 DOCUMENT TYPE: Patent
 LANGUAGE: Unavailable
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
NL 6410555		19650315	NL 1964-10555	19640910
PRIORITY APPLN. INFO.:			DE	19630909

AB Azo dyes contg 2,3-dichloro-6-quinoxaliny, dior trichloropyrimidyl, or

substituted s-triazinyl groups were prepared p-H2NC6H2NHMe (9.85 parts) diazotized and coupled with 32 parts 8,4,6,3-HO(HO3S)2C10H4NHC6H3SO3H and the mixture treated at 35° with 2,3-dichloroquinoxaline-6-carbonyl chloride (I) 19 and Na2CO3 7.7 in H2O 40 parts yielded II. 4,3-H2N(HO3S)C6H3CH2NHMe (12.5 parts) diazotized and coupled with 21 parts 8,3,6,1-HO(HO3S)2C10H4NHAc, and the mixture treated with 15.2 parts I gave a dye which dyed cotton and regenerated cellulose fabrics brilliant bluish red shades of good wet- and lightfastness. 3,4-H2N(HO3S)C6H3CH2NHMe (III) (12.5 parts) diazotized and coupled with 21.8 parts 5,2,1,7-HO(H2NCONH)C10H4(SO3H)2, and the mixture treated at 40-5° with I gave an orange-red dye. III (12.5 parts) diazotized and coupled with 19.9 parts 5,7,2-HO(HO3S)C10H5NH2 (IV), and the mixture coupled with 10.7 parts cyanuric chloride in 90 parts Me2CO gave a brilliant orange dye. III (12.5 parts) diazotized and coupled with 12 parts, 4,1-HO3SC10H6OH and then condensed with 13 parts 2,4-dichloropyrimidine-6-carbonyl chloride gave a brilliant scarlet dye. III (12.5 parts) diazotized and coupled with 19.9 parts IV and then condensed with 12.7 parts tetrachloropyrimidine gave a light orange dye. 5,2-H2N(HO3S)C6H3CH2NHMe (12.5 parts) diazotized and coupled with 27.3 parts p-O2NC5H4NH2 8,3,6,1-HO(HO3S)2C10H4NH2 and then condensed with 15.2 parts I gave a blue dye. III (12.5 parts) diazotized and coupled with 16.4 parts 5,7,2-HO(SO3H)C10H5NHCONH2 and then condensed successively with 10.7 parts cyanuric chloride and 10 parts m-HO3SC6H4NH2 gave a brilliant orange dye.


IT 2533-09-7, 2,7-Naphthalenedisulfonic acid, 5-acetamido-3-[[α-(2,3-dichloro-N-methyl-6-quinoxalinedicarboxamido)-3-sulfo-p-tolyl]azo]-4-hydroxy-, triisodium salt 2752-29-6, 1,7-Naphthalenedisulfonic acid, 6-[[α-(2,3-dichloro-N-methyl-6-quinoxalinedicarboxamido)-p-tolyl]azo]-5-hydroxy-4-(2-sulfoacetamido)-, triisodium salt (preparation of)

RN 2533-09-7 CAPLUS
 CN 2,7-Naphthalenedisulfonic acid, 5-acetamido-3-[[α-(2,3-dichloro-N-methyl-6-quinoxalinedicarboxamido)-3-sulfo-p-tolyl]azo]-4-hydroxy-, triisodium salt (7CI, 8CI) (CA INDEX NAME)



●3 Na

RN 2752-29-6 CAPLUS
 CN 1,7-Naphthalenedisulfonic acid, 6-[[p-((2,3-dichloro-N-methyl-6-quinoxalinedicarboxamido)methyl)phenyl]azo]-5-hydroxy-4-(2-sulfoacetamido)-, triisodium salt (8CI) (CA INDEX NAME)



●3 Na


L13 ANSWER 170 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1965:439122 CAPLUS
 DOCUMENT NUMBER: 63:39122
 ORIGINAL REFERENCE NO.: 63:7011c-f
 TITLE: Some substituted 1,2,3,4-tetrahydroquinoxalines and Hofmann degradation of a quaternary ammonium hydroxide derived from N,N'-dimethyltetrahydroquinoxaline
 Elina, A. S.; Musatova, I. S.
 S. Ordzhonikidze All-Union Chem.-Pharm. Research Inst., Moscow
 Khimiya Geterosiklicheskikh Soedinenii (1965), (2), 291-5
 CODEN: KGSSAQ; ISSN: 0132-6244
 DOCUMENT TYPE: Journal
 LANGUAGE: Russian

AB N,N,N'-Trimethyl-1,2,3,4-tetrahydroquinoxalinium iodide (I) undergoes a degradation in alkaline solution with the formation of methylated o-phenylenediamine deriva. N-Acetyl-1,2,3,4-tetrahydroquinoxaline and PhCH2Cl yielded a mixture of products: N-Acetyl-N'-benzyl-1,2,3,4-tetrahydroquinoxaline, m. 60.5-2.5°; N-benzyl-1,2,3,4-tetrahydroquinoxaline (II), b2 195-205° (HCl salt 3.170-1°); 1,2,3,4-tetrahydroquinoxaline, m. 92-8°; and N,N'-dibenzyl-1,2,3,4-tetrahydroquinoxaline, b2 210-30° (HCl salt m. 175-7°). II and MeI formed N,N'-dimethyl-N'-benzyl-1,2,3,4-tetrahydroquinoxalinium iodide m. 175-6°. I and 40% solution of NaOH refluxed 6 hrs. gave N,N,N'-trimethyl-o-phenylenediamine (III), b1 62-4° (picrate m. 112-13°; HCl salt m. 165-7°) and N,N'-dimethyl-1,2,3,4-tetrahydroquinoxaline, b1 92-4°; picrate m. 122-4°. Distillation of I gave III. MeI and III gave


2-methylaminophenyltrimethylammonium iodide, m. 207-8°. Exhaustive methylation of III yielded 2-dimethylaminophenyltrimethylammonium iodide, m. 219-20° (decomposition), the product of the reaction of N,N,N',N'-tetramethyl-o-phenylenediamine with MeI.

IT 2427-91-0, 6,7-Quinoxalinedicarboxylic acid, 5,8-dihydroxy-, diethyl ester (preparation of)

RN 2427-91-0 CAPLUS
 CN 6,7-Quinoxalinedicarboxylic acid, 5,8-dihydroxy-, diethyl ester (7CI, 8CI) (CA INDEX NAME)




L13 ANSWER 171 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1965:439121 CAPLUS
 DOCUMENT NUMBER: 63:39121
 ORIGINAL REFERENCE NO.: 63:7011a-c
 TITLE: Chelating reagents containing nitrogen heterocycles.
 IV. Syntheses of 5,8-dihydroxyquinoline derivatives
 Oguchi, Shoichi
 AUTHOR(S): Tokyo Gakugei Univ.
 CORPORATE SOURCE: Nippon Kagaku Zasshi (1965), 86(4), 435-7
 SOURCE: CODEN: NPKZAZ; ISSN: 0369-5387
 DOCUMENT TYPE: Journal
 LANGUAGE: Japanese
 AB cf. CA 63, 4295c. 2,3-Dichloro-5,8-dimethoxyquinoline (I) (1 mole) and 2.4 g. atom Na in appropriate alc. was heated 3 hrs. to give the following 2,3-dialkoxy-5,8-dimethoxyquinolines (alkyl in alkoxy, m.p., and % yield given): Et, 178°, 73; EtO(CH₂)₂, 114°, --. Heating I with 40% aqueous MeNH₂ at 120-130° for 3 hrs. gave 2,3-bis(dimethylamino)-5,8-dimethoxyquinoline, m. 116-117°. Using 1 mole 2,3-dichloro-5,8-dihydroxyquinoline and 4.4 g. atom Na in alc. gave the following 2,3-dialkoxy-5,8-dihydroxyquinolines (alkyl, m.p. and m.p. of acetate given): Et, 148°-168°; EtO(CH₂)₂, 65°, 78°. Similarly, 2,3-bis(ethylthio)-5,8-dihydroxyquinoline, m. 151-152°, and 2,3-bis(dimethylamino)-5,8-dihydroxyquinoline, m. approx. 80° (unstable), were prepared (CH₂CO₂Et) (II) (4.2 g.), 4.4 g. di-Et pyrazine-2,3-dicarboxylate, b₃ 165°, n_D20° 1.5059, and 1.0 g. powdered Na in xylene was heated 1 hr. at 110° to give 2.7 g. di-Et 5,8-dihydroxyquinoline-6,7-dicarboxylate, m. 149°; diacetate m. 205-6°. Similarly, di-Me 5,8-dihydroxyquinoline-6,7-dicarboxylate, m. 183-4°, was obtained in 30% yield. Similar treatment of di-Et 2,3-dimethylpyrazine-5,6-dicarboxylate with II failed to give quinoline derivative but gave di-Et cyclohexane-2,5-dione-1,4-dicarboxylate.
 IT 2427-91-0, 6,7-Quinoxalinedicarboxylic acid, 5,8-dihydroxy-, diethyl ester 2452-36-0, 6,7-Quinoxalinedicarboxylic acid, 5,8-dihydroxy-, diethyl ester, diacetate (ester) (preparation of)
 RN 2427-91-0 CAPLUS
 CN 6,7-Quinoxalinedicarboxylic acid, 5,8-dihydroxy-, diethyl ester (7CI, 8CI) (CA INDEX NAME)




RN 2452-36-0 CAPLUS
 CN 6,7-Quinoxalinedicarboxylic acid, 5,8-bis(acetyloxy)-, diethyl ester (9CI)


(CA INDEX NAME)




L13 ANSWER 172 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1965:424143 CAPLUS
 DOCUMENT NUMBER: 63:24143
 ORIGINAL REFERENCE NO.: 63:4295c-f
 TITLE: Chelating reagents containing nitrogen heterocycles.
 I. Syntheses of 1,4-dihydroxyphenazine derivatives
 Oguchi, Shoichi
 AUTHOR(S): Tokyo Gakugei Univ.
 CORPORATE SOURCE: Nippon Kagaku Zasshi (1965), 86(2), 246-9
 SOURCE: CODEN: NPKZAZ; ISSN: 0369-5387
 DOCUMENT TYPE: Journal
 LANGUAGE: Japanese
 AB The optimal conditions for condensation of di-Et quinoxaline-2,3-dicarboxylate (I) with (CH₂CO₂Et)₂ (II) were sought. Using NaOEt, NaNH₂, Ph₃CN, and NaH as base, 32% di-Et cyclohexane-2,5-dione-1,4-dicarboxylate, quinoxaline-2,3-dicarboxamide, tar, and 35% di-Et 1,4-dihydroxyphenazine-2,3-dicarboxylate (III) were obtained, resp. The best yield (51%) of III was obtained when 1:1:2 molar ratio of I, II, and Na or K was used. 4,5-(O₂N)₂CH₂(O₂N)2-1,2 (9.2 g.), 25 g. Sn, 130 ml. concentrated HCl, and 60 ml. EtOH was heated to give 4,5-(H₂N)₂CH₂(O₂N)2-1,2 (IV), m. approx. 98° (decomposition); di-Ac derivative m. 162°. Treatment of IV with [HO₂C(CH₂)₂CO₂H] gave 6,7-dibutoxyquinoline-2,3-dicarboxylic acid, m. 148-50° (decomposition); di-Et ester b₇ 234°, m. 95-6°. Similarly, 6,7-dimethoxyquinoline-2,3-dicarboxylic acid, m. 238-40° [di-Et ester (V) m. 146°; di-Bu ester m. 97-8°], was prepared. V, II, and Na in 1:1:2 molar ratio was treated as above to give 47% di-Et 1,4-dihydroxy-7,8-dimethoxyphenazine-2,3-dicarboxylate, m. 276-7°; diacetate, m. 218-19°. Similarly, di-Bu 1,4-dihydroxy-7,8-dimethoxyphenazine-2,3-dicarboxylate, m. 233-4° (diacetate m. 165-6°), and di-Et 1,4-dihydroxy-7,8-dibutoxyphenazine-2,3-dicarboxylate, m. 181-2° (diacetate, m. 226-7°), were prepared. Similar treatment of I with (CH₂CO₂Et)₂ gave 74% 2,3-dicyano-1,4-dihydroxyphenazine, m. >360°; monoacetate m. 274-6°. I and (AcCH₂)₂ gave 33% 2,3-diacetyl-1,4-dihydroxyphenazine, m. 241-2°. IT 1770-41-8, 2,3-Phenazinedicarboxylic acid, 1,4-dihydroxy-7,8-dimethoxy-, diethyl ester 1910-93-6, 2,3-Phenazinedicarboxylic acid, 1,4-dihydroxy-7,8-dimethoxy-, dibutyl ester, diacetate (ester) 1910-94-7, 2,3-Phenazinedicarboxylic acid, 7,8-dibutoxy-1,4-dihydroxy-, diethyl ester 1910-95-8, 2,3-Phenazinedicarboxylic acid, 7,8-dibutoxy-1,4-dihydroxy-, diethyl ester, diacetate (ester) 1983-91-1, 2,3-Phenazinedicarboxylic acid, 1,4-dihydroxy-7,8-dimethoxy-, dibutyl ester 3684-53-5, 2,3-Phenazinedicarboxylic acid, 1,4-dihydroxy-7,8-dimethoxy-, diethyl ester, diacetate (ester) (preparation of)
 RN 1770-41-8 CAPLUS
 CN 2,3-Phenazinedicarboxylic acid, 1,4-dihydroxy-7,8-dimethoxy-, diethyl ester (7CI, 8CI) (CA INDEX NAME)




RN 1910-93-6 CAPLUS
 CN 2,3-Phenazinedicarboxylic acid, 1,4-dihydroxy-7,8-dimethoxy-, dibutyl ester, diacetate (ester) (8CI) (CA INDEX NAME)




RN 1910-94-7 CAPLUS
 CN 2,3-Phenazinedicarboxylic acid, 7,8-dibutoxy-1,4-dihydroxy-, diethyl ester (7CI, 8CI) (CA INDEX NAME)




RN 1910-95-8 CAPLUS
 CN 2,3-Phenazinedicarboxylic acid, 1,4-bis(acetyloxy)-7,8-dibutoxy-, diethyl ester (9CI) (CA INDEX NAME)



RN 1983-91-1 CAPLUS
 CN 2,3-Phenazinedicarboxylic acid, 1,4-dihydroxy-7,8-dimethoxy-, dibutyl ester (7CI, 8CI) (CA INDEX NAME)



RN 3684-53-5 CAPLUS
 CN 1,2,3,4-Phenazinetetrol, 7,8-dimethoxy-, tetraacetate (ester) (8CI) (CA INDEX NAME)




L13 ANSWER 173 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
 ACCESSION NUMBER: 1963:448886 CAPLUS
 DOCUMENT NUMBER: 59:48886
 ORIGINAL REFERENCE NO.: 59:8910e-h
 TITLE: Water-soluble anthraquinone reactive dyes
 Singer, Josef; Schwechten, Heinz W.
 PATENT ASSIGNER(S): Farbenfabriken Bayer A.-G.
 SOURCE: 18 pp.
 DOCUMENT TYPE: Patent
 LANGUAGE: Unavailable
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
BE 622569		19630115	BE	
GB 1014055			GB	
US 3251844		19660517	US 1962-222937	19620911
			DE	19610933

PRIORITY CLAIM INFO:
 GI For diagram(s), see printed CA issue.
 AB Dyes of the formula I are suitable for dyeing and printing silk, wool, polyamides, polyurethanes, and especially natural and regenerated cellulose fast blue shades. Thus, 1-amino-4-[4-(methylaminomethyl)anilino]anthraquinone-2-sulfonic acid 43.7 and NaOH 4 were dissolved in H₂O 1000, 2,3-dichloro-6-quinoxalinecarboxyl chloride 27 parts added at 40° by stirring while maintaining pH 6-8 by adding NaOH solution, the dye salted, filtered, washed with NaCl solution, and dried at 40-50° to give a blue powder, blue in H₂O, which dyed cotton light- and wetfast blue shades in the presence of Na₂CO₃. Similarly, the following I were prepared (V, W, X, Y, and Z are given): H, 3-NaO₃S, 4-CH₂, Me, 2,3-trichloro-6-quinoxalinecarboxyl (II); H, 3-NaO₃S, 4-CH₂CH₂, H, II; H, 3-NaO₃S, 4-(CH₂)₃, Me, II; H, NaO₃S, 3-CH₂CH₂CH₂(Me), Me, II; H, NaO₃S, 3-CH₂CH₂CH₂(CH₂CH₂CH₂Me), Me, II; H, NaO₃S, H, 4-CH₂, Me, II; H, 3-NaO₃S, 4-O(CH₂)₄, Me, II; H, NaO₃S, 4-S(CH₂)₄, Me, II; NaO₃S, H, 3-SO₂(CH₂)₄, Me, II; H, NaO₃S, 3-CH₂, Me, 4,6-dichloro-s-triazin-2-yl; H, NaO₃S, 4-CH₂, Me, 4-chloro-6-methoxy-s-triazin-2-yl.
 IT 10009-44-7, 2-Anthracenesulfonic acid, 1-amino-4-[4-(2,3-dichloro-6-quinoxalinecarboxamido)ethyl]-3-sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt 104242-65-1, 2-Anthracenesulfonic acid,

1-amino-4-[α -(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)-p-toluidino]-9,10-dihydro-9,10-dioxo-, sodium salt 104601-65-2, 2,6-Anthracenedisulfonic acid, 1-amino-4-[α -(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)-p-toluidino]-9,10-dihydro-9,10-dioxo-, disodium salt 104601-66-3, 2-Anthracenesulfonic acid, 1-amino-4-[α -(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)-3-sulfo-p-toluidino]-9,10-dihydro-9,10-dioxo-, disodium salt 105232-45-9, 2-Anthracenesulfonic acid, 1-amino-4-[3-[3-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)-5-methylhexyl]sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt 106303-94-0, 2-Anthracenesulfonic acid, 1-amino-4-[4-[3-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)propyl]-3-sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt 106337-79-5, 2-Anthracenesulfonic acid, 1-amino-4-[4-[4-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)butyl]thio]-3-sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt 106381-99-0, 2-Anthracenesulfonic acid, 1-amino-4-[4-[4-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)butoxy]-3-sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt 106381-99-1, 2,6-Anthracenedisulfonic acid, 1-amino-4-[m-[4-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)butyl]sulfonyl]anilino]-9,10-dihydro-9,10-dioxo-, disodium salt 107062-63-5, 2-Anthracenesulfonic acid, 1-amino-4-[3-[3-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)butyl]sulfonyl]anilino]-9,10-dihydro-9,10-dioxo-, disodium salt (preparation of)

RN 104099-44-7 CAPLUS
CN 2-Anthracenesulfonic acid, 1-amino-4-[4-[2-(2,3-dichloro-6-quinoxalinecarboxamido)ethyl]-3-sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt (7CI) (CA INDEX NAME)




PAGE 1-A

NH₂


● Na

RN 104242-65-1 CAPLUS
CN 2-Anthracenesulfonic acid, 1-amino-4-[α -(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)-p-toluidino]-9,10-dihydro-9,10-dioxo-, sodium salt (7CI) (CA INDEX NAME)




● Na

RN 104601-65-2 CAPLUS
CN 2,6-Anthracenedisulfonic acid, 1-amino-4-[α -(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)-p-toluidino]-9,10-dihydro-9,10-dioxo-, disodium salt (7CI) (CA INDEX NAME)




● 2 Na

RN 104601-66-3 CAPLUS
CN 2-Anthracenesulfonic acid, 1-amino-4-[4-[4-(2,3-dichloro-6-quinoxaliny)carbonyl]methylamino]methyl]-3-sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt (9CI) (CA INDEX NAME)




● 2 Na

RN 105232-45-9 CAPLUS
CN 2-Anthracenesulfonic acid, 1-amino-4-[3-[3-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)-5-methylhexyl]sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt (7CI) (CA INDEX NAME)

D1-SO₃H


● 2 Na

RN 106303-94-0 CAPLUS
CN 2-Anthracenesulfonic acid, 1-amino-4-[4-[3-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)propyl]-3-sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt (7CI) (CA INDEX NAME)




● 2 Na

RN 106337-79-5 CAPLUS
CN 2-Anthracenesulfonic acid, 1-amino-4-[4-[4-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)butyl]thio]-3-sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt (7CI) (CA INDEX NAME)




● 2 Na

RN 106381-98-0 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[4-[(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)butoxy]-3-sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt (7CI) (CA INDEX NAME)




● 2 Na

RN 106381-99-1 CAPLUS
 CN 2,6-Anthracenedisulfonic acid, 1-amino-4-[m-[[4-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)butyl]sulfonyl]anilino]-9,10-dihydro-9,10-dioxo-, disodium salt (7CI) (CA INDEX NAME)



● 2 Na

RN 107062-63-5 CAPLUS
 CN 2-Anthracenesulfonic acid, 1-amino-4-[3-[(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)butyl]sulfoanilino]-9,10-dihydro-9,10-dioxo-, disodium salt (7CI) (CA INDEX NAME)

D1-SO₃H

● 2 Na

L13 ANSWER 174 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1963:429002 CAPLUS
 DOCUMENT NUMBER: 59:29002
 ORIGINAL REFERENCE NO.: 59:5299d-f,5300a-c
 TITLE: Quinoxaline dyes
 INVENTOR(S): Siegel, Edgar; Sasse, Klaus
 PATENT ASSIGNER(S): Farbenfabriken Bayer A.-G.
 SOURCE: 83 pp.
 DOCUMENT TYPE: Patent
 LANGUAGE: Unavailable
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
BE 613586		19620807	BE	19610207

PRIORITY APPLN. INFO.:

AB Condensation compds. of 3- or 2-chloro- and 2,3-dichloroquinoxaline derivs. with azo, anthraquinone, or phthalocyanine dyes are H₂O-soluble and suitable for dyeing or printing cellulosic and other materials. Thus, 2,4,8-H₂N(NaO₃S)2C10H₅ (I) 34.7 was diazotized and coupled with 3-H₂NC₆H₄Me (II) 10.7 parts. The product in aqueous NaOH was mixed and stirred with 26.5 parts 2,3-dichloro-6-quinoxalinecarbonyl chloride (III) (m. 116°, b.p. 144°) in 100 parts C₆H₆, heated to 35-40°, neutralized with Na₂CO₃, and 80 parts NaCl added. The dye was filtered and dried at 40-50°. Printed on cellulose, it gave a reddish yellow color, fast to washing and light. Similarly, dyes were prepared (reactants and color of dye on cellulosic fabric given): I → II, 2,3-dichloro-6-isocyanatoquinoxaline, reddish yellow; I → 3-MeHNC₆H₄Me, 2,3-dichloro-6-quinoxalinesulfonyl chloride (IV), reddish yellow; 2-H₂NC₆H₄SO₃H (V) → [1,8,3,6-H₂N(HO)(NaO₃S)2C10H₄ (VI), III], bluish red [2,4-(H₂N)2C₆H₃SO₃Na (VII), III] → 1,8,3,6-(BzNH)(HO)(NaO₃S)2C10H₄, bluish red; Cu complex of [3,4-H₂N(HO)C₆H₃SO₃H → 2,5,7-H₂N(HO)(HO₃S)C10H₅, III, ruby; Cu phthalocyaninetetra-sulfonyl chloride, VII, III, blue; 1-amino-4-bromo-2-anthraquinonesulfonic acid, [4,2-H₂N(HO₃S)C₆H₃]2, III, blue; Cr complex of [3,4,5-Cl(HO)(H₂N)C₆H₂SO₃H → 1- [3- [(3-aminophenyl)sulfonyl]sulfonyl]phenyl]-3-methyl-5-pyrazolone], III, yellow brown; V → [VI, 2- or 3-chloro-6-quinoxalinecarbonyl chloride], red; V → [VI, III, H₂NC₆H₄SO₃Na], red; 6-amino-2,3-dichloroquinoxaline (VIII) → 1-[2,5-dichloro-4-sulfonyl]-3-methyl-5-pyrazolone, reddish yellow; 2,1,7-H₂N(HO₃S)2C10H₅ [2,5,7-H₂N(HO)(NaO₃S)C10H₅, III], reddish orange; 2,5-H₂N(AcNH)C₆H₃SO₃H → 2,5,7-H₂N(HO₃S)2C10H₅, III, yellowish orange; 2,3,6,8-H₂N(HO₃S)2C10H₄ → 3-AcNH₂C₆H₄NH₂, III, reddish yellow; 2,4,6-H₂N(HO₃S)2C₆H₂OH → 2,8,6-H₂N(HO)(HO₃S)C10H₅, III, ruby; [VII, III] → 1,8,3,6-(AcNH)(HO)(HO₃S)2C10H₄, bluish red; 2,5-H₂N(O₂N)C₆H₃SO₃H → 1-[2-chloro-5-sulfonyl]-3-methyl-5-pyrazolone, III, yellow; 2,5-(4-H₂NC₆H₄NH)(2,4-O₂N(NaO₃S)C₆H₃SO₃Na, III, brown violet; PhNH₂ → [VI, III, 3-H₂NC₆H₄SO₃Na], bluish red; V → [VI, 2,3-dichloro-5-quinoxalinecarbonyl chloride], bluish red; VIII → PhN(CH₂CH₂OH)₂, orange; VIII → PhOH, reddish yellow; di-Me 1-amino-4-(2'-methyl-3'-aminophenyl)anthraquinone-2,5'-disulfonate, III, blue; equimolar mixture of the N-(3-amino-4-sulfonyl) mono- and diamides of Cu 3,3',3''-phthalocyaninetrisulfonic acid, III, turquoise; V → [VI, N-methyl-N-(2,3-dichloro-6-quinoxalinecarbonyl)glycyl chloride], bluish red; 1,4-HO(HO₃S)C10H₆ → [3,4,6-Me(H₂N)(HO₃S)C₆H₃]2 → PhOH, IV, scarlet.


IT 14573-57-0, 2,7-Naphthalenedisulfonic acid, 5-[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]-4-hydroxy-3-[(o-sulfonyl)azo]- (preparation of)

RN 14573-57-0 CAPLUS
 CN 2,7-Naphthalenedisulfonic acid, 5-[2-(2,3-dichloro-N-methyl-6-quinoxalinecarboxamido)acetamido]-4-hydroxy-3-[(o-sulfonyl)azo]- (7CI, ACT) (CA INDEX NAME)


113 ANSWER 175 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STD
ACCESSION NUMBER: 1961:17956 CAPLUS
DOCUMENT NUMBER: 55:17956
ORIGINAL REFERENCE NO.: 55:3597g-1,3598a-b
TITLE:
Synthesis of diethyl 1,4-dihydroxy-5,8-
dialkoxypheazine-2,3-dicarboxylates
Kawel, Sinichi; Otsuka, T.; Fujiki, Shun; Shibata,
Kiyoiko; Otaki, Atsuko; Sakakibara, Yoshiaki; Oguchi,
Shoshichi
CORPORATE SOURCE:
Tokyo Kyoiku Univ.
SOURCE:
Nippon Kagaku Zasshi (1959), 80, 788-91
CODEN: NPKZAZ; ISSN: 0369-5387
JOURNAL
DOCUMENT TYPE:
Unavailable
LANGUAGES:
AB Hydroquinone (I) (230 g.), 569 g. PrBr, 2.1 l. dry acetone, and 750 g.
K2CO3 refluxed 50 hrs., the solvent removed, 4. H2O added, and the mixture
extracted with ether gave 271 g. hydroquinone dipropyl ether (II), m.
50.5° (on evaporation of ether). Nitration (with HNO3 in AcOH) of
27 g. hydroquinone dipropyl ether (II) yielded 28.6 g. crude nitro compound
m. 176-177°. Mechanical separation of the crystals gave the dinitro
compound (IV) of III, m. 141.5°. Similarly, II gave the dinitro
compound (V) of II, m. 69.0-9.5°. IV shaken 50 min. with H in MeOH
over Raney Ni (in an autoclave, under 43 kg./sq. cm.) at 130-140°C
gave 20 g. 2,3-dimethoxyhydroquinone diethyl ether (VII), m.
79.7-75°. V was converted to the tetracetate (VIII), m.
139.5-46°, by refluxing it with Ac2O. VII refluxed 20 min. with
KOH in 95% EtOH gave the diacetate (VIII) of VI, m. 178-9°. V
(reduced with SnCl2 or Sn-HCl) gave 388 g. 2,3-diaminohydroquinone dipropyl
ether (IX), m. 27-8°, bs 176-8°. VI was condensed with
dihydroxytartaric acid-Na to give 931 g. 5,8-diethoxyquinoxaline-2,3-
dicarboxylic acid (X), m. 158-6° (decomposition). It was converted to
the diethyl ester (XI), m. 91-1.5°, according to the procedure of
Adachi (CA 51, 17936b). IX condensed with dihydroxytartaric acid gave 924
g. 5,8-dipropoxyquinoxaline-2,3-dicarboxylic acid (XII), m. 157°
(decomposition), which was converted to the diethyl ester (XIII), m.
68.5-69°, by the usual method. Diethyl quinoxaline-2,3-
dicarboxylate (6.2 g.) + diethyl succinate, 15 cc. xylene, and 1.5
g. finely powdered Na heated at 150-160° (5 hrs.) yielded diethyl
1,4-dihydroxypheazine-2,3-dicarboxylate, m. 165-7°. Similar runs
with diethyl 1,4-dihydroxy-5,8-dimethoxypheazine-2,3-dicarboxylate, X,
and XI gave diethyl 1,4-dihydroxy-5,8-dimethoxypheazine-2,3-
dicarboxylate, m. 181-2°, ethyl 1,4-dihydroxy-5,8-dimethoxypheazine-
2,3-dicarboxylate, m. 157°, and diethyl 1,4-dihydroxy-5,8-
dipropoxypheazine-2,3-dicarboxylate, m. 122°, resp.

IT 110081-11-3, 2,3-Phenazinedicarboxylic acid, 1,4-dihydroxy-6,9-
dimethoxy-, diethyl ester 113752-03-7, 2,3-Phenazinedicarboxylic
acid, 6,9-diethoxy-1,4-dihydroxy-, diethyl ester 114399-10-3,
2,3-Phenazinedicarboxylic acid, 1,4-dihydroxy-6,9-dipropoxy-, diethyl
ester
(preparation of)


RN 110081-11-3 CAPLUS
CN 2,3-Phenazinedicarboxylic acid, 1,4-dihydroxy-6,9-dimethoxy-, diethyl ester (5CI) (CA INDRY NAME)



RN 113752-03-7 CAPLUS
CN 2,3-Phenazinedicarboxylic acid, 6,9-diethoxy-1,4-dihydroxy-, diethyl ester
(6CI) (CA INDEX NAME)




RN 114399-30-3 CAPLUS
CN 2,3-Phenazinedicarboxylic acid, 1,4-dihydroxy-6,9-dipropoxy-, diethyl ester (6CI) (CA INDEX NAME)



113 ANSWER 176 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1957:1859 CAPLUS
DOCUMENT NUMBER: 51:1859
ORIGINAL REFERENCE NO.: 51:433f-1,434a-c
TITLE: Quinoxaline N-Oxides. VI. N-Oxides of
2,3-polymerchrysenoquinaxalines
AUTHOR(S): Landquist, Justus K.
CORPORATE SOURCE: Imperial Chem. Ltd., Manchester, UK
SOURCE: Journal of the Chemical Society (1956) 2551-3
CODEN: JCSOAS; ISSN: 0368-1769
DOCUMENT TYPE: Journal
LANGUAGES: Unavailable
GI For diagram(s), see printed CA issue.
AB 4,2-Cl(2M)C6H9N3AC (130 g.) added gradually to 110 g. "pin dust" iron,
350 cc. H2O, and 10 cc. glacial AcOH at 70 ± 2°, the temperature
raised to 80°, 30 g. CaCO3 added, the mixture kept 10 min. at
80°, filtered, and the filter cake extracted with EtOH gave
4,2-Cl(2M)C6H9N3AC (XXXIII), needles, m. 164° (from EtOH);
4,2-H2N(MeO)C6H9N3AC (XXXIV) was similarly prepared. XXXIII (120 g.) in 120

cc. each of gaseous AcOH and EtOH treated in 5 hrs. with 70 g. PhNO gave 70 g. 5,2-Cl (AcNH)C6H3N.NPH (XXIV), yellow needles, m. 180° (from alc.); similarly was prepared 244 2,5-AcNH(MeO)C6H3N.NPH (XXV), orange blades, m. 174-5° (from EtOAc). XXXV (68 g.) and 700 cc. 5% alc. KOH refluxed 9 hrs. and the mixture poured into 2 l. H₂O gave 70g 2-HZN analog, red needles, m. 131° (from petr. ether); similarly was prepared 69g 2,5-HZN(MeO)C6H3N.NPH, red laminae, m. 38.0-9.5° (from petr. ether). 4,2-(HZN)2C6H3NCl (28 g.), 23 g. 1,2-cyclohexanedione, and 240 cc. 10% aqueous AcOH heated 1 hr. at 98-100°, the mixture cooled, made alkaline with aqueous NaOH, and the precipitated solid filtered off, washed with H₂O, dried, and extracted with petr. ether left an insol. compound (XXVII), C18H18N4Cl2, pale brown prisms, m. 274-6° (from EtOH) [presumably either 1,2-(2-amino-x-chlorophenyl)phenyl)cyclohexane (XXVII) or R:C:C(R):R' (CH2)4 (R = x-chloro-o-phenylene, R' = amino-phenylene)]; the petr. ether exts. 70-70° and 170-170° of 2,3,4,5-tetrahydrophenazine (XXIII), m. 94°. 2-HN(C6H4)N2PH (10 g.), 100 cc. cyclohexanone (XL), and 0.5 cc. concentrated HCl refluxed 2 hrs. and distilled until the temperature of the mixture rose to 175°, the residue dissolved in Et2O, extracted with N HCl, and the HCl exts. made alkaline gave 6.9 g. 1,2,3,4-tetrahydrophenazine, b₇ 125-56°, m. 92-3° (from petr. ether); similarly, 4-[5,2-Me(HZN)C6H3N2]C6H4Me and XL gave 1,2,3,4-tetrahydro-7-methylphenazine (XLII), b₅ 163-4°, m. 80-2° (from petr. ether). Alkaline exts. and were 2,3-cyclopentenoquinoxaline (XLIII), b₄ 130-50°, m. 99-100°, 6-methyl-2,3-cyclopentenoquinoxaline (XLIII), b₄ 135-45°, m. 103-4°, 1,2,3,4-tetrahydro-7-methoxycyclohexenylphenazine (XLIV), m. 115-15°, and 2,3-cycloheptenoquinoxaline (XLV), m. 81-82° (4.25 g.) in 75 cc. Et₂O. 170 and 170-170° of 6-HO2C2C6H4O3CH in Et2O kept in the dark 3 days, the N-oxide phthalate filtered off, washed with Et2O, dissolved in H2O, the H2O solution adjusted to pH 7 with Na2CO3, extracted with CHCl3, and the CHCl3 exts. dried and concentrated gave the 1,4-dioxide, green-yellow leaflets, m. 179° (from petr. ether) (from C6H6); oxidation of XLIII gave only tars and XLII and XLIII were unstable to XVIII. XVIII by the above procedures gave 1,4-dioxides of the following comds.: XLI, yellow microcrystals, m. 186-8° (from EtOH); XLIV, m. 204-6° (from EtOH); XLV, brown needles, m. 172-3° (from C6H6); and bornylene[2',3',2,3]quinoxaline, green prisms, m. 137-8° (from petr. ether).

IT 099319-89-1, 6-Quinoxalinecarboxamide, 2,3-dimethyl-N-(3-piperidinopropyl)-, 1,4-dioxide
(preparation of)
1099319-69-1 CAPULUS
6-Quinoxalinecarboxamide, 2,3-dimethyl-N-(3-piperidinopropyl)-, 1,4-dioxide (GCII) [CA INDEX NAME]




L13 ANSWER 177 OF 181 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1957:1858 CAPLUS
DOCUMENT NUMBER: 51:1858
ORIGINAL REFERENCE NO.: 51:432e-1,433a-f
TITLE: Quinoxaline N-oxides. V. Further derivatives


AUTHOR(S): Silik, J. A.
CORPORATE SOURCE: Imperial Chemicals. Ltd., Manchester, UK
SOURCE: Journal of the Chemical Society (1956) 2058-63
CODEN: JCSOAG; ISSN: 0368-1769
DOCUMENT TYPE: Chemical
LANGUAGE: Unavailable
ABSTRACT: AD 4,3-H2N(O2N)C6H3OH [from 4,3-AcNH(O2N)C6H3OAc and 6M HCl] (15 g.) in 150 cc. MeOH hydrogenated over Raney Ni, filtered from the catalyst, the filtrate concentrated to 75 cc. in vacuo, mixed with 10 g. anhydrous AcONa and
28.5 g. glyoxal (XVIII) bisulfite in 140 cc. warm H2O, and the mixture heated 2.6 hrs. at 60° gave 7 g. 6-hydroxyquinoxaline (XIX), m. 252-4° (from H2O); Ac2 in place of XVIII gave the 2,3-Me2 derivative (X) of XIX, m. 247-9° (from H2O). Na (0.46 g.) in 30 cc. EtOH reacted with 2.92 g. XIX, then with 3.0 g. CH3CO2Et, the mixture heated 3 hrs., cooled, concentrated, poured into H2O, extracted with C6H6, and the C6H6 exts. concentrated gave 0.9 g. 6-EtO2CCH2 derivative of XIX, m. 99-100° (from H2O).
5-Ethoxy-2,3-dimethylquinoxaline (5 g.) in 125 cc. C6H6 and 5 g. crushed AlCl3 refluxed 16 hr., the mixture cooled, concentrated with ice H2O, the C6H6 evaporated by azeotropic distillation, the emulsion could not be broken, the solid (XXI) which separated filtered off, the filtrate adjusted to pH 4, extracted with C6H6, the C6H6 exts. extracted with hot dilute aqueous NaOH, the NaOH exts. neutralized, the solid (XXII) filtered off, and XXI and XXII recrystd. from H2O gave 0.89 g. 5-HO analog (XXIII), needles, m. 146-7°.
4,3-H2N(O2N)C6H3CO2H, m. 140-2° [obtained in 70-85% yield from 4,3-AcNH(O2N)C6H3CO2H with (a) EtOH and 33 (volume/volume) H2SO4 or (b) with EtOH-HCl, hydrogenated, gave 60% of 4,3-(H2N)C6H3CO2H (XXIV), needles, m. 112-14° (from dilute alic.). XXIV (4.5 g.) and (CHO)2 (from 6.1 g. sulfate (XV), 40 cc. H2O, and BaCO3) stirred vigorously 1 hr. at 60°, an equal portion of (CHO)2 added, and the stirring continued 1 hr. gave 2.3 g. Et 6-quinoxalinecarboxylate (XXVI), m. 48-70° (from C6H6-analog of Al2O3); XVI (1.5 g.) in 10 cc. EtOH at room temperature or at 50° gave an unidentified solid, m. about 340°. XXIV (42 g.), 22 g. Ac2, and 500 cc. 33% EtOH refluxed 30 min. gave 48 g. 2,3-Me2 derivative (XVII) of XXVI, feathery needles, m. 102-4°. XXVII (5 g.) and 15 g. Et2N(CH2)2OH refluxed 16 hr. and distilled gave 2.5 g. 6-CH2CH2 derivative (XVIII) of XVII, m. 44-5° (from EtOH).
6-CH2CH2CH2 ester (from 6-CH2CH2 analog of Al2O3) chromatography on Al2O3 on Pd-C, the mixture filtered, the filtrate treated with (CHO)2 (from 6 g. XVI), and the mixture heated 1 hr. at 60° gave 1.47 g. 6-acetylquinoxaline (XXVIII), m. 106-8° (from cyclohexane, Al2O3); XXVIII was prepared from 2,3-Me2 derivative of XVII, m. 116-18° (from aqueous EtOH). LiAlH4 (0.28 g.) in 50 cc. dry Et2O treated with 5 g. XXVII in 100 cc. dry Et2O in 10 min., the mixture stirred 10 min., 2 cc. EtOAc added, then 50 cc. H2O, the mixture filtered, the Et2O layer separated, the aqueous layer extracted with Et2O, and the combined Et2O exts. and solution dried and concentrated gave 0.7 g. 6-HOCH2 analog of XXVII, m. 113-14°.
6-Methoxyquinoxaline 1,4-dioxide (1 g.) and 2 g. AlCl3 in 25 cc. C6H6 as above gave the 6-HO analog (XXIX), yellow needles, m. 247-50° (from EtOH). 6-HOCH2 analog of XXIX (2 g.) and 2 g. 6-methylquinoxaline and 75 cc. EtOH gave 1.5 g. 6-MeOCH2 analog of XXIX, m. 247-50° (from EtOH).
VIIIA kept 17 hrs. at 60° gave 3 g. 1,4-dioxide (XXIXA), m. 249-50°. 5-Methoxy-2,3-dimethylquinoxaline 1,4-dioxide (2.33 g.), 7 g. AlCl3, and 50 cc. C6H6 stirred 17 hrs., the C6H6 decanted, the tar stirred with ice H2O and 10 cc. concentrated HCl, the solid ground with 2N

the mixture filtered, and the filtrate acidified gave 5-hydroxy-2,3-dimethylquinoxaline 1-oxide, cream needles, m. 143-4.5° (from C6H6-cyclohexane); the reaction repeated in PhNO₂ 16 hrs. at 60-5°, the mixture cooled, treated with ice H₂O and 10 cc. 10N NaOH, and the sparingly soluble Na salt filtered off and decomposed with dilute AcOH gave


5-hydroxy-2,3-dimethylquinoxaline 1,4-dioxide, m. 171-3° (from C6H6). XXIXA (0.2 g.) in 0.5 cc. HNO₃ (d. 1.4) and 0.5 cc. concentrated H₂SO₄, kept 1.5 hrs. at 0°, and the mixture poured on ice gave the 7(7)-nitro derivative, m. 258° (decomposition) (from 50% AcOH). With 1:1 HNO₃-H₂O, XXIXA gave the HNO₃ salt, m. 97° (decomposition). XXIXA (1.42 g.), 50 cc. saturated NaHCO₃, 5.25 g. iodine, and 70 cc. 10% KI kept 10 days at room temperature, filtered, and the filtrate saturated with SO₂ gave 1.3 g. 7(7)-iodo derivative (XXX), m. 148-50°; a similar procedure gave the 7(7)-Br derivative, golden needles, m. 180° (decomposition); XXX lost iodine when recrystn. was attempted. XXIXA (2.06 g.) in 100 cc. saturated NaHCO₃ and 3.7 g. Br in 30 cc. 15% KBr kept 0.25 hr. gave 1.26 g. di-Br derivative, dark red, m. 138° (explodes). XXVII (9 g.), 70 cc. 1.7M VIIIA containing 0.3% w/v H₂SO₄, and 0.1% Na₂P₂O₇ (XXXI) kept 7 hrs. at room temperature and 9 hrs. at 55°, concentrated in vacuo, and the residue treated with saturated NaHCO₃ gave 4.5 g. 1,4-dioxide (XXXII), m. 134-5° (from C₆H₆); HCO₃H and H₂O₂ in Me₂CO were unsatisfactory for this oxidation while the addition of XXXI gave more consistent results than only VIIIA and H₂SO₄. XXXII and 10N NaOH kept 0.5 hr. at room temperature and acidified with HCl gave the 6-HO₂C analog, m. 243° (decomposition) (from EtOCH₂CH₂OH). XXXII (2 g.) and 20 cc. MeOH-NH₃ kept 4 days at room temperature gave 0.95 g. 6-H₂NCO analog, m. 266° (decomposition) (from H₂O) (some hydrolysis occurs during recrystn.); the procedures described above gave the 6-Me₂NCO compound-H₂O, yellow needles, m. 215° (decomposition) (from MeOH-EtOAc); the 6-(3-piperidinopropylcarbonyl) analog, m. 172-3° (from C₆H₆), and the 6-HONHCO analog-H₂O, m. 230-2° (from aqueous AcOH). 6-Acetyl-2,3-dimethylquinoxaline (5 g.) and 36 cc. 2.1M VIIIA kept 10 hrs. at room temperature and 8 hrs. at 50° gave 2 g. 1,4-dioxide, m. 160-2° (from H₂O); oxime, m. 244-6° (from 50% AcOH). 6924-72-7, 6-Quinoxalinecarboxylic acid, ethyl ester 107419-21-6, Ethanol, 2-diethylamino-, 2,3-dimethyl-6-quinoxalinecarboxylate 109939-89-1, 6-Quinoxalinecarboxamide, 2,3-dimethyl-N-(3-piperidinopropyl)-, 1,4-dioxide (preparation of)



RN 107419-21-6 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2,3-dimethyl-, 2-diethylaminoethyl ester (6CI) (CA INDEX NAME)



RN 109939-89-1 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dimethyl-N-(3-piperidinopropyl)-, 1,4-dioxide (6CI) (CA INDEX NAME)




L13 ANSWER 178 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1956:40424 CAPLUS
DOCUMENT NUMBER: 50:40424
ORIGINAL REFERENCE NO.: 50:7812h-1,7813a-d
TITLE: Syntheses in the series of phenazine derivatives. I. 2,2'-Biphenazine and its derivatives
AUTHOR(S): Rozum, Yu. S.
SOURCE: Ukraine'skii Khimichnii Zhurnal (1955), 21, 491-5
CODEN: UKHZAS; ISSN: 0372-4190
DOCUMENT TYPE: Journal
LANGUAGE: Russian
AB (p-H₂NCO₂H)2 (I) (18.4 g.), 37 g. PhNO₂ (II), and 72 g. KOH (in 3-5 mm. granules) heated on the oil bath 4 hrs. at 90-110°, then 3 hrs. at 150°, the volatile fraction steam distilled, the residue filtered, boiled in 5-10% HCl, dried, and extracted with (CH₂Cl)₂, and the extract evaporated

gave, after chromatographing twice in C₆H₆ on Al₂O₃, 1.2 g. 2,2'-biphenazine (III), orange plates, m. 229° (from C₆H₆), insol. in Et₂O, H₂O or petr. ether, giving brightly colored, readily hydrolyzed salts in concentrated mineral acids. Analogously was obtained: from I and o-O₂NCO₂H₄OMe, 5.5% 9,9'-(MeO)₂ derivative of III, red plates or prisms, m. 181° (from C₆H₆), dark red in acids, the color fading eventually; from 1,4-HO-MeO(H₂N)CO₂H (IV) and II, 13.6% 4,4'-(MeO)₂ derivative, dark red needles, m. 174° (from C₆H₆), red in acids; from o-ClC₆H₄NO₂ (V) and I and from V and IV the 9,9'-Cl₂ (5.7%), red plates, m. 175° (from C₆H₆), and the 9,9',4,4'-Cl₂(MeO)₂ derivative (2.1%), red needles, m. 116° (from MeOH), resp. Both gave red acid solns. Similarly, I (36.8 g.) heated to 160° with 56 g. p-O₂NCO₂H₄OMe (VI) and 14 g. KOH, the product washed with ligroine and MeOH, the filter cake suspended in 3 l. H₂O, heated to 90-5°, 86 g. KOH added under agitation, the precipitated MnO₂ removed by filtration, the filtrate condensed to 300-400 ml., treated with C and glacial AcOH, and the dense yellow precipitate filtered, gave, after washing with EtOH and Et₂O and drying, 9.5% 2,2'-biphenazine-7,7'-dicarboxylic acid, yellow needles, m. 320° (from glacial AcOH), soluble in bases and concentrated acids (yellow), insol. in common organic solvents; diamide (75%), m. 360-3° (decomposition) (from HCOONH₂), insol. in common organic solvents, yellow-green in concentrated H₂SO₄; Me ester (67%), orange plates, m. 210-12° (from BuOH and active Cl), insol. in EtOH and Et₂O, yellow in concentrated H₂SO₄; Et ester (30%), red-orange needles, m. 130° (from EtOH), yellow in concentrated H₂SO₄; iso-amyl ester (34%), pink plates, m. 116° (from iso-AmOH), yellow in acid. Most of these retained 1-4 moles of crystallization solvent. Other derivs. of III prepared were: 7,7',4,4'-Me₂(MeO)₂ (9.4%) (from IV, VI, and KOH), red plates, m. 238° (from MeCHCl₂), blue in concentrated H₂SO₄; and 4,4',9,9'-(MeO)₄ (2.1%) (from IV, o-O₂NCO₂H₄OMe, and KOH), red-purple needles, m. 264° (from C₆H₆), red in H₂SO₄. Absolute maximum are given for all of these in toluene (320-390 mμ) and H₂SO₄ (410-610). IT 855639-47-3, Isopentyl alcohol, [2,2'-biphenazine]-7,7'-dicarboxylate 858239-92-6, Ethyl alcohol, compound with di-Et [2,2'-biphenazine]-7,7'-dicarboxylate

(preparation of)
RN 855639-47-3 CAPLUS
CN Isopentyl alcohol, [2,2'-biphenazine]-7,7'-dicarboxylate (5CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B

-CH₂-CH₂-CHMe₂

RN 858239-92-6 CAPLUS
CN Ethyl alcohol, compd. with di-Et [2,2'-biphenazine]-7,7'-dicarboxylate (5CI) (CA INDEX NAME)

CM 1

CRN 858239-91-5
CMP C30 H22 N4 O4



CM 2

CRN 64-17-5
CMP C2 H6 O

H₃C-CH₂-OH


L13 ANSWER 179 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1955:8306 CAPLUS
DOCUMENT NUMBER: 49:8306
ORIGINAL REFERENCE NO.: 49:1734e-i,1735a-b
TITLE: Neuphenazine derivatives and their tuberculostatic action
AUTHOR(S): Birkhofer, Leonhard; Widmann, Arno
CORPORATE SOURCE: Max-Planck-Inst., Heidelberg, Germany
SOURCE: Chemische Berichte (1953), 86, 1295-1302
CODEN: CHBEAM; ISSN: 0009-2940
DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

AB Some phenazine derivs. are prepared to be tested for their tuberculostatic action. Heating slowly 13 g. o-H₂NCO₂H₄CO₂H, 16 g. o-O₂NCO₂H₄CO₂H (I), and 25 g. finely powdered KOH to 85°, dissolving the melt in H₂O, and concentrating the solution give di-K 1,6-phenazinedicarboxylate (II) (C.A. numbering), which on acidification gives 1.5 g. free acid, charring at 300-30° without melting (di-Et ester, prepared by dissolving 1.5 g. II in 20 cc. 100% H₂SO₄, pouring the solution into absolute EtOH, neutralizing the mixture with NaOH to pH 8-9, and extracting with ether, green-yellow needles,


m. 143°). Treating 0.6 g. Me 1-phenazinecarboxylate with H₂NOH from 2 g. HCl salt] gives 1-phenazinehydroxamic acid, yellow needles, m. 207°. Adding 3.5 g. 1-phenazinecarboxamide to 240 cc. H₂O containing 9 g. NaOH and 2.5 g. Br and heating the mixture 5 min. at 70° give 74% 1-aminophenazine, red needles, m. 176°, which (0.5 g.), refluxed 2.5 hrs. with 2 g. anhydrous glucose and 20 mg. NH₄Cl in 30 cc. absolute MeOH, gives 50% N-D-glucoside, vermilion needles, m. 195° (tetra-Ac derivative, prepared by heating 0.4 g. glucoside in 10 cc. C₂H₅SH and 5 cc. Ac₂O 0.5 hr. on a water bath, orange needles, m. 181°). Heating 5 g. 1,4-g. 2-ClOH₂NH₂ and 15 g. KOH to 80°, raising the temperature 5° every 20 min. until 130° is reached, keeping the mixture 3 hrs. at 130°, dissolving the cold melt in H₂O, concentrating the filtered solution, and adding a little MeOH give the K benzo[α]phenazine-11-carboxylate (free acid, yellow needles, m. 256°). Mixing 2 g. 1,2-naphthoquinone (III) and 2.4 g. 3,4-(H₂N)2C₆H₃CO₂Et, each in 20 cc. AcOH, gives 80% Et benzo[α]phenazine-9(or 10)-carboxylate (IV), yellow needles, m. 205°, which, saponified with 30% aqueous KOH and acidified with AcOH, gives the free acid, yellow needles, m. 366°. Heating 0.5 g. IV and 6 cc. 40% NH₄H₂O in 15 cc. dioxane and 5 cc. EtOH 2 hrs. on a water bath gives the hydrazide, yellow needles, darkening at 270°, charring at 320°. Heating 0.52 g. III and 0.6 g. 3,4-(H₂N)2C₆H₃CH₂CH₂CO₂H in 10 cc. AcOH 5 min. on a water bath gives benzo[α]phenazine-9(or 10)-propionic acid, yellow needles, m. 212°, soluble in concentrated H₂SO₄ with a violet color. Heating 1.5 g. 3,4-(H₂N)2C₆H₃CO₂Et and 2.5 g. [CO₂]2 2H₂SO₃ in 15 cc. H₂O 1 hr. on a water bath gives 54% Et 6-quinoxalinecarboxylate, fine needles, m. 66° (free acid, needles, m. 266°). Refluxing 1 g. Et 2,3-diphenyl-6-quinoxalinecarboxylate in 30 cc. EtOH 5 hrs. with 15 cc. 40% NH₄H₂O gives the hydrazide, fine needles, m. 235°. 1-phenazinecarboxylic acid hydrazide, prepared from the Me ester, silky yellow needles, m. 231°, when refluxed (5.5 g.) in 20 cc. C₂H₅SH with 6 cc. PhSO₂Cl 3 hrs., gives 80% phenazine-1-carboxylic acid benzenesulfonylhydrazide, lacelike pale yellow needles, m. 233°. The results of the bacterio-static tests of these compds. are given in a table.

IT 6924-72-7, 6-Quinoxalinecarboxylic acid, ethyl ester (preparation of)
RN 6924-72-7 CAPLUS
CN 6-Quinoxalinecarboxylic acid, ethyl ester (6CI, 7CI, 8CI, 9CI) (CA INDEX NAME)




L13 ANSWER 180 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1954:64341 CAPLUS
DOCUMENT NUMBER: 48:64341
ORIGINAL REFERENCE NO.: 48:11426h-1,11427a-b

TITLE: Derivatives of 2-phenazinecarboxylic acid
AUTHOR(S): Pietra, Silvio; Maffei, Silvio; Rivotta, Angelamaria
CORPORATE SOURCE: Univ. Pavia, Italy
SOURCE: Annali di Chimica (Rome, Italy) (1953), 43, 227-31
CODEN: ANCRAL; ISSN: 0003-4592
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable
AB The synthesis of 2-phenazinecarboxylic acid (I) has already been described [cf. ibid. 42, 519(1952)]. Various deriva. are now described. Et2NH and MeOH with the acyl chloride of I yield 2-(diethylcarbamoyl)phenazine (II), m. 97.5-98°, and Me 2-phenazinecarboxylate (III), m. 153°. N2H4 and II yield 2-phenazinecarboxylic acid hydrazide (IV), m. 269-70° (decomposition). IV and o-HOCH4CHO 30 min. at 120°, give salicylaldehyde 2-phenazinecarbonylhydrazone, yellow needles, m. 275°. IV and PhCOMe give acetophenone 2-phenazinecarbonylhydrazone m. 259° (decomposition). IV (15 g.) in 900 cc. HCl (1:2) treated with 9 g. NaNO2 in 50 cc. H2O gives 2-phenazinecarboxylic acid azide (V), m. 135° (violent decomposition). V (2.49 g.) with 150 cc. absolute EtOH at 60-70° yields Et 2-phenazinecarbamate (VI), m. 196° [from ligroine (b. 90-130°)]. Likewise, Me2CHOH V yield the isopropyl urethan, m. 156°. V (0.997 g.) in 100 cc. xylene, decomposed with dry NH3 and heated to 120°, give 2-phenazinylurea, m. 261° (decomposition). VI (2.47 g.) and 50 cc. H2SO4 heated at 150-60°, cooled, diluted, and made alkaline with NH3 yield 2-aminophenazine (VII), m. 290-1° (from xylene), obtained also from V by heating in xylene with HCl.
IT 37648-82-1. 2-Phenazinecarboxamide, N,N-diethyl- (preparation of)
RN 37648-82-1 CAPLUS
CN 2-Phenazinecarboxamide, N,N-diethyl- (9CI) (CA INDEX NAME)




L13 ANSWER 181 OF 181 CAPLUS COPYRIGHT 2006 ACS ON STN
ACCESSION NUMBER: 1954:4690 CAPLUS
DOCUMENT NUMBER: 48:4690
ORIGINAL REFERENCE NO.: 48:884c-f
TITLE: The therapy of experimental peitacosis and lymphogranuloma venereum (inguinale). II. The activity of quinoxaline 1,4-dioxides and substituted and related compounds, with a note on the morphological changes induced in lymphogranuloma virus by these compounds and by antibiotics
AUTHOR(S): Hurst, E. Weston; Landquist, J. K.; Melvin, P.; Peters, J. M.; Senior, M.; Silk, J. A.; Stacey, G. J.
CORPORATE SOURCE: Imperial Chem. Inds., Ltd., Manchester, UK
SOURCE: British Journal of Pharmacology and Chemotherapy (1953), 5, 297-305
CODEN: BJPCAL; ISSN: 0366-0826
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable
AB cf. C.A. 45, 3084h. Representative mono- and disubstituted quinoxaline 1,4-dioxides, substituted 2,3-dimethylquinoxaline 1,4-dioxides, phenazine di-N-oxides, biquinoxaline tetra-N-oxides, and N-oxides of miscellaneous N-heterocyclic compds. possessed some degree of activity against the largest viruses of the peitacosis-lymphogranuloma group. Quinoxaline 1,4-dioxide (I) and its substituted deriva. were most potent, the best

equaling Aureomycin in their activity against lymphogranuloma venereum in the mouse. Relatively few were active against this disease in the chick embryo or peitacosis in the mouse. Therapeutic activity in man was noted but toxic side reactions preclude their use. These compds. did not inactivate virus in vitro but greatly restricted its growth in the mouse and altered its morphological appearance in the chick embryo. Therapeutic activity was not abolished by simultaneous administration of vitamin K. I deriva. did not influence infections with the viruses of herpes febrilis, ectromelia, mouse-adapted poliomyelitis, influenza A, equine encephalomyelitis, or louping-ill.
IT 108239-47-0. 6-Quinoxalinecarboxylic acid, 2,3-dimethyl-, ethyl ester, 1,4-dioxide (effect on peitacosis-lymphogranuloma group viruses)
RN 108239-47-0 CAPLUS
CN 6-Quinoxalinecarboxylic acid, 2,3-dimethyl-, ethyl ester, 1,4-dioxide (6CI) (CA INDEX NAME)



IT 109939-89-1. 6-Quinoxalinecarboxamide, 2,3-dimethyl-N-(3-piperidinopropyl)-, 1,4-dioxide (effect on peitacosis-lymphogranuloma group viruses)
RN 109939-89-1 CAPLUS
CN 6-Quinoxalinecarboxamide, 2,3-dimethyl-N-(3-piperidinopropyl)-, 1,4-dioxide (6CI) (CA INDEX NAME)



=> LOGOFF
ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF
LOGOFF? (Y/N/HOLD:Y)

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
FULL ESTIMATED COST	ENTRY	SESSION
	926.29	1305.50
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
CA SUBSCRIBER PRICE	ENTRY	SESSION
	-135.75	-135.75

STN INTERNATIONAL LOGOFF AT 12:23:58 ON 10 MAY 2006